

## Nutrient composition and feeding value of *Salicornia bigelovii* torr meal in broiler diets

F.M. Attia, A.A. Alsobayel, M.S. Kriadees, M.Y. Al-Saiady,  
M.S. Bayoumi

*Department of Animal Production, College of Agriculture, King Saud University, P.O. Box 2460, Riyadh 11451, Saudi Arabia*

Accepted 13 June 1996

---

### Abstract

Salicornia meal (SM) is a by-product of the oil extraction of the seeds of *Salicornia bigelovii* Torr, an oil-seed crop tolerant to seawater irrigation. Chemical analysis showed that it contained  $340 \text{ g kg}^{-1}$  of crude protein,  $64.5 \text{ g kg}^{-1}$  of ether extract and  $36 \text{ g kg}^{-1}$  of crude fiber with a gross energy of  $19.4 \text{ MJ kg}^{-1}$ . Salicornia is rich in linoleic acid. Lysine and sulfur containing amino acids are the most limiting in SM. In comparison with soybean meal, SM is inferior with respect to the amino acid profile. SM contains high levels of sodium ( $30.4 \text{ g kg}^{-1}$ ), chlorine ( $45.8 \text{ g kg}^{-1}$ ), and ash ( $150 \text{ g kg}^{-1}$ ). A feeding trial was conducted using broiler chickens to evaluate the nutritional value of SM. SM was incorporated in the diet at levels of 0, 30, 60 and  $90 \text{ g kg}^{-1}$ . The  $60 \text{ g kg}^{-1}$  SM diet was fed either with or without cholesterol ( $5 \text{ g kg}^{-1}$ ) whereas the  $90 \text{ g kg}^{-1}$  diet was fed only with cholesterol ( $5 \text{ g kg}^{-1}$ ). SM in broiler diets caused a depression ( $P \leq 0.01$ ) in growth and feed intake proportional to the level of SM in the diet. However, the growth depressing activity of SM was counteracted by the inclusion of cholesterol. These results indicate that SM may be used as an unconventional feed ingredient in broiler diets when supplemented with cholesterol. © 1997 Elsevier Science B.V.

*Keywords:* Salicornia meal; Chemical analysis; Antigrowth factor; Cholesterol; Broiler performance

---