

Growth Response of Cucumber to Hydrophilic Polymer Application Under Different Soil Moisture Levels

A. R. Al-Harbi
A. M. Al-Omran
A. Shalaby
H. Wahdan
M. I. Choudhary

ABSTRACT. The effect of hydrophilic polymer addition on the growth of cucumber (*Cucumis sativus* L., cv. Dina) was investigated under different levels of soil moisture. The plants were grown in pots filled with 7 kg of moderately calcareous loam soil. The soil was mixed with five concentrations of hydrophilic polymer Broad Leaf (P-4). These concentrations were 0.0% (control), 0.1, 0.2, 0.3, and 0.4% on a dry-weight basis. Three levels of soil moisture, namely, 25, 50, and 75% of field capacity of untreated soil (control), were used in this study. The results showed a significant increase in the vegetative growth with increasing polymer concentration up to 0.3%. Further increase to 0.4% of P-4 had no significant effect on plant growth. A significant interaction was observed between polymer concentration and soil moisture level. Growing plants under 50% of field capacity with 0.2% polymer concentration resulted in more vegetative growth than did other treatments. Addition of hydrophilic polymers was most effective when the plants were grown

A. R. Al-Harbi is affiliated with the Department of Plant Production, A. M. Al-Omran and A. Shalaby are affiliated with the Department of Soil Science, H. Wahdan is affiliated with the Department of Plant Production, and M. I. Choudhary is affiliated with the Department of Soil Science, College of Agriculture, King Saud University, P.O. Box 2460, Riyadh 11451, Kingdom of Saudi Arabia.

The financial support of the Agricultural Research Center, College of Agriculture, King Saud University is gratefully acknowledged.