

## Isometry of Kaehlerian manifolds to complex projective spaces

By Kentaro YANO and Hitosi HIRAMATU

(Received March 24, 1979)

### §1. Introduction.

Let  $M$  be a complex  $n$ -dimensional connected Kaehlerian manifold covered by a system of real coordinate neighborhoods  $\{U; x^h\}$ , where, here and in the sequel, the indices  $h, i, j, k, \dots$  run over the range  $\{1, 2, \dots, 2n\}$  and let  $g_{ji}$ ,  $F_i^h$ ,  $\{j^h_i\}$ ,  $\nabla_i$ ,  $K_{kji}^h$ ,  $K_{ji}$  and  $K$  be respectively the Hermitian metric tensor, the complex structure tensor, the Christoffel symbols formed with  $g_{ji}$ , the operator of covariant differentiation with respect to  $\{j^h_i\}$ , the curvature tensor, the Ricci tensor and the scalar curvature of  $M$ .

A vector field  $v^h$  is called a *holomorphically projective* (or *H-projective*, for brevity) vector field [2, 3, 5, 7] if it satisfies

$$(1.1) \quad \begin{aligned} L_v\{j^h_i\} &= \nabla_j \nabla_i v^h + v^k K_{kji}^h \\ &= \delta_j^h \rho_i + \delta_i^h \rho_j - \rho_i F_j^t F_i^h - \rho_t F_i^t F_j^h \end{aligned}$$

for a certain covariant vector field  $\rho_i$  on  $M$ , called the associated covariant vector field of  $v^h$ , where  $L_v$  denotes the operator of Lie derivation with respect to  $v^h$ . In particular, if  $\rho_i$  is zero vector field then  $v^h$  is called an affine vector field. When we refer in the sequel to an *H-projective* vector field  $v^h$ , we always mean by  $\rho_i$  the associated covariant vector field appearing in (1.1).

Recently, the present authors [9, 10] and one of the present authors [1] proved a series of integral inequalities in a compact Kaehlerian manifold with constant scalar curvature admitting an *H-projective* vector field and then obtained necessary and sufficient conditions for such a Kaehlerian manifold to be isometric to a complex projective space with Fubini-Study metric.

The purpose of the present paper is to continue the joint work [9, 10] of the present authors and to prove the following theorem.

**THEOREM A.** *If a complex  $n > 1$  dimensional, compact, connected and simply connected Kaehlerian manifold  $M$  with constant scalar curvature  $K$  admits a non-affine H-projective vector field  $v^h$ , then  $M$  is isometric to a complex projective space  $CP^n$  with Fubini-Study metric and of constant holomorphic sectional cur-*

vature  $\frac{K}{n(n+1)}$ .

In the sequel, we need the following theorem due to Obata [4]. (See also [6].)

**THEOREM B.** *Let  $M$  be a complete, connected and simply connected Kaehlerian manifold. In order for  $M$  to admit a non-trivial solution  $\varphi$  of a system of partial differential equations*

$$(1.2) \quad \nabla_j \nabla_i \varphi_n + \frac{c}{4} (2\varphi_j g_{ih} + \varphi_i g_{jh} + \varphi_n g_{ji} - F_{ji} F_n^t \varphi_t - F_{jh} F_i^t \varphi_t) = 0,$$

where  $\varphi_n = \nabla_n \varphi$  and  $F_{ji} = F_j^t g_{ti}$ ,  $c$  being a positive constant, it is necessary and sufficient that  $M$  is isometric to a complex projective space  $CP^n$  with Fubini-Study metric and of constant holomorphic sectional curvature  $c$ .

We assume in this paper that Kaehlerian manifolds under consideration are connected.

## § 2. Preliminaries.

Let  $M$  be a complex  $n$ -dimensional Kaehlerian manifold. The complex structure tensor  $F_i^h$  and the Hermitian metric tensor  $g_{ji}$  of  $M$  satisfy

$$(2.1) \quad F_i^h F_i^t = -\delta_i^h, \quad \nabla_j F_i^h = 0, \quad \nabla_j F_{ih} = 0,$$

$$(2.2) \quad g_{jt} F_i^t + g_{ti} F_j^t = 0$$

and

$$(2.3) \quad g_{jt} - g_{ts} F_j^t F_i^s = 0.$$

We have [7, 9, 10], for the curvature tensor  $K_{kji}^h$ ,

$$(2.4) \quad K_{kji}^t F_t^h - K_{kjt}^h F_i^t = 0,$$

$$(2.5) \quad K_{kji}^h + K_{kjt}^s F_i^t F_s^h = 0,$$

$$(2.6) \quad K_{kjit} F_n^t + K_{kjt} F_i^t = 0$$

and

$$(2.7) \quad K_{kjih} - K_{kjt} F_i^t F_n^s = 0,$$

where  $K_{kjih} = K_{kjt}^t g_{tn}$ .

Using (2.6) and the first Bianchi identity

$$K_{kjih} + K_{ikjh} + K_{jikh} = 0,$$

we have

$$\begin{aligned} 2K_{kt}F_j^t &= 2g^{ut}K_{kuts}F_j^s = -2g^{ut}K_{kusj}F_t^s \\ &= -2K_{ktsj}F^{ts} = -(K_{ktsj} - K_{kstj})F^{ts} \\ &= K_{tskj}F^{ts} = K_{kjts}F^{ts}, \end{aligned}$$

from which

$$(2.8) \quad K_{kjts}F^{ts} = 2K_{kt}F_j^t,$$

$$(2.9) \quad K_{tjis}F^{ts} = -K_{jt}F_i^t$$

and

$$(2.10) \quad K_{ktsj}F^{ts} = -K_{kt}F_h^t,$$

$g^{ji}$  being contravariant components of  $g_{ji}$  and  $F^{kj} = g^{kt}F_t^j$ .

From (2.8), we have, for the Ricci tensor  $K_{ji} = K_{tji}^t$ ,

$$(2.11) \quad K_{jt}F_i^t + K_{ti}F_j^t = 0,$$

$$(2.12) \quad K_{ji} - K_{ts}F_j^tF_i^s = 0,$$

$$(2.13) \quad K_t^hF_i^t - K_i^tF_t^h = 0$$

and

$$(2.14) \quad K_i^h + K_t^sF_i^tF_s^h = 0,$$

where  $K_i^h = K_{it}g^{th}$ .

A Kaehlerian manifold  $M$  has the constant holomorphic sectional curvature  $k$  if and only if

$$(2.15) \quad K_{kji}^h = \frac{k}{4}(\delta_k^h g_{ji} - \delta_j^h g_{ki} + F_k^h F_{ji} - F_j^h F_{ki} - 2F_{kj}F_i^h).$$

We define tensor fields  $G_{ji}$  and  $Z_{kji}^h$  [9, 10] on  $M$  by

$$(2.16) \quad G_{ji} = K_{ji} - \frac{K}{2n}g_{ji}$$

and

$$(2.17) \quad Z_{kji}^h = K_{kji}^h - \frac{K}{4n(n+1)}(\delta_k^h g_{ji} - \delta_j^h g_{ki} + F_k^h F_{ji} - F_j^h F_{ki} - 2F_{kj}F_i^h)$$

respectively. If  $G_{ji} = 0$  for  $n > 1$  then  $M$  is a Kaehler-Einstein manifold and  $K$  is a constant and if  $Z_{kji}^h = 0$  for  $n > 1$  then  $M$  is of constant holomorphic sectional curvature  $\frac{K}{n(n+1)}$ .

We easily see that the tensor fields  $G_{ji}$  and  $Z_{kji}^h$  satisfy

$$(2.18) \quad G_{ji} = G_{ij}, \quad G_{ji}g^{ji} = 0, \quad Z_{tji}^t = G_{ji},$$

$$(2.19) \quad Z_{kjt}{}^h = -Z_{jkt}{}^h, \quad Z_{kji}{}^h = Z_{ihkj}$$

and

$$(2.20) \quad Z_{kji}{}^h + Z_{ikj}{}^h + Z_{jtk}{}^h = 0,$$

where  $Z_{kji}{}^h = Z_{kji}{}^t g_{th}$ .

The tensor fields  $G_{ji}$  and  $Z_{kji}{}^h$  also satisfy

$$(2.21) \quad G_{jt}F_i{}^t + G_{ti}F_j{}^t = 0,$$

$$(2.22) \quad G_{ji} - G_{ts}F_j{}^t F_i{}^s = 0,$$

$$(2.23) \quad G_t{}^h F_i{}^t - G_i{}^t F_t{}^h = 0,$$

$$(2.24) \quad G_i{}^h + G_t{}^s F_i{}^t F_s{}^h = 0,$$

$$(2.25) \quad Z_{kji}{}^t F_t{}^h - Z_{kjt}{}^h F_i{}^t = 0,$$

$$(2.26) \quad Z_{kji}{}^h + Z_{kjt}{}^s F_i{}^t F_s{}^h = 0,$$

$$(2.27) \quad Z_{kjit}F_h{}^t + Z_{kjtih}F_i{}^t = 0,$$

$$(2.28) \quad Z_{kji}{}^h - Z_{kjt}{}^s F_i{}^t F_h{}^s = 0,$$

$$(2.29) \quad Z_{kjt}{}^s F^{ts} = 2G_{kt}F_j{}^t,$$

$$(2.30) \quad Z_{tjis}F^{ts} = -G_{jt}F_i{}^t$$

and

$$(2.31) \quad Z_{kts}{}^h F^{ts} = -G_{kt}F_h{}^t,$$

where  $G_i{}^h = G_{it}g^{th}$ .

If the scalar curvature  $K$  is a constant, then, from (2.1), (2.17) and the second Bianchi identity

$$\nabla_l K_{kji}{}^h + \nabla_j K_{lki}{}^h + \nabla_k K_{jli}{}^h = 0,$$

we have

$$(2.32) \quad \nabla_l Z_{kji}{}^h + \nabla_j Z_{lki}{}^h + \nabla_k Z_{jli}{}^h = 0,$$

from which and (2.18) and (2.19),

$$(2.33) \quad \nabla_t Z_{kji}{}^t = \nabla_k G_{ji} - \nabla_j G_{ki}.$$

A vector field  $u^h$  on  $M$  is said to be *contravariant analytic* if

$$(2.34) \quad (\nabla_j u_i)F_i{}^t + (\nabla_t u_i)F_j{}^t = 0$$

or equivalently if

$$(2.35) \quad \nabla_j u_i - (\nabla_t u_s)F_j{}^t F_i{}^s = 0,$$

where  $u_i = g_{ih}u^h$ . A vector field  $u^h$  on  $M$  is contravariant analytic if and only if

$$(2.36) \quad L_u F_i^h = 0,$$

where  $L_u$  denotes the operator of Lie derivation with respect to  $u^h$ . It is known [7] that if  $M$  is compact then a necessary and sufficient condition for a vector field  $u^h$  on  $M$  to be contravariant analytic is that

$$(2.37) \quad \nabla^i \nabla_i u^h + K_i^h u^i = 0$$

holds, where  $\nabla^i = g^{ih} \nabla_h$ .

For an  $H$ -projective vector field  $v^h$  on  $M$  defined by (1.1), we have

$$(2.38) \quad \nabla_j \nabla_i v^t = 2(n+1) \rho_j$$

and

$$(2.39) \quad \nabla^i \nabla_i v^h + K_i^h v^i = 0.$$

(2.38) shows that the associated covariant vector field  $\rho_j$  is gradient. Putting

$$(2.40) \quad \rho = \frac{1}{2(n+1)} \nabla_t v^t,$$

we have

$$(2.41) \quad \rho_j = \nabla_j \rho.$$

We have, for the  $H$ -projective vector field  $v^h$ , from (1.1),

$$(2.42) \quad \nabla_j L_v g_{ih} = 2\rho_j g_{ih} + \rho_i g_{jh} + \rho_h g_{ji} - F_{ji} F_h^t \rho_t - F_{jh} F_i^t \rho_t,$$

from which

$$(2.43) \quad \nabla^j L_v g_{ih} = 2\rho^j g_{ih} + \rho_i \delta_h^j + \rho_h \delta_i^j + F_i^j F_h^t \rho_t + F_h^j F_i^t \rho_t,$$

$$(2.44) \quad \nabla_j L_v g^{ih} = -2\rho_j g^{ih} - \rho^i \delta_j^h - \rho^h \delta_j^i + F_j^i F^{ht} \rho_t + F_j^h F^{it} \rho_t$$

and

$$(2.45) \quad \nabla^j L_v g^{ih} = -2\rho^j g^{ih} - \rho^i g^{jh} - \rho^h g^{ji} + F^{ji} F^{ht} \rho_t + F^{jh} F^{it} \rho_t,$$

where  $\rho^h = \rho_i g^{ih}$ .

Substituting (1.1) into the well known formula [7, 8]

$$L_v K_{kji}^h = \nabla_k L_v \{j^h i\} - \nabla_j L_v \{k^h i\},$$

we find

$$(2.46) \quad L_v K_{kji}^h = -\delta_k^h \nabla_j \rho_i + \delta_j^h \nabla_k \rho_i \\ + (F_k^h \nabla_j \rho_i - F_j^h \nabla_k \rho_i) F_i^t + (F_k^t \nabla_j \rho_t - F_j^t \nabla_k \rho_t) F_i^h,$$

from which, contracting with respect to  $h$  and  $k$ ,

$$(2.47) \quad L_v K_{ji} = -2n \nabla_j \rho_i - 2(\nabla_i \rho_s) F_j^s F_i^s.$$

Suppose that an  $H$ -projective vector field  $v^h$  on  $M$  is contravariant analytic. Then, applying the operator  $L_v$  of Lie derivation with respect to  $v^h$  to both sides of (2.12), we have

$$L_v K_{ji} = (L_v K_{is}) F_j^s F_i^s,$$

from which and (2.47), we see that if  $n > 1$  then  $\rho^h$  is also contravariant analytic and

$$(2.48) \quad L_v K_{ji} = -2(n+1) \nabla_j \rho_i$$

holds.

If a Kaehlerian manifold  $M$  is compact, then we see, from (2.39), that an  $H$ -projective vector field  $v^h$  on  $M$  is contravariant analytic or equivalently  $L_v F_i^h = 0$  holds and moreover if  $n > 1$  then the associated vector field  $\rho^h$  is also contravariant analytic and (2.48) holds.

For a contravariant analytic  $H$ -projective vector field  $v^h$  on a complex  $n > 1$  dimensional Kaehlerian manifold  $M$  with constant scalar curvature  $K$ , we have [9, 10], for the tensor field  $G_{ji}$ ,

$$(2.49) \quad L_v G_{ji} = -\nabla_j w_i - \nabla_i w_j,$$

where we have put

$$(2.50) \quad w^h = (n+1) \rho^h + \frac{K}{2n} v^h$$

and  $w_i = g_{ih} w^h$ , and, for the tensor field  $Z_{kji}^h$ ,

$$(2.51) \quad L_v Z_{kji}^h = \frac{1}{2(n+1)} \{ \delta_k^h L_v G_{ji} - \delta_j^h L_v G_{ki} - F_k^h (L_v G_{jt}) F_i^t + F_j^h (L_v G_{kt}) F_i^t - F_k^t (L_v G_{jt}) F_i^h + F_j^t (L_v G_{kt}) F_i^h \}.$$

### § 3. Proof of Theorem A.

In this section, we prove Theorem A. For this purpose, we need a series of lemmas. We use freely formulas (2.1)~(2.51) in the proofs of all lemmas and Theorem A in this section.

LEMMA 1 (Yano and Hiramatu [9]). *If, in a compact Kaehlerian manifold  $M$ , a non-constant function  $\varphi$  satisfies*

$$(3.1) \quad \nabla_j \nabla_i \varphi_n + \frac{c}{4} (2\varphi_j g_{ih} + \varphi_i g_{jh} + \varphi_n g_{ji} - F_{jt} F_h^t \varphi_t - F_{jh} F_i^t \varphi_t) = 0,$$

where  $\varphi_n = \nabla_n \varphi$ ,  $c$  being a constant, then the constant  $c$  is necessarily positive.

PROOF. Transvecting (3.1) with  $g^{in}$ , we have

$$\nabla_j \Delta \varphi + (n+1)c\varphi_j = 0,$$

from which

$$c \int_M \varphi_j \varphi^j dV = -\frac{1}{n+1} \int_M (\nabla_j \Delta \varphi) \varphi^j dV = \frac{1}{n+1} \int_M (\Delta \varphi)^2 dV,$$

where  $\Delta = g^{j\bar{i}} \nabla_j \nabla_{\bar{i}}$ ,  $\varphi^j = g^{j\bar{i}} \varphi_{\bar{i}}$  and  $dV$  denotes the volume element of  $M$ . Since  $\varphi$  is a non-constant function, two inequalities

$$\int_M \varphi_j \varphi^j dV > 0, \quad \int_M (\Delta \varphi)^2 dV > 0$$

hold and consequently  $c$  is necessarily positive.

LEMMA 2 (Yano and Hiramatu [9]). *If a complete and simply connected Kaehlerian manifold  $M$  with positive constant scalar curvature  $K$  admits a non-affine  $H$ -projective vector field  $v^h$  and if the vector field  $w^h$  defined by (2.50) is a Killing vector field, then  $M$  is isometric to a complex projective space  $CP^n$  with Fubini-Study metric of constant holomorphic sectional curvature  $\frac{K}{n(n+1)}$ .*

PROOF. We have, from (1.1),

$$(3.2) \quad \nabla_j (\nabla_i v_h + \nabla_h v_i) = 2\rho_j g_{ih} + \rho_i g_{jh} + \rho_h g_{ji} - F_{ji} F_h^t \rho_t - F_{jh} F_i^t \rho_t.$$

If  $w^h$  is a Killing vector field then

$$\nabla_i w_h + \nabla_h w_i = 0$$

holds and consequently

$$2(n+1)\nabla_i \rho_h + \frac{K}{2n} (\nabla_i v_h + \nabla_h v_i) = 0,$$

from which and (3.2), we find

$$\nabla_j \nabla_i \rho_h + \frac{K}{4n(n+1)} (2\rho_j g_{ih} + \rho_i g_{jh} + \rho_h g_{ji} - F_{ji} F_h^t \rho_t - F_{jh} F_i^t \rho_t) = 0.$$

Thus the lemma follows from Theorem B.

REMARK. Using Lemma 1, we see that if, in Lemma 2,  $M$  is compact then we can remove the positiveness of the scalar curvature  $K$  from the assumption.

LEMMA 3 (Yano and Hiramatu [9]). *For an  $H$ -projective vector field  $v^h$  on a complex  $n > 1$  dimensional compact Kaehlerian manifold  $M$  with constant scalar curvature  $K$ , we have*

$$(3.3) \quad \int_M (\nabla_i w^t)^2 dV = \frac{1}{2} \int_M (\nabla_j w_i + \nabla_i w_j) (\nabla^j w^i + \nabla^i w^j) dV.$$

PROOF. By using a well known formula [7, 8] on a compact orientable Riemannian manifold, we have

$$\int_M (\nabla^i \nabla_i w^h + K_i^h w^i) w_h dV - \int_M (\nabla_i w^t)^2 dV + \frac{1}{2} \int_M (\nabla_j w_i + \nabla_i w_j) (\nabla^j w^i + \nabla^i w^j) dV = 0.$$

On the other hand, as was stated in Section 2, the associated vector field  $\rho^h$  is contravariant analytic and hence satisfies

$$\nabla^i \nabla_i \rho^h + K_i^h \rho^i = 0.$$

Consequently (3.3) follows from (2.39), (2.50) and the above relations since  $K$  is a constant.

LEMMA 4 (Yano and Hiramatu [9]). *For an  $H$ -projective vector field  $v^h$  on a complex  $n > 1$  dimensional compact Kaehlerian manifold  $M$  with constant scalar curvature  $K$ , we have*

$$(3.4) \quad \int_M G_{ji} \rho^j w^i dV = \frac{1}{4(n+1)} \int_M (\nabla_j w_i + \nabla_i w_j) (\nabla^j w^i + \nabla^i w^j) dV.$$

PROOF. The associated vector field  $\rho^h$  is contravariant analytic and hence satisfies

$$\nabla^j \nabla_j \rho^i + K_j^i \rho^j = 0,$$

from which and the equality

$$\nabla_i \nabla_i \rho^t = \nabla^t \nabla_i \rho_i - K_{ji} \rho^j$$

we find

$$\nabla_i \nabla_i \rho^t = -2K_{ji} \rho^j.$$

Using the above relation, (2.40), (2.41), (2.50) and Lemma 3, we have

$$\begin{aligned} \int_M G_{ji} \rho^j w^i dV &= \int_M K_{ji} \rho^j w^i dV - \frac{K}{2n} \int_M \rho_i w^i dV \\ &= -\frac{1}{2} \int_M (\nabla_i \nabla_i \rho^t) w^t dV - \frac{K}{2n} \int_M \rho_i w^i dV \\ &= -\frac{1}{2} \int_M (\nabla_i \nabla_i \rho^t) w^t dV - \frac{K}{4n(n+1)} \int_M (\nabla_i \nabla_i v^t) w^t dV \\ &= -\frac{1}{2(n+1)} \int_M \left[ \nabla_i \nabla_i \left\{ (n+1) \rho^t + \frac{K}{2n} v^t \right\} \right] w^t dV \\ &= -\frac{1}{2(n+1)} \int_M (\nabla_i \nabla_i w^t) w^t dV = \frac{1}{2(n+1)} \int_M (\nabla_i w^t)^2 dV \\ &= \frac{1}{4(n+1)} \int_M (\nabla_j w_i + \nabla_i w_j) (\nabla^j w^i + \nabla^i w^j) dV. \end{aligned}$$

LEMMA 5 (Yano and Hiramatu [9]). *For an  $H$ -projective vector field  $v^h$  on a complex  $n > 1$  dimensional compact Kaehlerian manifold  $M$  with constant scalar curvature  $K$ , we have*

$$(3.5) \quad \int_M (\nabla^j L_v G_{ji}) w^i dV = \frac{1}{2} \int_M (\nabla_j w_i + \nabla_i w_j) (\nabla^j w^i + \nabla^i w^j) dV.$$

PROOF. Integrating

$$\begin{aligned}\nabla^j \{(L_\nu G_{ji})w^i\} &= (\nabla^j L_\nu G_{ji})w^i + (L_\nu G_{ji})\nabla^j w^i \\ &= (\nabla^j L_\nu G_{ji})w^i + \frac{1}{2}(L_\nu G_{ji})(\nabla^j w^i + \nabla^i w^j)\end{aligned}$$

over  $M$  and using (2.49), we have (3.5).

LEMMA 6 (Yano and Hiramatu [9]). *For an  $H$ -projective vector field  $v^h$  on a complex  $n > 1$  dimensional compact Kaehlerian manifold  $M$  with constant scalar curvature  $K$ , we have*

$$(3.6) \quad \int_M g^{kj}(L_\nu \nabla_k G_{ji})w^i dV = \frac{n}{2(n+1)} \int_M (\nabla_j w_i + \nabla_i w_j)(\nabla^j w^i + \nabla^i w^j) dV.$$

PROOF. Substituting (1.1) into

$$g^{kj}(L_\nu \nabla_k G_{ji})w^i = (\nabla^j L_\nu G_{ji})w^i - g^{hj}(L_\nu \{^t_j\})G_{ti}w^i - g^{hj}(L_\nu \{^t_i\})G_{jt}w^i,$$

we have

$$g^{kj}(L_\nu \nabla_k G_{ji})w^i = (\nabla^j L_\nu G_{ji})w^i - 2G_{ji}\rho^j w^i.$$

Integrating this over  $M$  and using Lemmas 4 and 5, we have (3.6).

LEMMA 7 (Yano and Hiramatu [10]). *For an  $H$ -projective vector field  $v^h$  on a complex  $n > 1$  dimensional compact Kaehlerian manifold  $M$  with constant scalar curvature  $K$ , we have*

$$(3.7) \quad \int_M (\nabla^k L_\nu Z_{kji}{}^h)g^{ji}w_h dV = \frac{1}{n+1} \int_M (\nabla_j w_i + \nabla_i w_j)(\nabla^j w^i + \nabla^i w^j) dV.$$

PROOF. By using (2.51), we have

$$\begin{aligned}(\nabla^k L_\nu Z_{kji}{}^h)g^{ji}w_h &= \frac{1}{2(n+1)} \{(\nabla^k L_\nu G_{ji})g^{ji}w_k - (\nabla^j L_\nu G_{ji})w^i \\ &\quad - F_k{}^h(\nabla^k L_\nu G_{jt})F_i{}^t g^{ji}w_h + F_j{}^h(\nabla^k L_\nu G_{kt})F_i{}^t g^{ji}w_h \\ &\quad - F_k{}^t(\nabla^k L_\nu G_{jt})F_i{}^h g^{ji}w_h + F_j{}^t(\nabla^k L_\nu G_{kt})F_i{}^h g^{ji}w_h\}.\end{aligned}$$

Here we notice that

$$-F_k{}^h(\nabla^k L_\nu G_{jt})F_i{}^t g^{ji}w_h = -F_k{}^h(\nabla^k L_\nu G_{jt})F^{jt}w_h = 0,$$

$$F_j{}^h(\nabla^k L_\nu G_{kt})F_i{}^t g^{ji}w_h = (\nabla^j L_\nu G_{ji})w^i,$$

$$-F_k{}^t(\nabla^k L_\nu G_{jt})F_i{}^h g^{ji}w_h = F_j{}^t(\nabla^k L_\nu G_{tk})F_i{}^h g^{ji}w_h = (\nabla^j L_\nu G_{ji})w^i$$

and

$$F_j{}^t(\nabla^k L_\nu G_{kt})F_i{}^h g^{ji}w_h = (\nabla^j L_\nu G_{ji})w^i$$

hold. Therefore we have

$$(\nabla^k L_v Z_{kji}{}^h) g^{ji} w_h = \frac{1}{2(n+1)} \{ (\nabla^k L_v G_{ji}) g^{ji} w_k + 2(\nabla^j L_v G_{ji}) w^i \}.$$

Integrating this over  $M$  and using (2.49) and Lemmas 3 and 5, we find

$$\begin{aligned} \int_M (\nabla^k L_v Z_{kji}{}^h) g^{ji} w_h dV &= \frac{1}{2(n+1)} \left\{ \int_M (\nabla^k L_v G_{ji}) g^{ji} w_k dV + 2 \int_M (\nabla^j L_v G_{ji}) w^i dV \right\} \\ &= \frac{1}{2(n+1)} \left\{ - \int_M (L_v G_{ji}) g^{ji} \nabla_i w^t dV + 2 \int_M (\nabla^j L_v G_{ji}) w^i dV \right\} \\ &= \frac{1}{n+1} \left\{ \int_M (\nabla_i w^t)^2 dV + \int_M (\nabla^j L_v G_{ji}) w^i dV \right\} \\ &= \frac{1}{n+1} \int_M (\nabla_j w_i + \nabla_i w_j) (\nabla^j w^t + \nabla^t w^j) dV. \end{aligned}$$

LEMMA 8 (Yano and Hiramatu [10]). *For an  $H$ -projective vector field  $v^h$  on a complex  $n > 1$  dimensional compact Kaehlerian manifold  $M$  with constant scalar curvature  $K$ , we have*

$$(3.8) \quad \int_M (\nabla^k L_v Z_{kji}{}^h) g^{ji} w^h dV = \frac{3}{2(n+1)} \int_M (\nabla_j w_i + \nabla_i w_j) (\nabla^j w^t + \nabla^t w^j) dV.$$

PROOF. By using  $Z_{kji}{}^h g^{ji} = G_k{}^h$  and

$$(\nabla^k Z_{kji}{}^h) g^{ji} = \nabla^k G_k{}^h = \frac{n-1}{2n} \nabla^h K = 0,$$

we have

$$\begin{aligned} (\nabla^k L_v Z_{kji}{}^h) g^{ji} w^h &= \{ \nabla^k L_v (Z_{kji}{}^t g_{th}) \} g^{ji} w^h \\ &= (\nabla^k L_v Z_{kji}{}^h) g^{ji} w_h + G_k{}^t (\nabla^k L_v g_{th}) w^h. \end{aligned}$$

Substituting (2.43) into this, we find

$$(\nabla^k L_v Z_{kji}{}^h) g^{ji} w^h = (\nabla^k L_v Z_{kji}{}^h) g^{ji} w_h + 2G_{ji} \rho^j w^i,$$

and consequently, integrating this over  $M$  and using Lemmas 4 and 7, we have (3.8).

Now we prove Theorem A. Using

$$\nabla^k Z_{kji}{}^h = \nabla_k Z_{hi}{}^k$$

and

$$(L_v Z_{kji}{}^h) g^{ji} = L_v G_{kh} - Z_{kji}{}^h L_v g^{ji},$$

we have

$$(\nabla^k L_v Z_{kji}{}^h) g^{ji} w^h = (\nabla^k L_v G_{kh}) w^h - (\nabla_k Z_{hi}{}^k) (L_v g^{ji}) w^h - Z_{kji}{}^h (\nabla^k L_v g^{ji}) w^h.$$

Substituting (2.33) and (2.45) into this, we find

$$\begin{aligned} (\nabla^k L_v Z_{kji h}) g^{ji} w^h &= (\nabla^j L_v G_{ji}) w^i - (\nabla_h G_{ij})(L_v g^{ji}) w^h \\ &\quad + (\nabla_i G_{hj})(L_v g^{ji}) w^h + 4G_{ji} \rho^j w^i, \end{aligned}$$

from which, integrating over  $M$ ,

$$\begin{aligned} \int_M (\nabla^k L_v Z_{kji h}) g^{ji} w^h dV &= \int_M (\nabla^j L_v G_{ji}) w^i dV - \int_M (\nabla_h G_{ij})(L_v g^{ji}) w^h dV \\ &\quad + \int_M (\nabla_j G_{ih})(L_v g^{ji}) w^h dV + 4 \int_M G_{ji} \rho^j w^i dV. \end{aligned}$$

Here we notice that we have, using (2.44) and (2.49),

$$\begin{aligned} - \int_M (\nabla_h G_{ji})(L_v g^{ji}) w^h dV &= \int_M G_{ji} (\nabla_h L_v g^{ji}) w^h dV + \int_M G_{ji} (L_v g^{ji}) \nabla_i w^i dV \\ &= -4 \int_M G_{ji} \rho^j w^i dV - \int_M (L_v G_{ji}) g^{ji} \nabla_i w^i dV \\ &= -4 \int_M G_{ji} \rho^j w^i dV + 2 \int_M (\nabla_i w^i)^2 dV \end{aligned}$$

and, using

$$(\nabla_j G_{ih}) g^{ji} = \frac{n-1}{2n} \nabla_h K = 0,$$

$$\int_M (\nabla_j G_{ih})(L_v g^{ji}) w^h dV = - \int_M g^{ji} (L_v \nabla_j G_{ih}) w^h dV.$$

Consequently, we have

$$\begin{aligned} \int_M (\nabla^k L_v Z_{kji h}) g^{ji} w^h dV &= 2 \int_M (\nabla_i w^i)^2 dV + \int_M (\nabla^j L_v G_{ji}) w^i dV - \int_M g^{kj} (L_v \nabla_k G_{ji}) w^i dV, \end{aligned}$$

from which, using Lemmas 3, 5 and 6,

$$\int_M (\nabla^k L_v Z_{kji h}) g^{ji} w^h dV = \frac{2n+3}{2(n+1)} \int_M (\nabla_j w_i + \nabla_i w_j) (\nabla^j w^i + \nabla^i w^j) dV.$$

From this and Lemma 8, we have

$$\int_M (\nabla_j w_i + \nabla_i w_j) (\nabla^j w^i + \nabla^i w^j) dV = 0,$$

from which

$$\nabla_j w_i + \nabla_i w_j = 0,$$

that is, the vector field  $w^h$  is a Killing vector field. Thus, Theorem A follows from Lemma 2.

### Bibliography

- [ 1 ] H. Hiramatu, Integral inequalities in Kaehlerian manifolds and their applications, to appear in *Period. Math. Hungar.*
- [ 2 ] S. Ishihara, Holomorphically projective changes and their groups in an almost complex manifold, *Tôhoku Math. J.*, **9** (1959), 273-297.
- [ 3 ] S. Ishihara, On holomorphic planes, *Ann. Mat. Pura Appl.*, **47** (1959), 197-241.
- [ 4 ] M. Obata, Riemannian manifolds admitting a solution of a certain system of differential equations, *Proc. U.S.-Japan Seminar in Differential Geometry*, Kyoto, Japan, 1965, 101-114.
- [ 5 ] S. Tachibana and S. Ishihara, On infinitesimal holomorphically projective transformations in Kählerian manifolds, *Tôhoku Math. J.*, **12** (1960), 77-101.
- [ 6 ] S. Tanno, Some differential equations on Riemannian manifolds, *J. Math. Soc. Japan*, **30** (1978), 509-531.
- [ 7 ] K. Yano, *Differential geometry on complex and almost complex spaces*, Pergamon Press, 1965.
- [ 8 ] K. Yano, *Integral formulas in Riemannian geometry*, Marcel Dekker, New York, 1970.
- [ 9 ] K. Yano and H. Hiramatu, On Kaehlerian manifolds with constant scalar curvature admitting a holomorphically projective vector field, to appear in *J. Differential Geometry*.
- [10] K. Yano and H. Hiramatu, Integral inequalities and their applications in Kaehlerian manifolds admitting a holomorphically projective vector field, *Bull. Inst. Math. Acad. Sinica*, **6** (1978), 313-332.

Kentaro YANO  
Department of Mathematics  
Tokyo Institute of Technology  
Tokyo 152  
Japan

Hitosi HIRAMATU  
Department of Mathematics  
Faculty of Engineering  
Kumamoto University  
Kumamoto 860  
Japan