

*Case Report***ANESTHETIC CONSIDERATIONS OF GASTRO-BRONCHIAL FISTULA REPAIR**

ABDELAZEEM EL-DAWLATLY<sup>1</sup> M.D., KHALID ALKATTAN<sup>2</sup> F.R.C.S.,  
WASEEM HAJJAR<sup>3</sup> F.R.C.S. AND MOHAMED MAHDY<sup>4</sup> M.D.

Gastro-bronchial fistula (GBF) represents an extremely rare complication after surgical procedures. Anesthetic management of such case is challenging. We are presenting a case of GBF following gastric bypass surgery for treating morbid obesity. A 25-year-old female patient was admitted in the thoracic surgical floor with cough of gastric contents following Roux-in-Y surgery. Barium meal revealed gastro-bronchial fistulous tract. She underwent left thoracotomy under general anesthesia and the fistula was closed. Rapid sequence induction of anesthesia and proper isolation of the sound lung are important to minimize incidence of pulmonary aspiration and soiling of the non-operated lung at induction of anesthesia. All precautions of managing one lung ventilation during surgery are to be undertaken. In conclusion, anesthesia for repair of GBF is challenging. To the best of our knowledge this is the first report in literature describing the anesthetic management of surgical correction of GBF.

**Introduction**

GASTRO-BRONCHIAL FISTULA (GBF) represents an extremely rare complication after surgical procedures. Anesthetic management of such case is challenging. We are presenting a case of GBF following gastric bypass surgery for treating morbid obesity.

*Case report*

A 25-year-old female patient was admitted to our hospital for further investigations and management. She was in her usual state of health until two months back when she underwent in another hospital Roux-en-Y bypass gastrectomy for treatment of morbid obesity. However, three weeks later, she started to develop fever, chest pain and cough. She was

investigated and diagnosed to have leak from the anastomotic line. She underwent surgical exploration and the leak was closed. Three days later she started to have severe productive cough and fever. Chest-x-ray and CT scan showed left sided pleural effusion which was drained with needle aspiration. The drainage turned to be an empyema. She started to improve clinically and was discharged with mild fever and left sided chest pain. She was then admitted to our hospital complaining of productive cough mainly every time she eats and drink fluids.

On examination she was clammy, sweaty with low grade fever. Chest examination showed reduced air entry on the left base and bronchial breathing in the mid lateral zone on the left side with dullness on the left base with otherwise good air entry on the right side. Laboratory and biochemical analysis were within normal ranges. Chest-x-ray showed fluid collection in the left base, the right lung was normal. CT scan showed thickened parietal pleura in the left side with fluid collection. The upper abdomen showed large spleen with very small fluid collection in the left subphrenic space. Barium swallow showed free fluid in different esophageal parts with peri-anastomotic thin fistulous tract directed into the

<sup>1</sup>Professor, Department of Anesthesia, College of Medicine, King Saud University;

<sup>2</sup>Professor of Thoracic Surgery, King Saud University;

<sup>3</sup>Assistant Professor, College of Medicine, King Saud University;

<sup>4</sup>Senior Registrar Thoracic Surgery, King Khalid University Hospital, Riyadh, Saudi Arabia.

*Address for correspondence:* A. El-Dawlatly, Professor, Department of Anaesthesia, P.O. Box 2925, Riyadh 11461, Saudi Arabia.  
[dawlatly@ksu.edu.sa](mailto:dawlatly@ksu.edu.sa)

left lower intrathoracic pleura (Figure 1). The patient was scheduled to undergo left exploratory thoracotomy.



**Figure 1.** Barium meal shows gastro-bronchial fistula. Arrow at fistula.

Preoperative visit revealed 25-year-old female patient, body weight 78kg and height 167cm. All laboratory and biochemical analysis as well as pulmonary function tests were within normal ranges. She was kept fasting night of surgery. Premedication included oral lorazepam 2mg at night and 2mg two hr preoperatively. Also oral ranitidine 150mg was prescribed two hr preoperatively. In the operating room, routine monitoring included, lead II ECG, non-invasive and invasive blood pressure monitoring, tissue oxygen saturation, end tidal CO<sub>2</sub> and rectal temperature probe. Peripheral and radial arterial lines were established. Thoracic epidural D5-6 was performed before induction of general anesthesia. After pre-oxygenation, induction of anesthesia was achieved with sufentanil 10mic and propofol 200mg with cricoid pressure technique. Succinylcholine 100mg was given to facilitate endotracheal intubation. Left sided double lumen tube (DLT) 37Fr was used. Immediately after its insertion the left lung was isolated. Fiberoptic bronchoscope was used to verify the correct placement of DLT. Right sided internal jugular vein was cannulated. Anesthesia was maintained with 50% O<sub>2</sub> in air and 1MAC sevoflurane. Muscle relaxation was maintained with incremental doses of cis-atracurium. Analgesia was achieved with epidural bupivacaine 0.5% (7cc). Left exploratory thoracotomy, decortication and resection of fistula

tract through the diaphragm with segmentectomy were performed (Figure 2).



**Figure 2.** Two artery forceps, one in the bronchial fistulous tract and the other in the diaphragm.

The operation lasted for 90min and at the end reversal agents (atropine/neostigmine) in the usual doses were given i.v and the trachea was extubated. The patient was then sent to surgical intensive care unit for further observation.

### Discussion

Gastro-bronchial fistula is rare complication following gastric surgery. There are few and sporadic case reports published in the literatures. Most of the reported cases followed esophageal surgery for treating cancer esophagus.<sup>1, 2</sup> Also, benign GBF has been reported following esophagectomy.<sup>3</sup> Delayed perforation of the stomach following splenectomy leading to GBF although rare, has been reported.<sup>4</sup> In the literature only one case has been reported following laparoscopic gastric banding for treatment of morbid obesity.<sup>5</sup> The diagnosis of GBF should be suspected when a patient coughs gastric contents, develop recurrent lower respiratory infections or hemoptysis. Investigations include bronchoscopy, barium meal and CT chest to identify lung pathology. Surgical correction remains an option. However, conservative treatment is possible with control of sepsis, adequate drainage and adequate nourishing the patient.<sup>6</sup>

Anesthesia for repair of GBF is challenging. The challenges include, dealing with septic patient with repeated lower respiratory tract infection and possible pulmonary aspiration in malnourished patient.

Preoperative preparation of these patients for surgery is important for better outcome. Rapid sequence induction of anesthesia and proper isolation of the sound lung with double lumen tube are important to minimize incidence of pulmonary aspiration and soiling of the non-operated lung at induction of anesthesia. All precautions of managing one lung ventilation during surgery are to be undertaken.

In conclusion, and to the best of our knowledge this is the first report in literature describing the anesthetic management of surgical correction of GBF.

### References

1. Pramesh CS, Sharma S, Saklani AP and Sanghvi BV. Broncho-gastric fistula complicating transthoracic esophagectomy. *Dis Esoph* 2001;14:271-3.
2. Baisi A, Fumagalli U, Rosati R, et al. Successful primary treatment of bronchial fistula complicating esophagectomy. *Dis Esoph* 1994;7:209-11.
3. Devbhandari MP, Jain R, Galloway S, et al. Benign gastro-bronchial fistula- an uncommon complication of esophagectomy: case report. *BMC Surgery* 2005;5:16.
4. Jeganathan R, Pore N and Clements WDB. Gastrobronchial fistula-a complication of splenectomy. *The Ulster Medical J* 2004;73:126-28.
5. Rassias DJ, Fabian T and Ilves R. Gastrobronchial fistula secondary to laparoscopic gastric band surgery. *Chest* 2004;126:942S.
6. Li YD, Li MH, Han XW, et al. Gastrotracheal and gastrobronchial fistulas: Management with covered expandable metallic stents. *J Vasc Interv Radiol* 2006;17:1649-1656.