

## Sample of the question

In each of the following questions choose ONE best answer and shade the appropriate circle IN PART (I) of your answer sheet

1. The nerve that might be injured in case of extraction of the lower third molar tooth is:

- a. Inferior alveolar.
- b. Lingual.
- c. Hypoglossal.
- d. Nerve to mylohyoid.
- e. Deep temporal nerves.

**Ans: is B**

2) The muscle that is partially inserted into the disc of temporomandibular joint is:

- a. Medial pterygoid.
- b. Anterior fibers of temporalis.
- c. Masseter.
- d. Posterior fibers of temporalis.
- e. Lateral pterygoid.

**Ans is E**

3) The nerve the most liable to be injured in case of thyroidectomy is:

- a. Recurrent laryngeal.
- b. Superior laryngeal.
- c. Pharyngeal branch of vagus.
- d. Internal laryngeal.
- e. Sympathetic trunk.

**Ans: is A**

4) All the following structures are within the parotid gland, EXCEPT:

- a. Parotid lymph nodes.
- b. Facial nerve.
- c. Facial artery.
- d. Retromandibular vein.
- e. External carotid artery.

**Ans: is C**

5) Infection in the region of face drained by the angular vein may result in venous thrombosis of the cavernous sinus through:

- a. Infraorbital vein.
- b. Maxillary vein.
- c. Superior ophthalmic vein.
- d. Transverse facial vein.
- e. Supratrochlear vein.

**Ans: C**