

EXACT TRAVELLING WAVE SOLUTIONS FOR CATTANEO'S REACTION DIFFUSION FOR PRISONER'S DILEMMA GAME

E. S. FAHMY*, H. A. ABDUSALAM and M. LAHLOUH

*Mathematics Department
King Saud University
Women Students Medical Studies & Sciences Sections
P. O. Box 22452, Riyadh 11495,, Saudi Arabia
e-mail: esfahmy@operamail.com

Mathematics Department
College of Science
Kingdom of Saudi Arabia
P. O. Box 2455, Riyadh 11451, Saudi Arabia
e-mail: hosny@operamail.com

Abstract

In this work, we use factorization method to find explicit exact particular travelling wave solutions for or Cattaneo's reaction diffusion for Prisoner's dilemma (PD) game. Using the particular solutions for these equations we find the two-parameter solutions for the equation.

1. Introduction

It is known now that diffusion in many branches of sciences is better modelled by Cattaneo's equation (Telegraph reaction diffusion equation)

2000 Mathematics Subject Classification: **Kindly Provide.**

Keywords and phrases: factorization method, travelling wave solutions, Cattaneo's reaction diffusion, Prisoner's dilemma game.

Received April 14, 2007

where memory effects are included [1-5]. E. Ahmed and S. Z. Hassan, have found an implicit travelling wave solutions for HD and PD Cattaneo's reaction diffusion equations by transforming the equations into first order equations and use a special choice for the wave speed.

Recently, factorization of second-order linear differential equations became a well established technique to find solutions in an algebraic manner [6-8]. Rosu and Cornejo find one particular solution once the nonlinear equation is factorized with the use of two first order differential operators [6, 7]. They use the method for equations of types:

$$u'' + \gamma u' + f(u) = 0, \quad (1)$$

where γ is a constant and

$$u'' + g(u)u' + f(u) = 0, \quad (2)$$

where $'$ means the derivative $D = \frac{d}{dz}$, $g(u)$, $f(u)$ are polynomials in u .

Now, equation (2) can be factorized as

$$[D - \varphi_2(u)][D - \varphi_1(u)]u = 0, \quad (3)$$

which leads to the equation

$$u'' - \frac{d\varphi_1}{du} uu' - \varphi_1 u' - \varphi_2 u' + \varphi_1 \varphi_2 u = 0, \quad (4)$$

or

$$u'' - \left(\varphi_1 + \varphi_2 + \frac{d\varphi_1}{du} u \right) u' + \varphi_1 \varphi_2 u = 0. \quad (5)$$

Comparing (5) and (2) we find

$$g(u) = -\left(\varphi_1 + \varphi_2 + \frac{d\varphi_1}{du} u \right) \text{ and } f(u) = \varphi_1 \varphi_2 u. \quad (6)$$

If equation (2) can be factorized as in equation (3), then a first particular solution can be easily found by solving

$$[D - \varphi_1(u)]u = 0. \quad (7)$$

2. Explicit Exact Travelling Wave Solutions for Prisoner's Dilemma Game

Prisoner's dilemma is a 2×2 symmetric game in which two possible strategies cooperate (A) or defect (B), with payoff matrix

$$\text{matrix} \begin{bmatrix} & A & B \\ A & R & S \\ B & T & U \end{bmatrix}.$$

Cattaneo's reaction diffusion equation for Prisoner's dilemma game is given as [4]:

$$\tau \frac{\partial^2 P}{\partial t^2} + \left(1 - \tau \frac{df}{dP}\right) \frac{\partial P}{\partial t} = \frac{\partial^2 P}{\partial x^2} + f(P), \quad f(P) = -P(1-P)(\gamma - \alpha P), \quad (8)$$

where $\alpha = R + U - S - T$, $\gamma = U - S$ with and $T > R > U > S$, $2R > T + S$.

Using the coordinate transformation $z = x - ct$ (c is the propagation speed) in equation (8) we obtain the following nonlinear ordinary differential equation

$$P'' + g(P)P' + F(P) = 0, \quad (9)$$

where

$$g(P) = \frac{c(1 - \tau\gamma)}{(\tau c^2 - 1)} - \frac{2\tau c(\alpha + \gamma)}{(1 - \tau c^2)} P + \frac{3\alpha}{(1 - \tau c^2)} P^2,$$

$$F(P) = \frac{1}{(1 - \tau c^2)} P(1 - P)(\alpha P - \gamma), \quad 1 > \tau c^2.$$

Using operator notation, equation (9) takes the form

$$\left[D^2 + g(P)D + \frac{F(P)}{P} \right] P = 0. \quad (10)$$

The factorization of (10) leads to

$$[D - \Psi_2(P)][D - \Psi_1(P)]P = 0, \quad (11)$$

and then

$$P'' - \left[\Psi_2 + \Psi_1 + \frac{d\Psi_1}{dP} P \right] P' + \Psi_2 \Psi_1 P = 0. \quad (12)$$

Comparing (12) and (9) we obtain the conditions on Ψ_1 and Ψ_2 as:

$$-\left[\Psi_2 + \Psi_1 + \frac{d\Psi_1}{dP} P \right] = \frac{c(1 - \tau\gamma)}{(1 - \tau c^2)} - \frac{2\tau c(\alpha + \gamma)}{(1 - \tau c^2)} P + \frac{3\alpha}{(1 - \tau c^2)} P^2, \quad (13)$$

$$\Psi_2 \Psi_1 = \frac{1}{(1 - \tau c^2)} (1 - P)(\alpha P - \gamma). \quad (14)$$

Now, we choose Ψ_1 and Ψ_2 such that

$$\Psi_1(P) = \frac{\alpha}{\sqrt{(1 - \tau c^2)}} (1 - P), \quad \Psi_2(P) = \frac{1}{\alpha \sqrt{(1 - \tau c^2)}} (\alpha P - \gamma), \quad \alpha \neq 0. \quad (15)$$

From (15) and (13), we get

$$\sqrt{(1 - \tau c^2)} a^2 + c(1 - \tau\gamma)a + \gamma \sqrt{(1 - \tau c^2)} = 0 \quad (16)$$

which implies that

$$a = \frac{-c(1 - \tau\gamma) \pm \sqrt{c^2(1 - \tau\gamma)^2 - 4\gamma(1 - \tau c^2)}}{2\sqrt{(1 - \tau c^2)}}, \quad (17)$$

and

$$\left[D - \frac{1}{\alpha \sqrt{(1 - \tau c^2)}} (\alpha P - \gamma) \right] \left[D - \frac{\alpha}{\sqrt{(1 - \tau c^2)}} (1 - P) \right] P = 0, \quad (18)$$

and the compatible first order differential equation is

$$P' - \delta(1 - P)P = 0, \quad \delta = \frac{\alpha}{\sqrt{(1 - \tau c^2)}}. \quad (19)$$

By direct integration we get

$$P_1(z) = \frac{1}{1 \pm \exp[\delta(z - z_0)]}. \quad (20)$$

Now if we choose Ψ_1 and Ψ_2 such that

$$\Psi_1(P) = \frac{r}{\sqrt{(1-\tau c^2)}}(\alpha P - \gamma), \quad \Psi_2(P) = \frac{1}{r\sqrt{(1-\tau c^2)}}(1-P), \quad r \neq 0, \quad (21)$$

then we can find that

$$r = \frac{-c(1-\tau\gamma) + \sqrt{c^2(1-\tau\gamma)^2 + 4\gamma(1-\tau c^2)}}{2\gamma\sqrt{(1-\tau c^2)}}, \quad (22)$$

and

$$\left[D - \frac{1}{r\sqrt{(1-\tau c^2)}}(1-P) \right] \left[D - \frac{r}{\sqrt{(1-\tau c^2)}}(\alpha P - \gamma) \right] P = 0, \quad (23)$$

the compatible first order differential equation is

$$P' - \eta(\alpha P - \gamma)P = 0, \quad \eta = \frac{r}{\sqrt{(1-\tau c^2)}}, \quad (24)$$

then

$$P_1(z) = \frac{\alpha}{\gamma \pm \exp[\eta(z - z_0)]}. \quad (25)$$

3. Conclusions

In this paper, the efficient factorization method that was proposed by H. C. Rosu and O. Cornejo-Pérez [6, 7] has been applied to Cattaneo's Reaction Diffusion for Hawk-Dove Games. Exact particular solutions have been obtained.

References

- [1] H. A. Abdusalam, Asymptotic solution of wave front of the telegraph model of dispersive variability, *Chaos Solitons Fractals* 30 (2006), 1190-1197.
- [2] H. A. Abdusalam and E. S. Fahmy, Cross-diffusional effect in a telegraph reaction diffusion Lotka-Volterra two competitive system, *Chaos Solitons Fractals* 18 (2003), 259.
- [3] E. Ahmed, H. A. Abdusalam and E. S. Fahmy, On telegraph reaction diffusion and coupled map lattice in some biological systems, *Internat. J. Modern Phys. C* 2(5) (2001), 717.

- [4] E. Ahmed and S. Z. Hassan, On Diffusion in Some Biological and Economic Systems, *Z. Naturforsch* 55a (2000), 669.
- [5] E. S. Fahmy, Exact solution of the generalized time-delayed Burgers equation through the improved tanh-function method, *Journal of Mathematical Control Science and Applications* (accepted).
- [6] H. C. Rosu and O. Cornejo-Pérez, Supersymmetric pairing of kinks for polynomial nonlinearities, *Phys. Rev. E* 71 (2005), 046607.
- [7] O. Cornejo-Pérez and H. C. Rosu, Nonlinear second order ODE'S: Factorization and particular solutions, *Prog. Theor. Phys.* 114 (2005), 533.
- [8] E. S. Fahmy, Travelling wave solutions for some time-delayed equations through factorizations, *Chaos Solitons Fractals* (to appear).

Kindly return the proof after correction to:

*The Publication Manager
Pushpa Publishing House
Vijaya Niwas
198, Mumfordganj
Allahabad-211002 (India)*

along with the print charges*
by the fastest mail

***Invoice enclosed**

Proof read by:

Signature:

Date:

Tel:

Fax:

E-mail:

Number of additional reprints required

.....

Cost of a set of 25 copies of additional reprints @ U. S. Dollars 15.00 per page.
(25 copies of reprints are provided to the corresponding author ex-gratis)