Abd El-Ghani, M.M. and A.H. Marei. (2006) Vegetation associates of the endangered *Randonia africana* Coss. and its soil characteristics in an and desert ecosystem of western Egypt. Acta Botanica Croatica 65: 83-99

Abd El-Ghani, M.M., R. Bornkamm and F. Darius. (2003). Plant communities in two vegetation transects in the extreme desert of western Egypt. Phytocoenologia 33: 29-48

Caglar, G., S. Caglar, O. Ergin and M. Yarim. (2005). The influence of growth regulators on shoot proliferation and rooting of in vitro propagated caper. Journal of Environmental Biology 26: 479-485

Chalak, L. and A. Elbitar. (2006). Micropropagation of *Capparis spinosa* L. subsp *rupestris* Sibth. & Sm. by nodal cuttings. Indian Journal of Biotechnology 5: 555-558

Chalak, L., A. Elbitar, N. Cordahi, C. Hage and A. Chehade. (2003). In vitro propagation of *Capparis spinosa* L. Acta Horticulturae 616: 335-338

Cora, A., M.N. Bregaglio and R.O. Coirini. (2005). Short communication. Goat preferences for native woody shrubs in the Chaco Arido region of Cordoba, Argentina. Spanish Journal of Agricultural Research 3: 243-247

Goykovic, V., C. Botti and E. Doussoulin. (2003). Anatomical characterization of the roots of caper plant (*Capparis spinosa* L.) grown in two different agroecosystems of northern Chile. Idesia 21: 89-95

Hamed, A.R., K.A. Abdel-Shafeek, N.S. Abdel-Azim, S.I. Ismail and F.M. Hammouda. (2007). Chemical investigation of some *Capparis* species growing in Egypt and their antioxidant activity. Evid Based Complement Alternat Med. 4(1): 25–28

Inocencio, C., R.S. Cowan, F. Alcaraz, D. Rivera and M.F. Fay. (2005). AFLP fingerprinting in *Capparis* subgenus *Capparis* related to the commercial sources of capers. Genetic Resources and Crop Evolution 52: 137-144

Levizou, E., P. Drilias and A. Kyparissis. (2004). Exceptional photosynthetic performance of *Capparis spinosa* L. under adverse conditions of Mediterranean summer. Photosynthetica (Prague) 42: 229-235

Mali, R.G., S. Mahajan and K.S. Patil. (2005). Anthelmintic activity of root bark of *Capparis spinosa*. Indian Journal of Natural Products 21: 50-51

Mandeel, Q. and A. Taha. (2005). Assessment of in vitro antifungal activities of various extracts of indigenous Bahraini medicinal plants. Pharmaceutical Biology 43: 340-348

Pascual, B., A. San Bautista, A. Imbernon, S. Lopez-Galarza, J. Alagarda and J.V. Maroto. (2004). Seed treatments for improved germination of caper (*Capparis spinosa*). Seed Science and Technology 32: 637-642

Pascual, B., A. San Bautista, N. Ferreros, S. Lopez-Galarza and J.V. Maroto. (2003). Analysis of germination of caper seeds as influenced by the position of fruit on the mother plant, fruit maturation stage and fruit weight. Journal of Horticultural Science & Biotechnology 78: 73-78

Proestos, C., I.S. Boziaris, G.J.E. Nychas and M. Komaitis. (2006). Analysis of flavonoids and phenolic acids in Greek aromatic plants: Investigation of their antioxidant capacity and antimicrobial activity. Food Chemistry 95: 664-671

Qasem, J.R and A.A. Hassan. (2003). Herbicidal properties of some medicinal plants against *Malva sylvestris* and *Portulaca oleracea*. Dirasat Agricultural Sciences 30: 84-100

Rhizopoulou, S. and G.K. Psaras. (2003). Development and structure of drought-tolerant leaves of the Mediterranean shrub *Capparis spinosa* L. Annals of Botany (London). 92: 377-83

Rhizopoulou, S., E. Ioannidi, N. Alexandredes and A. Argiropouios. (2006). A study on functional and structural traits of the nocturnal flowers of *Capparis spinosa* L. Journal of Arid Environments 66: 635-647

Romeo, V., M. Ziino, D. Giuffrida, C. Condurso A. Verzera. (2007). Flavour profile of capers (*Capparis spinosa* L.) from the Eolian Archipelago by HS-SPME/GC-MS. Food Chemistry 101: 1272-1278

Sharaf, M., M.A. El-Ansari, N.A. Saleh. 91997). Flavonoids of four *Cleome* and three *Capparis* species. Biochem. Syst. Ecol. 25:161–166

Strack, D. and T. Fester. (2006). Isoprenoid metabolism and plastid reorganization in arbuscular mycorrhizal roots. New Phytologist 172: 22-34

Turkmen, N. and A. Duzenli. (2005). Changes in floristic composition of *Quercus coccifera macchia* after fire in the Qukurova region (Turkey). Annales Botanici Fennici 42: 453-460