

PEAK EXPIRATORY FLOW RATE IN A S AMPLE OF NORMAL SAUDI MALES AT RIYADH, SAUDI ARABIA

Ahmed A. Al-Taweel, **ABFM**, Khalid A. Kalantan, **ABFM**, Hamza A. Ghani, **ABFM**

Department of Family and Community Medicine, College of Medicine, King Saud University, Riyadh, Saudi Arabia

هدف الدراسة: هو إيجاد المعدل الطبيعي للقدرة القصوى على الزفير للسعوديين البالغين من الرجال ومقارنته بالمعدل الطبيعي لدى البريطانيين.

طريقة البحث : عملت دراسة مقطعية في خمسة مراكز صحية أولية تمثل مدينة الرياض في الفترة من ١٥ نوفمبر إلى نهاية ديسمبر ١٩٩٣م. ستمائة وثمانين رجلاً سعودياً والذين انطبقت عليهم صفات سترنجنت للأشخاص الطبيعيين تضمنتهم هذه الدراسة معظمهم تحت سن ٥٤ عاماً (٩٤٪).

نتائج البحث: كان معدل ومعامل التشنت للعمر والطول للمشاركين في هذه الدراسة هما ٢٨,٤ ± ١٣ و ١٦٧,٦ ± ٦,٤ على التوالي. وعمل تحليل للنتائج بخطوات متعددة لتحديد طبيعة خط الانحدار للقدرة القصوى على الزفير بحسب العمر والطول. رسمت خطوط الانحدار ووجد أن القدرة القصوى على الزفير لا تبدأ بالنزول حتى سن ٢٥ عاماً. وكلما زاد الطول تزداد القدرة القصوى على الزفير بعلاقة طردية.

الاستنتاجات : وجد في هذه الدراسة أن معدلات القدرة على الزفير لدى السعوديين أقل مقارنة بالبريطانيين. هذه النتائج ستخدم كقاعدة لإيجاد المعدلات الطبيعية للقدرة القصوى على الزفير لسكان المملكة.

الكلمات المرجعية : القدرة القصوى على الزفير، الربو الشعبي، المملكة العربية السعودية.

Objective: To find out the normal peak expiratory flow rate for adult Saudi males and to compare our

standards with British standards.

Methods: A cross-sectional study was carried out in five primary health care centers representing Riyadh city in the period between 15th November through December 1993. Six hundred and eighty Saudi men who satisfied stringent criteria of normality were included in the study.

Results: The mean and standard deviation of the subjects' age and height were 28.4 ± 13 and 167.6 ± 6.4 respectively. Linear regression analysis was performed through step-wise procedure to determine the form of regression of peak expiratory flow on age and height. Regression curves were obtained and it was found that peak expiratory flow rate did not begin to decline until about the age of 25 years; and as height increased the peak expiratory flow rate increased in a linear relationship.

Conclusion: It was demonstrated that our study group had lower peak expiratory flow rate compared with British people. These findings will serve as a basis for preparing flow rate values for our population.

Key Words: Peak expiratory flow rate, Bronchial asthma, Saudi Arabia.

Correspondence to:

Dr. Hamza Abdul Ghani, Consultant Family Physician, Department of Family & Community Medicine (34), College of Medicine, King Saud University, P.O. Box 2925, Riyadh 11461, Saudi Arabia

INTRODUCTION

Bronchial asthma is one of the most important and highly prevalent problems in Saudi Arabia. The prevalence of asthma in Saudi Arabia ranges between 10-17% (10% in Riyadh, 13% in Jeddah and Qassim and 17% in Abha); which is higher than the range reported in Western countries.^{1,2}

Pulmonary function tests are important in the detection and evaluation of pulmonary dysfunction. Peak expiratory flow rate (PEFR) is the simplest estimate for lung functions. It indicates the severity of airflow limitation.^{3,4} Peak expiratory flow rate is measured by peak expiratory flow meter (PEFM) which is a simple and relatively cheap device (cost ranges from 25-40 US dollars). It has a great diagnostic and prognostic value in patients with hyperactive airway disease.^{5,7} It is widely used in general practice in the UK and in family practice in the USA with proved importance in evaluating and monitoring patients with bronchial asthma.⁶ Although PEFM is a simple, practical and effective tool for use in the primary health care (PHC) setting, few of PHC practitioners in Saudi Arabia seem to use it.⁸

Many factors can affect the reading of peak expiratory flow rate (PEFR). The age, sex and height are important variables upon which peak expiratory flow depends.⁹ Other factors include the diurnal variation¹⁰ and ethnic differences.¹¹ The degree to which both the instructor and the patient understand the technique of using the flow meter is important and can greatly affect the result. Widely used standard peak flow rates have not been established for our population and the PEFR tables in use are based on studies in the Western countries. Not many local studies on the PEFR of Saudis in PHC clinics have been done.^{12,13} The objective of the current study was to identify the normal peak expiratory flow rate for a sample of Saudi males and to compare the findings with British standards.

MATERIAL AND METHODS

Riyadh (the capital of Saudi Arabia) is divided into five main sectors for the provision of the Health Services (i.e., North, South, East, West and Central). A number was given to each center in all sectors, and five primary health care centers were randomly selected to represent different geographical areas. Within each center, patients were enrolled randomly in a systematic way to ensure a good non-biased presentation. The sample size for the centers was nearly the same to make the allocation equal. Subjects included in the study were Saudi males aged 15 years and above, non-smokers who satisfied

stringent criteria of normality.¹⁰ These were as follows 1) had never smoked; 2) were not subject to recurrent or persistent expectoration; 3) were not subject to wheezing nor had had an episode of acute bronchitis; 4) had not had asthma or recurrent bronchitis during their childhood; 5) had not had any serious respiratory disease.

Since it was difficult for the subjects to recall one attack of acute bronchitis, this criterion was not used to exclude any. The number of subjects included in this study was 680. Each center was provided with mini-Wright peak flow meter (product of Ciement Clarke International Ltd, England), with disposable mouth pieces. All the mini-Wright peak expiratory flow meters were new and had been calibrated to make sure that they had similar readings. Only males were included in the study to ensure accessibility so that we could deal directly with the subjects and supervise more closely. It was agreed that all the tests be done in the afternoon sessions (4 -8 p.m.) to avoid the effect of the diurnal variation of peak expiratory flow¹⁰ and get more volunteers.

The Director of each health center allocated one of the male nurses to work with the investigators. To ensure a unified correct procedure, all the nurses were instructed and trained by the assigned investigator on the technique of using the device. A group of subjects were instructed to do the test in front of both the nurse and the investigator to ensure that they perform it correctly and to use this as demonstration for the nurse.

The instructions adopted¹⁴ for using the peak flow meter were as follows: 1) Every subject was asked to do the test three times. 2) Take a full deep breath in. 3) Hold the peak flow meter horizontally and close the lips tightly round the mouth piece. 4) Blow out as hard and as fast as you can in a short sharp blow. 5) Wait one minute between each blow. Take the best of 3 readings. This was recorded on the table along with subject's age and height.

Data were analyzed using Stat Pac Gold statistical analysis package. Linear regression step-wise analysis was performed. The mean was expressed as mean \pm 1 standard deviation.

RESULTS

The age distribution of a total of 680 subjects included in this study is shown in Table 1. The mean and standard deviation of their age and height were 28.4 ± 13 and 167.6 ± 6.4 respectively.

Linear regression analysis was performed through step-wise procedure to determine the form of regression of peak expiratory flow on age and height (Table 2). When the data was plotted it became evident that the relation between the variables was curvilinear (Figure 1). It showed that as age increased the PEFR decreased ($b = -0.5$, $T = -2.5$, p -value 0.012). However, it showed a positive association between height of the study population and PEFR ($b=4.3$, $T=11.7$, p -value < 0.0001). From the multiple regression equation $R^2 = 21\%$ and F _ratio + 46%, p _value = < 0.0001 , i.e., about 21% variability of the PEFR can be determined by the factors in the equation.

Table 1: Age distribution of subjects in series

Age in years	Number	%
< 25	358	52.6
25 – 34	142	20.9
35 – 44	93	13.7
45 – 54	49	7.2
55 – 64	25	3.7
≥ 65	13	1.9
Total	680	100

Table 2: Regression Equations

Parameter	Coefficient	Value
Constant	b_0	-338.5
Age	b_1	20.3
Age ²	b_2	-0.55

However, a significant positive association was found between the height of the study population and the PEF. The multiple regression equation $R^2 = 21\%$ and fraction = 46% with p-value <0.0001, indicates a significant dependency of the PEF of the study population on their age and height.

One limitation of the current study was the non-inclusion of female subjects since certain cultural difficulties in performing the measurements were envisaged. The other limitation was that the computed equation was reliable in predicting only 21%, for the whole range of age groups. However, this study has shown that Saudi Arabian male adults living in Riyadh have lower peak expiratory flow rate than adults of different racial origins living in UK.

There is a need for baseline normal values for peak expiratory flow rate for both children and adults in Saudi Arabia, and the data obtained from this study can be used as a base-line for future studies for the preparation of standard charts for our population. Further studies without the limitations of this study i.e., to include females and a wide range of patient characteristics are required.

ACKNOWLEDGMENT

The authors are grateful to Prof. Jamal S. Jarallah for reviewing the manuscript and Dr. Fayek Khwsky for his statistical comments. We would also like to thank the Directors of all PHCC and nurses assigned to do and supervise the project.

REFERENCES

1. Al Frayh A, Bener A, Al-Jawadi TQ. Prevalence of asthma among Saudi school - children. Saudi Med J 1992; 13 (6): 521-4.
2. Al Frayh A. Prevalence of asthma and allergic rhinitis in the Kingdom of Saudi Arabia. Proceeding of the first annual meeting of the Saudi Society of Allergy and Immunology Jan 1994.
3. Charlton I, Charton G, Broomfield J, Mullee M. Evaluation of peak flow and symptoms only self management plans for control of asthma in general practice. BMJ 1990; 301: 1355-9.

4. Cochrane GM. Asthma: diagnosis and role of PEFr monitoring. Update 1987; 8: 585-89.
5. British Thoracic Association. Death from asthma in two regions of England. BMJ 1982; 285: 1251-5.
6. Katz DN. The Mini Wright peak flow meter for evaluating airway obstruction in a family practice. J Fam Pract 1983; 17: 51-7.
7. Battu K, Collins-Williams C, Zaleskey C. Evaluation of home monitoring of asthmatic children with the mini-wright peak flow meter. J Asthma 1982; 19 (1): 33-7.
8. Khoja Ta, Al-Ansari LA. Asthma in Saudi Arabia: is the system appropriate for optimal case? J Public Health Management Practice 1998, 4(3): 64-72.
9. British Thoracic Society, Research Unit of the Royal College of Physicians of London. Guidelines for the management of asthma in adults chronic persistent asthma. BMJ 1990; 301: 651-3.
10. Gregg I, Nunn A.J. Peak expiratory flow in normal subjects. BMJ 1973; 3: 282-4.
11. Cinkotai FF, Sharpe TC, Gibbs AC. Circadian rhythms in peak expiratory flow rate in workers exposed to cotton dust. Thorax 1984; 39 (10): 759-65.
12. Graff-Lonnevig V, Harfi H, Tipirneni P. Peak expiratory flow rates in healthy Saudi Arabian children living in Riyadh. Ann Allergy 1993; 71 (5): 446-50.
13. Al Fayed SF, Kassimi MA, Ardawi MS. Normal spirometry values in Saudi Nationals in Western Region of Saudi Arabia. Saudi Med Journal; 1989; 56: 227-34.
14. Catterall J, Shapiro C. Nocturnal asthma. BMJ 1993; 306: 1189-93.
15. Koenigsberg M, Holden DM. Peak expiratory flow rate in the elderly. J Fam Pract 1989; 29 :503-6.
16. Cook NR, Evans DA, Scherr PA, Speizer FE, Vedal S, Branch LG. Peak expiratory flow rate in an elderly population. Am J Epidemiol 1989; 130 (1): 66-78.