

# ***Pindolol***

◆ Is a nonselective  $\beta$ -adrenergic blocking agent with high ISA , no membrane stabilizing activity, and moderate lipid solubility. hepatic dysfunction.

# ***PHARMACOKINETIC***

- *Pindolol is almost completely (95%) absorbed orally, with peak plasma concentrations range from 15 to 38 ng/MI*

# CAUTIONS:

- Pindolol may cause bradycardia, claudication, hypotension, syncope, tachycardia, fatigue, headache, dizziness, sleep disturbances, depression, tremors, diarrhea, constipation, abdominal discomfort, impotence and bronchospasm.

# *Pindolol*

- *Pindolol is 40% to 60% protein bound*
- *\*000000000000000000vd is 1.2 to 2 L/kg  
half-life is 3 to 4 hours*
- *Pindolol may cause bradycardia,  
claudication, hypotension, syncope,  
tachycardia, fatigue, headache, dizziness,  
sleep disturbances, depression ,tremors,  
diarrhea, constipation ,abdominal  
discomfort, impotence and bronchospasm.*

# Dosage

- *The usual adult dose is 5mg orally twice daily with incremental increases by 10mg/day every 3 to 4 weeks to achieve blood pressure control to a maximum daily dose of 60 mg .*