


## Work due to friction

If friction is involved in moving objects, work has to be done against the kinetic frictional force.


This work is:



$$W_f = f_k \cdot d = f_k d \cos 180^\circ = -f_k d$$

Dr. Abdallah M. Azzeer

### Example



A block of mass  $m$  is pulled by a force  $F$  for a distance  $d$  on a rough surface as shown. What is the total work done on the block?

$$\hat{y} : F_{net,y} = n + F \sin \theta - mg = 0$$

$$n = mg - F \sin \theta$$

$$\hat{x} : F_{net,x} = F \cos \theta - \mu_k n$$

$$= F \cos \theta - \mu_k (mg - F \sin \theta) = F (\cos \theta + \mu_k \sin \theta) - \mu_k mg$$

$$W_{net} = F_{net,x} \cdot d$$

N.B. find the work of each force and add the work, can you get the same answer? why

**WORK IS SCALAR QUANTITY**

Dr. Abdallah M. Azzeer

2

**Example**


$2.40 \times 10^2$  N force is pulling an 85.0-kg refrigerator across a horizontal surface. The force acts at an angle of  $20.0^\circ$  above the surface. The coefficient of kinetic friction is 0.200, and the refrigerator moves a distance of 8.00 m. Find

- (a) the work done by the pulling force, and  
 (b) the work done by the kinetic frictional force.

(a)  $W = F \cos \theta d = 1.8 \times 10^3$  J

(b)  $W_f = f_k d \cos \theta$   
 $f_k = \mu_k (mg - F \sin \theta)$   
 $= 1.5 \times 10^2$  N

so  $W_f = -1.2 \times 10^3$  J




Dr. Abdallah M. Azzeer

3

### 7.5 Work & Kinetic Energy...

A constant net Force



Work done by  $\vec{F} \Rightarrow W_{net} = \vec{F} \cdot \vec{d} = Fd$

$\therefore F = ma,$

$$v_f^2 = v_i^2 + 2ad \Rightarrow d = \frac{1}{2a}(v_f^2 - v_i^2)$$

$$\Rightarrow W_{net} = ma \cdot \frac{1}{2a}(v_f^2 - v_i^2)$$

$$W_{net} = \frac{1}{2}mv_f^2 - \frac{1}{2}mv_i^2$$

$$K \equiv \frac{1}{2}mv^2$$

$$W_{net} = K_f - K_i = \Delta K$$

{Net Work done on object}  
 =  
 {change in kinetic energy of object}

Dr. Abdallah M. Azzeer

4

## Kinetic Energy; Work-Energy Principle

- **Energy**  $\equiv$  The ability to do work
- **Kinetic Energy**  $\equiv$  The energy of motion
- “Kinetic”  $\equiv$  Greek word for motion
- An object in motion has the ability to do work
- Net work on an object = Change in KE.

$$W_{\text{net}} = \Delta K$$

### **The Work-Energy Principle**

- **Note:**  $W_{\text{net}}$  = work done by the net (total) force.
- $W_{\text{net}}$  is a scalar.
- $W_{\text{net}}$  can be positive or negative (because  $\Delta KE$  can be both + & -)
- Units are Joules for both work & KE.

Dr. Abdallah M. Azzeer

5

Table 7.1

### Kinetic Energies for Various Objects

Object	Mass (kg)	Speed (m/s)	Kinetic Energy (J)
Earth orbiting the Sun	$5.98 \times 10^{24}$	$2.98 \times 10^4$	$2.65 \times 10^{33}$
Moon orbiting the Earth	$7.35 \times 10^{22}$	$1.02 \times 10^3$	$3.82 \times 10^{28}$
Rocket moving at escape speed <sup>a</sup>	500	$1.12 \times 10^4$	$3.14 \times 10^{10}$
Automobile at 65 mi/h	2 000	29	$8.4 \times 10^5$
Running athlete	70	10	3 500
Stone dropped from 10 m	1.0	14	98
Golf ball at terminal speed	0.046	44	45
Raindrop at terminal speed	$3.5 \times 10^{-5}$	9.0	$1.4 \times 10^{-3}$
Oxygen molecule in air	$5.3 \times 10^{-26}$	500	$6.6 \times 10^{-21}$

<sup>a</sup> Escape speed is the minimum speed an object must reach near the Earth's surface in order to move infinitely far away from the Earth.


© 2004 Thomson/Brooks Cole

Dr. Abdallah M. Azzeer

6


### 7.6 the nonisolated system CONSERVATION OF ENERGY

A book sliding to the right on a horizontal surface slows down in the presence of a force of kinetic friction acting to the left. The initial velocity of the book is  $v_i$  and its final velocity is  $v_f$ . The normal force and the gravitational force are not included in the diagram



Dr. Abdallah M. Azzeer

7




Energy transfer mechanisms.  
 (a) Energy is transferred to the block by *work*;  
 (b) energy leaves the radio from the speaker by *mechanical waves*;  
 (c) energy transfers up the handle of the spoon by *heat*;  
 (d) energy enters the automobile gas tank by *matter transfer*;  
 (e) energy enters the hair dryer by *electrical transmission*; and  
 (f) energy leaves the lightbulb by *electromagnetic radiation*.

Dr. Abdallah M. Azzeer

8

# We will be back to this point later

Example 7.9, 7.10 & 7.11



Dr. Abdallah M. Azzeer

9

## A Question

- A box is pulled up a rough ( $\mu > 0$ ) incline by a rope-pulley-weight arrangement as shown below. How many forces are doing work on the box?

- (a) 2
- (b) 3
- (c) 4



### Solution


Draw FBD of box: Consider direction of motion of the box

Any force not perpendicular to the motion will do work:

- $N$  does **no** work ( $\perp v$ )
- $T$  does **positive** work
- $f$  does **negative** work
- $mg$  does **negative** work



3 forces do work



103 PHYS


Dr. Abdallah M. Azzeer

READ EXAMPLES 7.7 in your textbook

Example 7.8

A man loads a refrigerator onto a truck using a ramp. Ignore friction.

He claims he would be doing less work if the length of the ramp would be longer. Is this true?



$$W_{\text{net}} = W_{\text{by man}} + W_{\text{by gravity}} = 0$$


Dr. Abdallah M. Azzeer

11

### Example

A rescue helicopter lifts a 79-kg person straight up by means of a cable. The person has an upward acceleration of  $0.70 \text{ m/s}^2$  and is lifted from rest through a distance of 11 m.

- What is the tension in the cable?
- How much work is done by the tension in the cable?
- How much work is done by the person's weight?
- Use the work-energy theorem and find the final speed of the person.



a)  $T - mg = ma$ ,  $T = 8.3 \times 10^2 \text{ N}$

b)  $W_T = Td = 9.13 \times 10^3 \text{ J}$

c)  $W_w = -mgd = -8.5 \times 10^3 \text{ J}$

d)  $W_T + W_w = \frac{1}{2} mv_f^2 - \frac{1}{2} mv_0^2$   
 $v_f = 3.92 \text{ m/s}$

Dr. Abdallah M. Azzeer

12

**How would you attack this problem?**

$$v_{yi} = v_0 \sin \theta$$

$$a_y = -g$$

$$\Delta y = -h$$

$$v_{yf} = ?$$

$$v_{yf}^2 = v_{yi}^2 + 2a_y \cdot \Delta y$$

Eventually obtaining...

$$v_f = \sqrt{v_0^2 + 2gh}$$

Independent of  $\theta$ !...

**OLD WAY**

Given  $v_0$ ,  $\theta$ , and  $h$ , what is the speed of the projectile the instant before it lands?

*Dr. Abdallah M. Azzeer* 13

**WORK – ENERGY THEOREM**

$$\Delta y = -h$$

$$W_g = -mg\Delta y = mgh$$

$$\Sigma W = W_g = K_f - K_i$$

$$mgh = \frac{1}{2} m v_f^2 - \frac{1}{2} m v_0^2$$

$$v_f^2 = v_0^2 + 2gh$$

$$v_f = \sqrt{v_0^2 + 2gh}$$


**NEW WAY**

Given  $v_0$ ,  ~~$\theta$~~ , and  $h$ , what is the speed of the projectile the instant before it lands?

*Dr. Abdallah M. Azzeer* 14

**Example**

Two blocks have masses  $m_1$  and  $m_2$ , where  $m_1 > m_2$ . They are sliding on a frictionless floor and have the **same kinetic energy** when they encounter a long rough stretch (i.e.  $\mu > 0$ ) which slows them down to a stop. Which one will go farther before stopping?

(a)  $m_1$ (b)  $m_2$ (c) *they will go the same distance*

103 PHYS


Dr. Abdallah M. Azzeer

**Solution****for the first mass  $m_1$** 

- The work-energy theorem says that for any object  $W = \Delta K$
- In this example the only force that does work is **friction** (since both  $N$  and  $mg$  are perpendicular to the blocks motion).
- The net work done to stop the box is  $-f d = -\mu mg d$ .

This work “removes” the kinetic energy that the box had:

$$W = K_f - K_i = 0 - K_i$$



103 PHYS

Dr. Abdallah M. Azzeer




**for the second mass  $m_2$** 

- The net work done to stop a box is  $-fd = -\mu mgd$ .
  - This work “removes” the kinetic energy that the box had:
  - $W = K_f - K_i = 0 - K_i$
- This is the same for both boxes (same starting kinetic energy).

$$\Rightarrow \mu m_2 g d_2 = \mu m_1 g d_1 \quad \Rightarrow \quad m_2 d_2 = m_1 d_1$$

Since  $m_1 > m_2$  we can see that  $d_2 > d_1$



103 PHYS


Dr. Abdallah M. Azzeer

**Example**

0.075-kg arrow is fired horizontally. The bowstring exerts an average force of 65 N on the arrow over a distance of 0.90 m. With what speed does the arrow leave the bow?

$$W_{net} = F \cdot d = \frac{1}{2}mv_f^2 - \frac{1}{2}mv_i^2$$

$$v_f = 39 \text{ m/s}$$



Dr. Abdallah M. Azzeer

18

## 7.8 Power

Power is the rate at which work is done by a force

$$P_{AVG} = W/\Delta t \quad \text{Average Power}$$

$$P = dW/dt \quad \text{Instantaneous Power}$$

The unit of power is a Joule/second (J/s) which we define as a Watt (W)

$$1 \text{ W} = 1 \text{ J/s} = 1 \text{ kg}\cdot\text{m}^2/\text{s}^3$$

$$P = \frac{dW}{dt} = \frac{d}{dt} (\vec{F} \cdot \vec{x})$$

$$P = \vec{F} \cdot \frac{d\vec{x}}{dt} = \vec{F} \cdot \vec{v}$$


In general, power is defined for any type of energy transfer.

$$P = dE/dt$$

Where  $dE/dt$  is the rate at which energy is crossing the boundary of the system by a given transfer mechanism.

*Dr. Abdallah M. Azzeer*

19



$$\Delta W = F\Delta x \cos\theta$$

$$\Delta x = v\Delta t$$

$$P = \frac{\Delta W}{\Delta t} = \frac{F\Delta x \cos\theta}{\Delta t} = Fv \cos\theta$$

Note: power  $\times$  time = work, so work can be measured in units of kWh

$$(1 \text{ kWh} = (10^3 \text{ Watt}) \times (3600 \text{ s}) = 3.6 \times 10^6 \text{ W s} = 3.6 \times 10^6 \text{ J} = 3.6 \text{ MJ.})$$

- British units are hp (horse power)
- 1 hp = 550 ft.lb/s = 746 W

*Dr. Abdallah M. Azzeer*

20

**Example 7.12**

An elevator car has a mass of 1600 kg and is carrying passengers having a combined mass of 200 kg. A constant friction force of 4000 N retards its motion upward, as shown in the figure.

- (a) What power delivered by the motor is required to lift the elevator car at a constant speed of 3 m/s?  
 (b) What power must the motor deliver at the instant the speed of the elevator is  $v$  if the motor is designed to provide the elevator car with an upward acceleration of  $1 \text{ m/s}^2$ ?


(a)  $M_{\text{max}} = 1800 \text{ kg}$ ,  $f = 4000 \text{ N}$   
 $v = 3 \text{ m/s (constant)} \Rightarrow a = 0$

$$\sum F = T - f - Mg = 0$$

$$T = f + Mg = 4000 \text{ N} + 1800 \times 9.8 \text{ N} = 2.16 \times 10^4 \text{ N}$$

$$\text{Power: } P = \vec{T} \cdot \vec{v} = T v = (2.16 \times 10^4)(3) = 64.8 \text{ kW}$$

(b) Left for you to try



Physics

Dr. Abdallah M. Azzeer

21

**Example : Power Needs of Car**

Calculate the power needed for the car (a) to climb a hill. (b) to pass another car.

(a)

$$\sum F_x = 0$$

$$F - F_R - mg \sin \theta = 0$$

$$P = Fv$$

(b)


Now,  $\theta = 0$

$$\sum F_x = ma$$

$$F - F_R = 0$$

$$v = v_0 + at$$

$$P = Fv$$




Dr. Abdallah M. Azzeer

22

**Problem:** determine the power after an elapsed time of 3.0 s

$$\bar{P} = \frac{mgh}{\Delta t} = \frac{50 \text{ kg} \times 9.8 \text{ m/s} \times 2.0 \text{ m}}{3.0 \text{ s}} = 330 \text{ W}$$



*Dr. Abdallah M. Azzeer*

23