

Example 8.5 :

A spring-loaded popgun $x = 0.12 \text{ m}$, $m = 35 \text{ g}$, $h = x_c = 20 \text{ m}$,
Neglect all resistive forces , $k = ?$

$$\Delta K + \Delta U = 0$$

$$K_2 - K_1 + U_2 - U_1 = 0$$

$$K_2 + U_2 = K_1 + U_1 = \text{constant} = E \equiv \text{Total mechanical energy}$$

$$K_2 + U_2 = K_1 + U_1$$


$$0 + U_2 = 0 + U_1$$

$$U_2 = U_{2s} + U_{2g} = 0 + mgh$$

$$U_1 = U_{1s} + U_{1g} = \frac{1}{2}kx^2 + 0$$

$$mgh = \frac{1}{2}kx^2$$

$$k = \frac{2mgh}{x^2} = 953 \text{ N/m}$$



103 Phys

Dr. Abdallah M. Azzeer

What is the speed of projectile as it moves through equilibrium position of the spring

$$K_2 + U_2 = K_1 + U_1$$


$$K_2 + U_2 = 0 + U_1$$

$$U_2 = U_{2s} + U_{2g} = 0 + mgx$$

$$U_1 = U_{1s} + U_{1g} = \frac{1}{2}kx^2 + 0$$

$$\frac{1}{2}mv_2^2 + mgh = \frac{1}{2}kx^2$$

$$v_2 = \sqrt{\frac{kx^2}{m} - 2gx} = 19.7 \text{ m/s}$$



103 Phys

Dr. Abdallah M. Azzeer

Example 8.6


$m = 3 \text{ kg}$, $d = 1 \text{ m}$, $\theta = 30^\circ$,
 $v_i = 0$, $f_k = 5 \text{ N}$, $h = 0.5 \text{ m}$,
 $v_f = ?$

$$\Delta K + \Delta U = W_{nc}$$

$$K_f - K_i + U_f - U_i = W_{f_k}$$

$$\frac{1}{2}mv_f^2 - 0 + 0 - mgh = -f_k d$$

$$v_f = \sqrt{\frac{2}{m}(mgh - f_k d)} = 2.54 \text{ m/s}$$



What happen when you don't know h?

103 Phys

Dr. Abdallah M. Azzeer


3

Example 8.6

A child of mass m rides on an irregularly curved slide of height $h = 2.00 \text{ m}$, as shown in Figure 8.12. The child starts from rest at the top.

(A) Determine his speed at the bottom, assuming no friction is present.

(B) If a force of kinetic friction acts on the child, how much mechanical energy does the system lose? Assume that $v_f = 3.00 \text{ m/s}$ and $m = 20.0 \text{ kg}$.



103 Phys


Dr. Abdallah M. Azzeer

4

example 8.8

A skier starts from rest at the top of a frictionless incline of height 20.0 m, as shown. At the bottom of the incline, she encounters a horizontal surface where the coefficient of kinetic friction between the skis and the snow is 0.210.

(a) How far does she travel on the horizontal stretch.



103 Phys

Dr. Abdallah M. Azzeer

5

$$\Delta E = 0 = \Delta K + \Delta U$$

$$mgh = \frac{1}{2}mv^2$$

$$v = \sqrt{2gh}$$

$$v = \sqrt{2 \times 9.8 \times 20.0} = 19.8 \text{ m/s}$$

$$\Delta K = K_f - K_i = -f_k d$$

Since $K_f = 0$ $-K_i = -f_k d$; $f_k d = K_i$

$$f_k = \mu_k n = \mu_k mg$$

$$d = \frac{K_i}{\mu_k mg} = \frac{\frac{1}{2}mv^2}{\mu_k mg} = \frac{v^2}{2\mu_k g} = \frac{(19.8)^2}{2 \times 0.210 \times 9.80} = 95.2 \text{ m}$$


103 Phys

Dr. Abdallah M. Azzeer

6

Example 8.9

A block sliding on a smooth, horizontal surface collides with a light spring. (a) Initially the mechanical energy is all kinetic energy. (b) The mechanical energy is the sum of the kinetic energy of the block and the elastic potential energy in the spring. (c) The energy is entirely potential energy. (d) The energy is transformed back to the kinetic energy of the block. The total energy of the system remains constant throughout the motion.



(i) $\mu_k = 0$

$$E_i = \frac{1}{2} m v_i^2$$

$$E_i = \frac{1}{2} k x_m^2 \Rightarrow x_m = \sqrt{\frac{m}{k}}$$

$$E_i = \frac{1}{2} m v_f^2 \Rightarrow \vec{v}_f = -\vec{v}_i$$


(i) $\mu_k \neq 0$

$$E_i - \mu_k m g (x_m + x_0) = \frac{1}{2} k x_m^2$$

$$E_i - \mu_k m g 2(x_m + x_0) = \frac{1}{2} m v_f^2$$

Example 8.10

As the hanging block moves from its highest elevation to its lowest, the system loses gravitational potential energy but gains elastic potential energy in the spring. Some mechanical energy is lost because of friction between the sliding block and the surface.



MINI REVIEW: WORK – KE – PE

Work done by constant force: $W = \mathbf{F} \cdot \mathbf{d} = Fd \cos\theta$

e.g. Work done by gravity: $W_g = -mg \Delta y$

Change in gravitational PE: $\Delta U_g = -W_g = mg \Delta y$

Work done by variable force: $W = \int_{x_1}^{x_2} F_x dx$

e.g. Work *by* spring: $W_{x_1 \rightarrow x_2} = -\frac{1}{2}k(x_2^2 - x_1^2)$

Change in spring PE: $\Delta U = -W_s = \frac{1}{2}k(x_2^2 - x_1^2)$

Work – Energy Thm: $\Sigma W_{non-con} = \Delta E_{mech} = \Delta K + \Delta U$

103 Phys


Dr. Abdallah M. Azzeer

9

Example : Hotwheel

A toy car slides on the frictionless track shown below. It starts at rest, drops a distance d , moves horizontally at speed v_1 , rises a distance h , and ends up moving horizontally with speed v_2 .

Find v_1 and v_2 .



103 Phys

Dr. Abdallah M. Azzeer

10

- K+U energy is conserved, so $\Delta E = 0$

$$\Delta K = -\Delta U$$

- Moving down a distance d ,

$$\Delta U = -mgd, \quad \Delta K = \frac{1}{2}mv_1^2 \quad \Rightarrow$$

- Solving for the speed:

$$v_1 = \sqrt{2gd}$$



103 Phys

Dr. Abdallah M. Azzeer


11

- At the end, we are a distance $d - h$ below our starting point.

$$\Delta U = -mg(d - h), \quad \Delta K = \frac{1}{2}mv_2^2$$

Solving for the speed:

$$v_2 = \sqrt{2g(d - h)}$$



103 Phys

Dr. Abdallah M. Azzeer

12

Example:

With what speed does the weight have just before contact with the nail?

$$\Delta K + \Delta U = 0$$


$$U_i = mgh$$

$$U_f = 0$$

$$K_i = 0$$

$$K_f = \frac{1}{2}mv^2$$

$$v = \sqrt{2gh}$$



103 Phys

Dr. Abdallah M. Azzeer

13

What is the force of resistance between the nail and the block?

$$\Delta K + \Delta U = W_{nc} = -fd$$

$$U_i = 0$$


$$U_f = -mgd$$

$$K_i = mgh$$

$$K_f = 0$$

Solve you get

$$f = mg \left(\frac{h+d}{d} \right)$$



103 Phys


Dr. Abdallah M. Azzeer

14

Using Energy to Find Resistive Forces

Pendulum & Sliding Block

What is the work done by friction?



$$KE_i = KE_f = 0$$

$$U_i = mgh$$

$$= mgL(1 - \cos \theta)$$

$$U_f = 0$$

$$\Delta KE + \Delta U = W_{nc}$$

$$W_{nc} = \Delta U = mgL(1 - \cos \theta) = -fd$$

103 Phys


Dr. Abdallah M. Azzeer

15

Example:

A block slides down a frictionless ramp. Suppose the horizontal (bottom) portion of the track is rough, such that the coefficient of kinetic friction between the block and the track is μ_k .

How far, x , does the block go along the bottom portion of the track before stopping?



Using $W_{nc} = \Delta K + \Delta U$

As before, $\Delta U = -mgd$

W_{nc} = work done by friction = $-\mu_k mgx$.

$\Delta K = 0$ since the block starts out and ends up at rest.

$$W_{nc} = \Delta U \Rightarrow -\mu_k mgx = -mgd \Rightarrow x = d / \mu_k$$


103 Phys

Dr. Abdallah M. Azzeer

16

Example

Two blocks, A and B ($m_A=50\text{ kg}$ and $m_B=100\text{ kg}$), are connected by a string as shown. If the blocks begin at rest, what will their speeds be after A has slid a distance $d = 0.25\text{ m}$? Assume the pulley and incline are frictionless.



ANS: 1.51 m/s

103 Phys

Dr. Abdallah M. Azzeer


17

Example:

A skier ($m=58\text{ kg}$) is traveling down a 25 degree slope. His skies against the snow exert a frictional force of 70 N. He starts out with a velocity of 3.6 m/s. What velocity does he end up with after traveling 57 m downhill?

What is the net force along the direction of the displacement?

$$\sum F_s = mg \sin \theta + (-70\text{ N})$$



$$W = \sum F_s \times s = \frac{1}{2} m v^2 - \frac{1}{2} m v_0^2$$

$$[mg \sin \theta + (-70\text{ N})] \times s = \frac{1}{2} m v^2 - \frac{1}{2} m v_0^2$$

From this, we can solve for v!


103 Phys

Dr. Abdallah M. Azzeer

18

Sample Problem

A 61.0 kg bungee-cord jumper is on a bridge 45.0 m above a river. The elastic bungee cord has a relaxed length of $L = 25.0$ m. Assume that the cord obeys Hooke's law, with a spring constant of 160 N/m. If the jumper stops before reaching the water, what is the height h of her feet above the water at her lowest point?



$$\Delta K + \Delta U_e + \Delta U_g = 0$$

$$\Delta U_g = mg \Delta y = -mg(L + d)$$

$$\Delta U_e = \frac{1}{2}kd^2$$

103 Phys

Dr. Abdallah M. Azzeer

19

$$0 + \frac{1}{2}kd^2 - mg(L + d) = 0$$

$$\frac{1}{2}kd^2 - mgL - mgd = 0$$

$$\frac{1}{2}(160 \text{ N/m})d^2 - (61.0 \text{ kg})(9.8 \text{ m/s}^2)(25.0 \text{ m})$$

$$-(61.0 \text{ kg})(9.8 \text{ m/s}^2)d = 0$$

$$d = 17.9 \text{ m}$$

$$h = 45.0 \text{ m} - 42.9 \text{ m} = 2.1 \text{ m}$$


103 Phys

Dr. Abdallah M. Azzeer

20

Sample Problem

In Fig., a 2.0 kg package of tamale slides along a floor with speed $v_1 = 4.0$ m/s. It then runs into and compresses a spring, until the package momentarily stops. Its path to the initially relaxed spring is frictionless, but as it compresses the spring, a kinetic frictional force from the floor, of magnitude 15 N, acts on it. The spring constant is 10,000 N/m. By what distance d is the spring compressed when the package stops?



103 Phys

Dr. Abdallah M. Azzeer

21

SOLUTION:

$$E_{\text{mec},2} = E_{\text{mec},1} - \Delta E_{\text{th}}$$

$$E_{\text{mec},1} = K_1 + U_1 = \frac{1}{2}mv_1^2 + 0$$

$$E_{\text{mec},2} = K_2 + U_2 = 0 + \frac{1}{2}kd^2$$

$$\frac{1}{2}kd^2 = \frac{1}{2}mv_1^2 - f_k d$$

$$5000 d^2 + 15 d - 16 = 0$$

$$d = 0.055 \text{ m} = 5.5 \text{ cm}$$

103 Phys

Dr. Abdallah M. Azzeer

22

8.5 Relationship Between Conservative Forces and Potential Energy

A conservative force does not depend on the path

The work depends only on the initial and final coordinates.

► **Potential energy function U** such that the work done by a conservative force equals the decrease in the potential energy of the system.

The work done by a conservative force F

$$W_c = \int_{x_i}^{x_f} F_x dx = -\Delta U$$

where F_x is the component of F in the direction of the displacement.

the work done by a conservative force acting equals the negative of the change in the potential energy associated with that force

103 phys

Dr. Abdallah M. Azzeer

23

Where $\Delta U = U_f - U_i$

$$\Delta U = U_f - U_i = - \int_{x_i}^{x_f} F_x dx$$

Therefore, ΔU is **negative** when F_x and dx are in the same direction, as when an object is lowered in a gravitational field or when a spring pushes an object toward equilibrium.

The term **potential energy** implies that the system has the potential, or capability, of either gaining kinetic energy or doing work when it is released under the influence of a conservative force exerted on an object by some other member of the system.

We can then define the potential energy function as

$$U_f = - \int_{x_i}^{x_f} F_x dx + U_i$$

Physics

Dr. Abdallah M. Azzeer

24

- The value of U_i is often taken to be zero for the reference configuration. It really does not matter what value we assign to U_i because any nonzero value merely shifts $U_f(\mathbf{x})$ by a constant amount and only the *change* in potential energy is physically meaningful.

$$dU = -F_x dx$$

Therefore, the conservative force is related to the potential energy function through the relationship,

$$F_x = -\frac{dU}{dx}$$

That is, the x component of a conservative force acting on an object within a system equals the negative derivative of the potential energy of the system with respect to x .

We can easily check this relationship for the two examples already discussed. In the case of the deformed spring, $U_s = \frac{1}{2}kx^2$, and therefore

$$F_s = -\frac{dU_s}{dx} = -\frac{d}{dx}\left(\frac{1}{2}kx^2\right) = -kx$$

which corresponds to the restoring force in the spring (Hooke's law). Because the gravitational potential energy function is $U_g = mgy$, it follows from Equation 8.18 that $F_g = -mg$ when we differentiate U_g with respect to y instead of x .

We now see that U is an important function because a conservative force can be derived from it. Furthermore, Equation 8.18 should clarify the fact that adding a constant to the potential energy is unimportant because the derivative of a constant is zero.

Quick Quiz 8.11 What does the slope of a graph of $U(x)$ versus x represent? (a) the magnitude of the force on the object (b) the negative of the magnitude of the force on the object (c) the x component of the force on the object (d) the negative of the x component of the force on the object.

Energy Loss in Automobile

Automobile uses only at 13% of its fuel to propel the vehicle.

| | | |
|-------------|--|--|
| Why? | 67% in the engine: 1. Incomplete burning 2. Heat 3. Sound | 16% in friction in mechanical parts 4% in operating other crucial parts such as oil and fuel pumps, etc |
|-------------|--|--|

13% used for balancing energy loss related to moving vehicle, like air resistance and road friction to tire, etc

Two frictional forces involved in moving vehicles $m_{car} = 1450kg$ Weight $=mg = 14200N$

Coefficient of Rolling Friction; $\mu=0.016$ $\mu n = \mu mg = 227N$

Air Drag $f_a = \frac{1}{2} D \rho A v^2 = \frac{1}{2} \times 0.5 \times 1.293 \times 2v^2 = 0.647v^2$ **Total Resistance** $f_t = f_r + f_a$

Total power to keep speed $v=26.8m/s=60mi/h$ $P = f_t v = (691N) \cdot 26.8 = 18.5kW$

Power to overcome each component of resistance $P_r = f_r v = (227) \cdot 26.8 = 6.08kW$

$P_a = f_a v = (464.7) \cdot 26.8 = 12.5kW$

103 PhysDr. Abdallah M. Azzeer27

K:\103 Phys LECTURES SLIDES\103Phys_Slides_T1Y3839\CH8Flash

K:\103 Phys LECTURES SLIDES\103Phys_Slides_T1Y3839\CH8Flash

K:\103 Phys LECTURES SLIDES\103Phys_Slides_T1Y3839\CH8Flash

K:\103 Phys LECTURES SLIDES\103Phys_Slides_T1Y3839\CH8Flash

K:\103 Phys LECTURES SLIDES\103Phys_Slides_T1Y3839\CH8Flash

K:\103 Phys LECTURES SLIDES\103Phys_Slides_T1Y3839\CH8Flash

K:\103 Phys LECTURES SLIDES\103Phys_Slides_T1Y3839\CH8Flash

K:\103 Phys LECTURES SLIDES\103Phys_Slides_T1Y3839\CH8Flash

PhysicsDr. Abdallah M. Azzeer28

SUMMARY

If zero, then mechanical energy is conserved!

$\Delta K = \frac{1}{2} m(v_f^2 - v_i^2)$

NOT very good for:

- 1) times
- 2) directions
- 3) accelerations

$\sum W_{nc} = \Delta K + \Delta U$

e.g. work by friction $f_k d$

e.g. $\Delta U_g = mg\Delta y$
 $\Delta U_{el} = \frac{1}{2} k \Delta x^2$


Dr. Abdallah M. Azzeer

PROBLEM (?)

Mass is dropped on spring. ...
How far does spring compress?

Spring PE:
 $\frac{1}{2} kx^2$

Gravitational PE decreases by mgx



KE + Gravitational PE:
 $\frac{1}{2} mv^2 + mgh$

$$\frac{1}{2} mv^2 + mgh = -mgx + \frac{1}{2} kx^2$$

Dr. Abdallah M. Azzeer