



Fundamentals of Organic Chemistry

CHEM 109

For Students of Health Colleges

Credit hrs.: (2+1)

King Saud University

College of Science, Chemistry Department

Learning Objectives



At the end of this chapter, students will be able to:


- know the difference in structure between alcohols, phenols and ethers.
- Know the different classes of alcohols.
- Know how to name alcohols, phenols and ethers using IUPAC method.
- Recognize the basic properties (structure, physical and chemical properties) of alcohols, phenols and ethers .
- Recognize the effect of hydrogen bonds on their physical properties.
- Recognize the acidic properties of alcohols and phenols.
- know the different methods for the preparation of alcohols, phenols and ethers .
- Know the chemical reactions of these compounds.

Alcohols, Phenols and Ethers

3

- **Alcohols, phenols and ethers** may be viewed as organic *derivatives of water*.
- **Alcohols and phenols** have a common functional group, *the hydroxyl group, -OH*.

H-O-H	R-O-H	Ar-O-H	R-O-R	R-O-Ar	Ar-O-Ar
Water	Alcohol	Phenol	Ether		




- **Alcohols** are compounds whose molecules have a hydroxyl group attached to a *saturated* carbon atom.
- **Phenols** are compounds that have a hydroxyl group attached directly to a *benzene ring*.
- **Ethers** are compounds whose molecules have an oxygen atom bonded to **two** carbon atom.

Classification of Alcohols and Ethers

4

- **Alcohols** are classified as **primary** (1°), **secondary** (2°), or **tertiary** (3°), depending on whether one, two, or three organic groups are connected to the hydroxyl-bearing carbon atom.



- **Methyl alcohol**, which is not strictly covered by this classification, is usually grouped with the primary alcohols.

- **Ethers are classified as**

- **Symmetrical ethers;**

When the organic groups attached to the oxygen are **identical**.


- **Unsymmetrical ethers (mixed ethers);**

When the organic groups attached to the oxygen are **different**.

Nomenclature of Alcohols

5


- The **common names** for the simplest alcohols consist of alkyl group attached to the hydroxyl function followed by the word alcohol: *Alkyl alcohol*.
- In the **IUPAC system**, alcohols are named according to the following rules.
 1. Select the **longest continuous carbon chain that contains the -OH group**.
Drop the **-e** ending of the parent alkane and replace it by the suffix **-ol**: *Alkanol*
 2. When **isomers are possible**, the chain is numbered so as to give the functional group (-OH) the **lowest possible number**.




methanol
(methyl alcohol)



ethanol
(ethyl alcohol)




1-propanol
(*n*-propyl alcohol)




2-propanol
(isopropyl alcohol)




1-butanol
(*n*-butyl alcohol)



2-butanol
(*sec*-butyl alcohol)



2-methyl-1-propanol
(isobutyl alcohol)




2-methyl-2-propanol
(*tert*-butyl alcohol)

Nomenclature of Alcohols


6

3. When alkyl side chains or other groups are present; they are named alphabetically and their positions are indicated by a number.

The position of the functional group ($-OH$) is always given the **lowest possible number** at the end of the name.




4-Ethyl-2-hexanol
(not 3-Ethyl-5-hexanol)




5-Chloro-2-methyl-1-heptanol
(not 3-Chloro-6-methyl-7-heptanol)

For cyclic alcohols, numbering always starts from the carbon bearing the $-OH$ group.



3-Methylcyclohexanol
(not 1-Methyl-3-cyclohexanol)







3-Bromo-2-phenylcyclopentanol
(not 1-Bromo-2-phenyl-3-cyclopentanol)

Nomenclature of Alcohols



4. With Unsaturated Alcohols; If a molecule contains both an -OH group and a C=C or C-C triple bond, the -OH group takes preference before the double or triple bonds in getting the lower number.

The name should include (if possible) both the hydroxyl and the unsaturated groups, *even if this does not make the longest chain the parent hydrocarbon.*



Nomenclature of Alcohols





8

Alcohols with More Than One Hydroxyl Group

➤ Compounds with two adjacent alcohol groups are called **glycols**.

The most important example is ethylene glycol.

➤ Compounds with more than two hydroxyl groups are also known, and several, such as glycerol and sorbitol, are important commercial chemicals.








- **Ethylene glycol** is used as the “permanent” antifreeze in automobile radiators and as a raw material in the manufacture of Dacron.
- **Ethylene glycol** is completely miscible with water.
- **Glycerol** is a syrupy, colorless, water-soluble, high-boiling liquid with a distinctly sweet taste. Its soothing qualities make it useful in shaving and toilet soaps and in cough drops and syrups.



Nomenclature of Phenols

9


- **Phenols** are usually named as derivatives of the parent compounds.



- The hydroxyl group is named as a substituent when it occurs in the same molecule with carboxylic acid, aldehyde, or ketone functionalities, which have **priority in naming**.



but




Nomenclature of Ethers

10


Common Names

Ethers are usually named by giving the name of each alkyl or aryl group, in alphabetical order, followed by the word *ether*.


Methyl ether




Ethyl ether




Vinyl ether




Phenyl ether




Ethyl methyl ether




Ethyl-*n*-propyl ether



t-Butyl methyl ether




Methyl phenyl ether
(anisole)




IUPAC system


- For ethers with more complex structures, it may be necessary to name the **-OR** group as an **alkoxy** group.
- In the IUPAC system, the smaller alkoxy group is named as a substituent.



2-Methoxypentane



3-Ethoxyhexane



2-Methoxyethanol

Physical Properties of Alcohols and Ethers

11


○ Physical State

- The simplest **alcohol**, methanol, is a liquid at room temperature. In contrast, alkanes from methane to butane are gases.
- **Phenol** is a colorless, crystalline, and low-melting solid and other phenols also are solids, .
- **Ethers** are colorless compounds with characteristic, relatively pleasant odors.


○ Boiling Points

- **Ethers** have lower boiling points (bps) than **alcohols** with an equal number of carbon atoms.
- **Ether** has nearly the same b.p. as the corresponding hydrocarbon in which a $-\text{CH}_2-$ group replaces the ether's oxygen.


Because of their structures (no O-H bonds), ether molecules cannot form hydrogen bonds with one another.



Alkanes: No hydrogen bonding between molecules; low boiling points



Ethers: No hydrogen bonding between molecules; low boiling points



Alcohols: Hydrogen bonding between molecules; high boiling points


Compound	Formula	bp	mol wt	Water solubility (g/100 mL, 20°C)
1-butanol	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$	118°C	74	7.9
diethyl ether	$\text{CH}_3\text{CH}_2-\text{O}-\text{CH}_2\text{CH}_3$	35°C	74	7.5
pentane	$\text{CH}_3\text{CH}_2-\text{CH}_2-\text{CH}_2\text{CH}_3$	36°C	72	0.03

Physical Properties of Alcohols and Ethers



○ Boiling Points

- *Series of normal alcohols*; The boiling points increase with increasing molecular weights.
- A comparison of boiling points among *isomeric alcohols*; The boiling points decrease as the number of alkyl branches from the carbinol group increases.



- **Phenol** and most other phenols have high boiling points.


Physical Properties of Alcohols and Ethers

13


○ Solubility

- The lower *alcohols* are completely miscible with water.
- As the number of carbons in the alcohol increases, the solubility in water decreases.
- Low-molecular-weight ethers, such as dimethyl ether, are quite soluble in water.


Ether molecules can form hydrogen bonds to water.



Alkanes: No hydrogen bonding with water; insoluble



Ethers: Hydrogen bonding with water; soluble



Alcohols: Hydrogen bonding with water; soluble

Structure	Name	Mol.wt.	Bp (°C)	Solubility in H ₂ O At 20 °C
CH ₃ CH ₂ CH ₃	propane	44	-42	insoluble
CH ₃ OCH ₃	methyl ether	46	-24	soluble
CH ₃ CH ₂ OH	ethanol	46	78	soluble
CH ₃ CH ₂ CH ₂ CH ₃	<i>n</i> -butane	58	-0.5	insoluble
CH ₃ CH ₂ OCH ₃	ethyl methyl ether	60	8	soluble
CH ₃ CH ₂ CH ₂ OH	1-propanol	60	97	soluble
CH ₃ (CH ₂) ₃ CH ₃	<i>n</i> -pentane	72	35	insoluble
CH ₃ CH ₂ OCH ₂ CH ₃	ethyl ether	74	36	7.5 g/100 g
CH ₃ (CH ₂) ₂ CH ₂ OH	1-butanol	74	118	7.9 g/100 g
CH ₃ (CH ₂) ₅ CH ₃	<i>n</i> -heptane	100	98	insoluble
CH ₃ (CH ₂) ₂ O(CH ₂) ₂ CH ₃	<i>n</i> -propyl ether	102	91	0.2 g/100 g
CH ₃ (CH ₂) ₄ CH ₂ OH	1-hexanol	102	157	0.6 g/100 g

- **Phenol** and most other phenols are slightly soluble in water .

Hydrogen Bonding in Alcohols and Ethers


14

- The **boiling points** (bp's) of alcohols are much higher than those of ethers or hydrocarbons with similar molecular weights.

	CH ₃ CH ₂ OH	CH ₃ OCH ₃	CH ₃ CH ₂ CH ₃
mol wt	46	46	44
bp	+78.5°C	-24°C	-42°C

Why? Because alcohols form hydrogen bonds with one another.

The O-H bond is polarized by the high electronegativity of the oxygen atom and places a partial positive charge on the hydrogen atom and a partial negative charge on the oxygen atom.




Two or more alcohol molecules thus become loosely bonded to one another through hydrogen bonds.

- Consequently, **alcohols** have relatively high boiling points because they must supply enough heat to break the hydrogen bonds before each molecule.
- Hydrogen bonds are weaker than ordinary covalent bonds.

Hydrogen Bonding in Alcohols and Ethers

15

- The lower molecular-weight alcohols and ethers can form H-bond with water molecules.
- This accounts for the complete miscibility of the lower alcohols and ethers with water.



- However, as the organic chain lengthens and the alcohol becomes relatively more hydrocarbon like, its water solubility decreases.

Table 7.1 Boiling Point and Water Solubility of Some Alcohols



Name	Formula	bp, °C	Solubility in H ₂ O g/100 g at 20°C
methanol	CH ₃ OH	65	completely miscible
ethanol	CH ₃ CH ₂ OH	78.5	completely miscible
1-propanol	CH ₃ CH ₂ CH ₂ OH	97	completely miscible
1-butanol	CH ₃ CH ₂ CH ₂ CH ₂ OH	117.7	7.9
1-pentanol	CH ₃ CH ₂ CH ₂ CH ₂ CH ₂ OH	137.9	2.7
1-hexanol	CH ₃ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ OH	155.8	0.59

The Acidity of Alcohols and Phenols

16

- Like water, alcohols and phenols are weak acids.


The hydroxyl group can act as a proton donor, and dissociation occurs in a manner similar to that for water



The Acidity of Alcohols and Phenols

17

- **Phenols are stronger acids than alcohols** mainly because the corresponding phenoxide ions are stabilized by resonance.




The negative charge of an **alkoxide ion** is concentrated on the oxygen atom, but the negative charge on a phenoxide ion can be delocalized to the ortho and para ring positions through resonance.

Because **phenoxide ions** are stabilized in this way, the equilibrium for their formation is more favorable than that for alkoxide ions

The Acidity of Alcohols and Phenols

18


- All **electron-withdrawing groups increase acidity** by stabilizing the conjugate base. **Electron-donating groups decrease acidity** because they destabilize the conjugate base.



The Acidity of Alcohols and Phenols

19

- **Alkoxides**, the conjugate bases of alcohols, can be prepared by the reaction of an alcohol with sodium or potassium metal.




- Treatment of alcohols with sodium hydroxide does not convert them to their alkoxides.

This is because alkoxides are stronger bases than hydroxide ion, so the reaction goes in the reverse direction.

Since alcohols are weaker acids than water, it is not possible to form the salt of an alcohol in aqueous alkaline solutions.

- Treatment of phenols with sodium hydroxide converts them to phenoxide ions.




Preparation of Alcohols


20

1. Hydration of Alkenes

a. Addition of water to a double bond in the presence of an *acid catalyst, H⁺*.




b. The addition follows *Markovnikov's rule*.



c. It is *not possible to prepare primary alcohols* except Ethanol.

2. Oxidation of Cycloalkenes


Alkenes react with alkaline potassium permanganate to form glycols.



Preparation of Alcohols



21

○ Nucleophilic Substitution of Alkyl Halide



○ Reduction of Ketones, and Aldehydes

Aldehydes and ketones are easily reduced to primary and secondary alcohols, respectively.






Preparation of Alcohols



○ Addition of Grignard's Reagent to Aldehydes and Ketones

General equations




Preparation of Phenols

○ *The Alkali Fusion of Sulfonates*

The alkali fusion of sulfonates involves the following steps;

1. **Sulfonation** of an aromatic ring.
2. **Melting (fusion)** of the aromatic sulfonic acid with sodium hydroxide to give a **phenoxide salt**.
3. **Acidification** of the phenoxide with HCl to produce the **phenol**.



Uses of Alcohols and Phenols




- Drinks – The “alcohol” in alcoholic drinks is simply ethanol.
- As a fuel – Ethanol burns to give carbon dioxide and water and can be used as a fuel in its own right, or in mixtures with petrol (gasoline).
- As a solvent – Ethanol is widely used as a solvent. It is relatively safe, and can be used to dissolve many organic compounds which are insoluble in water. It is used, for example, in many perfumes and cosmetics.
- Isopropanol; rubbing alcohol, rapid evaporation, Antiseptic, more toxic than ethanol, but induces vomiting and used for manufacture of acetone.
- Phenol is used for resins, and pharmaceuticals.

Preparation of Ethers




1) Dehydration of Alcohols

- It takes place in the presence of acid catalysts (H_2SO_4 , H_3PO_4) (intermolecular reaction)




- **Example;**

The most important commercial ether is diethyl ether. It is prepared from ethanol and sulfuric acid.



- When ethyl alcohol is dehydrated by **sulfuric acid at 180°C** , the dominant product is **ethylene**.




Preparation of Ethers




2) Williamson Synthesis

- This method has two steps;
 - 1) An alcohol is converted to its alkoxide by treatment with a reactive metal (sodium or potassium).



- 2) Displacement is carried out between the alkoxide and an alkyl halide.



- To obtain the best yields of mixed dialkyl ethers, we select a 1° rather than a 2° or 3° alkyl halide and react it with a sodium alkoxide
- To prepare an alkyl aryl ether, we must be careful not to pick a combination in which one of the reagents has a halogen directly attached to an aromatic ring.

Preparation of Ethers

27

○ Example 1; Preparation of *t*-butyl methyl ether, $(\text{CH}_3)_3\text{C-O-CH}_3$.


➤ In theory, this could be done by either of two reactions.

1. You could react sodium methoxide, $\text{CH}_3\text{O}^-\text{Na}^+$, with *t*-butyl chloride, $(\text{CH}_3)_3\text{C-Cl}$.

This combination leads to dehydrohalogenation to an alkene, an elimination reaction.

2. You could react sodium *t*-butoxide, $(\text{CH}_3)_3\text{C-O}^-\text{Na}^+$, with methyl chloride, CH_3Cl .

This route gives the desired ether by substitution.





Preparation of Ethers

28

Example 2; Assume you need to synthesize **methyl phenyl ether (anisole)**, $\text{CH}_3\text{-O-C}_6\text{H}_5$, by the Williamson method.

➤ *In theory, you could obtain anisole in either of two ways.*



Reactions of Alcohols, Phenols and Ethers




29

- **Alcohols** undergo two kinds of reactions:
 - Those that involve the breaking of the oxygen-hydrogen bond (CO-H).
 - Those that involve the rupture of the carbon-oxygen bond (C-OH).
- **Phenols** do not participate in reactions where the C-OH bond is broken.
- **Ethers** are quite stable compounds.
 - The **ether** linkage does not react with bases, reducing agents, oxidizing agents, or active metals.
 - *Ethers react only under strongly acidic conditions.*



A) Those that involve the breaking of the oxygen-hydrogen bond (CO-H).

1) Reactions of Alcohols and Phenols as Acids: Salt Formation.



Reactions of Alcohols



A) Those that involve the rupture of the carbon-oxygen bond (C-OH).


1) Nucleophilic Substitution Reaction; The Reaction of Alcohols with Hydrogen Halides: Alkyl Halides

Alcohols react with hydrogen halides (HCl, HBr and HI) to give alkyl halides.



2) Dehydration of Alcohols: Formation of Alkenes

Alcohols can be dehydrated by heating them with strong acid.




Reactions of Alcohols

32


B) Oxidation Reactions

Alcohols with at least one hydrogen attached to the hydroxyl-bearing carbon can be oxidized to carbonyl compounds.

- **Primary alcohols** give **aldehydes**, which may be further oxidized to **carboxylic acids**.



- **Primary alcohols**, oxidation can be stopped at aldehyde stage by special reagents, such as “**pyridinium chlorochromate (PCC)**”.




Reactions of Alcohols




B) Oxidation Reactions

- Primary alcohols yield aldehydes when treated with mild oxidizing agents such as hot metallic copper or CrO_3 in pyridine.



- Primary alcohols; when treated with stronger oxidizing agents, such as chromic acid, $\text{H}_2\text{Cr}_2\text{O}_7$, or neutral potassium permanganate, KMnO_4 , the intermediate aldehydes formed initially are oxidized further to carboxylic acids.




Reactions of Alcohols



B) Oxidation Reactions


- **Secondary alcohols**, when treated with *any of the oxidizing agents* mentioned previously, yield ketones.



- **Tertiary alcohols**, having no hydrogen atom on hydroxyl-bearing carbon, do not undergo oxidation.

Reactions of Phenols

- Halogenation takes place *without catalyst*.



➤ **The products depend on the solvent used.**


- In *aprotic solvents* (solvents that do not release protons) (CCl_4 , CS_2)-bromination gives a mixture of *o*- and *p*-bromophenol.
- In *protic solvents* (solvents that can release protons) (H_2O)-halogenation gives a trisubstituted phenol is produced.

Reactions of Ethers

36

○ Cleavage of Ethers by Hot Concentrated Acids


- When **ethers** are heated in concentrated acid solutions, the ether linkage is broken.



- The acids most often used in this reaction are HI, HBr, and HCl.
- If an excess of acid is present, the alcohol initially produced is converted into an alkyl halide by the reaction.



For example,



Uses of Ethers



- Ether is used as a mild anesthetic and as a solvent in industries
- It is used as an antiseptic to prevent infection when an injection is administered.
- Dimethyl ether is used as refrigerant and as solvent at low temperature.
- Diethyl ether is a common ingredient as an aesthesia in surgery.
- Diethyl ether is common solvent for oils, gums, resins etc.
- We use phenyl ether as a heat transfer medium because of its high boiling point.