

A Steroidal Na⁺/K⁺ ATPase Inhibitor Triggers Pro-apoptotic Signaling and Induces Apoptosis in Prostate and Lung Tumor Cells

Sabina Honisch^{1,†}, Saad Alkahtani^{2,3,‡}, Michalis Kounenidakis², Guilai Liu¹, Saud Alarif³, Hamad Al-Yahya³, Konstantinos Dimas⁴, Abdullah A. AlKahtane³, Kyriakos C. Prousis⁵, Bader Al-Dahmash³, Theodora Calogeropoulou⁵, Konstantinos Alevizopoulos⁶, Florian Lang¹ and Christos Stournaras^{1,2,*}

¹Department of Physiology, University of Tübingen, Gmelinstr. 5, 72076 Tübingen Germany; ²Department of Biochemistry, University of Crete Medical School, Voutes, 70013 Heraklion, Greece; ³Department of Zoology, Science College, King Saud University, Riyadh, Saudi Arabia; ⁴Laboratory of Pharmacology, Faculty of Medicine, University of Thessaly, Biopolis, 41500 Larissa, Greece; ⁵Institute of Biology, Medicinal Chemistry and Biotechnology, National Hellenic Research Foundation, 11635, Athens, Greece; ⁶Medexis SA, 14568, Kryoneri, Attica, Greece

Abstract: Recently we have reported potent anti-cancer actions of various steroidal Na⁺/K⁺ ATPase inhibitors in multiple cell lines. Furthermore, the most powerful compound identified in this study, the 3-[(R)-3-pyrrolidinylo]xime derivative (3-R-POD), was highly effective in various tumor cell lines *in vitro*, and exhibited significant tumor growth inhibition in prostate and lung xenografts *in vivo*. In the present study we have addressed the molecular mechanisms implicated in the anti-cancer actions of 3-R-POD. We report here that 3-R-POD induces strong apoptotic responses in A549 lung- and in DU145 prostate- cancer cells. These effects are accompanied by significant upregulation of caspase-3 activity. Focussing on A549 cells, we further demonstrate late downregulation of BCL-2- and upregulation of c-Fos- gene transcription. In addition, the steroidal Na⁺/K⁺ ATPase inhibitor induced late de-phosphorylation of Focal Adhesion Kinase (FAK) and activation of p38 MAPK. Our findings suggest that the steroidal Na⁺/K⁺ ATPase inhibitor 3-R-POD induces apoptosis, paralleled by altered BCL-2 and c-Fos gene transcription, inhibition of the pro-survival FAK signalling, up-regulation of the pro-apoptotic p38 MAPK pathway and stimulation of caspase-3 activity.

Keywords: Apoptosis, lung cancer, prostate cancer, signaling, steroidal Na⁺/K⁺ ATPase inhibitor.

1. INTRODUCTION

The sodium potassium pump (Na⁺/K⁺ ATPase) is a transmembrane protein composed of multiple isoforms of catalytic α , regulatory β and modulatory γ subunits. The ion transport function of Na⁺/K⁺ ATPase is mainly regulating ionic and osmotic balance in cells (reviewed in [1]). Recent experimental evidence points to a multi-functional role of Na⁺/K⁺ ATPase in various cellular systems. Interestingly, this enzyme has been linked as well to cancer therapy. Indeed, recent studies established aberrant expression of some of its subunits in various malignancies including prostate, lung, colorectal, renal and pancreatic cancer [2-8]. In particular, the alpha 1 subunit of the Na⁺/K⁺ ATPase has been reported to represent a novel target to combat glioblastoma, melanoma and non-small cell lung cancers [4-6, 8].

In a recent study, we have tested for the first time the anti-cancer activity of 17 steroidal Na⁺/K⁺ ATPase inhibitors previously characterized as novel cardiotherapeutic agents [9]. Our results showed strong anti-cancer potencies of most compounds *in vitro* and nominated the 3-[(R)-3-pyrrolidinylo]xime derivative (3-R-POD) as the most active agent in 12 different cancer cell lines; its activity peaked at GI₅₀/IC₅₀ and LC₅₀ values of about 10 and 100 nM respectively in proliferation assays. More importantly, the compound exhibited significant tumor growth inhibition and lack of toxicity in prostate and lung cancer xenografts. On the contrary, digoxin, the prototype cardiac Na⁺/K⁺ ATPase inhibitor used as a control, was highly toxic and exhibited no anti-cancer activity at the doses tested [9]. In agreement with these observations, 3-R-POD was at least 30 times more cytotoxic to cancer cells versus normal fibroblasts in comparison to digoxin [9].

In this study, we sought to further analyze the anti-cancer properties of the 3-R-POD derivative in prostate and lung cancer cells. We have specifically focused on a series of key regulators of proliferation and apoptosis such as caspase 3, BCL-2, c-fos, FAK and p38 MAPK as these are typically involved in cancer development and progression in multiple cell systems. Our results provide novel mechanistic insights induced by 3-R-POD-dependent Na⁺/K⁺ ATPase inhibition and collectively support further development of the compound as a novel anti-cancer agent.

2. MATERIALS AND METHODS

2.1. Chemical Synthesis and Cell Cultures

Compound 3-R-POD was synthesized in seven steps from DHEA as previously described [9]. DU145 prostate- and A549 non small cell lung- cancer cell lines were obtained from the American Type Culture Collection (Manassas, VA) or the National Cancer Institute (NCI), NIH (Bethesda, MD, USA) and were adapted to grow in RPMI 1640 (Gibco, USA) supplemented with 10% foetal bovine serum (PAA, Austria) and 1% of penicillin/streptomycin (PAA, Austria) in a 5% CO₂ humidified atmosphere at 37°C. For each experiment, cells were seeded at 2 x 10⁵ in 6 well plates and cultured overnight. After 2 x washing with PBS, 3-R-POD (500 nM) or DMSO in fresh medium was applied for the indicated time.

2.2. FACS analysis and Caspase 3 activity

To determine the apoptotic response we used the Annexin V Apoptosis Detection Kit (MabTag, Germany). For that cells were harvested from the 6 well plate by treatment with trypsin-EDTA (Sigma-Aldrich, USA) for 10 minutes and washed once with cell culture medium. After a further wash with PBS and centrifugation at 1600 RPM for 3 minutes at RT cells were suspended in 100 μ l binding buffer containing Annexin V-FITC and propidium iodide and incubated for 20 minutes in the dark at RT. Subsequently, the cells were washed once, resuspended in 200 μ l binding buffer

*Address correspondence to this author at the Department of Biochemistry, University of Crete Medical School, GR-71110, Heraklion, Greece; Tel/Fax: +302810 394563/394530; E-mail: cstourn@med.uoc.gr

[†]Contributed equally and thus share first authorship

and measured immediately using the BD FACS Calibur (BD Biosciences, USA).

Active caspase-3 was ascertained by CaspGlow Fluorescein Active Caspase-3 Staining Kit (BioVision, USA) according to the manufacturers instructions. Briefly, cells detached and washed once with cell culture medium were suspended in 300µl of complete DMEM (Gibco, USA) including 1µl of FITC conjugated inhibitor of active Caspase-3 (FITC-DEVD-FMK). After 1h incubation and two washes with supplied wash buffer, cells were resuspended in 300µl of wash buffer and analyzed by flow cytometry (BD FACS Calibur, BD sciences, USA).

2.3. Western Blotting

Protein expression levels were analyzed by Western blotting. Subsequent to incubation with 500nM 3-R-POD for the indicated time periods, A549 cells were washed twice with ice-cold PBS and suspended in ice-cold RIPA lysis buffer (Thermo Fisher Scientific, USA) containing PhosSTOP phosphatase and cOmplete protease inhibitors (Roche Diagnostics, Germany). The extracts were centrifuged at 15000 RPM for 15 min at 4 °C and the protein concentration of the supernatant was determined by the Bradford

assay (BioRad, USA). 40 µg of total proteins were boiled with Roti-Load sample buffer (Carl Roth, Germany) for 5 min at 95°C and separated in 10% SDS-PAGE. Proteins were transferred to a PVDF-membrane (Thermo Fisher Scientific, USA) and blocked for 1h at room temperature with 5% BSA (Carl Roth, Germany) in TBST. For immunostaining membranes were incubated overnight at 4 °C with antibodies directed against phospho-p38 MAPK (Thr180/Tyr182), p38 MAPK, phospho-FAK (Tyr397) and FAK (all received from Cell Signaling, USA) each diluted 1:1000 in 5% BSA. GAPDH (1:3000, Cell Signaling, USA) was used for a loading control. To detect the specific proteins membranes were incubated for 1h at RT with a 1:2000 dilution of anti-rabbit IgG conjugated to horseradish peroxidase (Cell Signaling, USA). After washing, bands were visualized using the ECL western blotting detection reagent (GE Healthcare, USA) and quantified by Quantity One Software (ChemiDoc XRS, Bio-Rad, USA).

2.4. Real-time PCR

Gene expression of BCL-2 and c-Fos was determined by real-time PCR. Total RNA was extracted from A549 cells with TriFast (Peqlab, Erlangen, Germany) according to the manufacturer's




Fig. (1). Early and late apoptosis of A549 lung cancer cells treated by 3-R-POD. **A.** Chemical structure of the steroidal Na⁺/K⁺ ATPase inhibitor 3-[(R)-3-pyrrolidinyloxy]oxime derivative (3-R-POD). **B.** Flow cytometry results after a 24h treatment in the absence (control) or presence of 500nM 3-R-POD or 1µl solvent (DMSO) following staining with FITC conjugated Annexin V and propidium iodide (PI). Presented are arithmetic means ± SEM (n=6) of the percentage gated A549 cells binding: on the left bar graph Annexin V but no propidium iodide (early apoptosis), on the right bar graph both Annexin V and propidium iodide (late apoptosis). *** (p < 0,001) indicates a significant difference to respective value of untreated control (unpaired t-test). **C.** Original dot-plots (PI/Annexin V) of a representative experiment demonstrating an increase of cell events in the lower right quadrant (early apoptosis) and upper right quadrant (late apoptosis) in 3-R-POD treated group.

instructions. Reverse transcription of 2.5 µg total RNA was performed using GoScript™ Reverse Transcription System (Promega, USA). Polymerase chain reaction (PCR) of the respective genes was set up in a total volume of 20 µl using 500 nM forward and reverse primer and 2x GoTaq® qPCR Master Mix SYBR Green (Promega, USA) according to the manufacturer's instructions. Cycling conditions were as follows: initial denaturation at 95 °C for 2 min, followed by 40 cycles of 95 °C for 15 sec, 59 °C for 15 sec and 72 °C for 30 sec. For the amplification the following primers was used (5'→3' orientation):

BCL-2 fw: TGGATGACTGAGTACCTGAACCG
 BCL-2 rev: TGAGCAGAGTCTTCAGAGACAGC
 FOS fw: AGACTCCTTCTCCAGCATGGG
 FOS rev: TCCTGTCATGGTCTTCACAACG
 GAPDH fw: TGAGTACGTCGTGGAGTCCACTG
 GAPDH rev: GGTGCTAAGCAGTTGGTGGTG

Analysis of a melting curve confirmed specificity of PCR products, while real-time PCR amplifications were performed on a CFX96 Real-Time System (Bio-Rad, USA). All experiments were done in duplicate. As described earlier [10] relative quantification of gene expression was achieved using the Δct method. The amount

of sample RNA was standardized by amplification of the house-keeping gene GAPDH.

2.5. Statistical Analysis

All results are presented as means ± SEM, while *n* represents the number of independent experiments. Significance was assessed using unpaired student's t-test or ANOVA as appropriate. *p*-values < 0.05 were considered statistically significant.

3. RESULTS

3.1. 3-R-POD Induces Strong Pro-apoptotic Responses and Caspase 3 Activation in A549 and DU145 Cancer Cells

In a previous study, using the SRB- and MTT-cell growth assays the steroidal Na⁺/K⁺ ATPase inhibitor 3-R-POD ((*E, Z*) 3-((*R*)-pyrrolidin-3-yloxyimino)-6-methylene-5α-hydroxy-androstane-17-one, (chemical structure shown in Fig. 1A) showed potent anti-cancer activity in a large number of tumor cell lines [9]. These effects were attributed to its potent inhibition on the Na⁺/K⁺ ATPase (IC₅₀: 53 nM) [9]. In the present study, we addressed the pro-apoptotic effects of the compound in A549 lung- and DU145 prostate- cancer cell lines by using FACS analysis. As shown in Fig. 1 B, C, strong early and late apoptosis of A549 cells became




Fig. (2). Early and late apoptosis of DU145 prostate cancer cells treated by 3-R-POD. A. Flow cytometry results after a 24h treatment in the absence (control) or presence of 500nM 3-R-POD or 1µl solvent (DMSO) following staining with FITC conjugated Annexin V and propidium iodide (PI). Presented are arithmetic means ± SEM (n=6) of the percentage gated DU145 cells binding on the left bar graph Annexin V but no propidium iodide (early apoptosis), on the right bar graph both Annexin V and propidium iodide (late apoptosis). *** (*p* < 0,001) indicates a significant difference to respective value of untreated control (unpaired t-test). B. Original dot-plots (PI/Annexin V) of a representative experiment demonstrating an increase of cell events in the lower right quadrant (early apoptosis) and upper right quadrant (late apoptosis) in 3-R-POD treated group.




Fig. (3). Caspase 3 activation of A549 lung- and DU145 prostate cancer cells treated by 3-R-POD. After 24h exposure in the absence (control) or presence of 500 nM 3-R-POD or 1 μ l solvent (DMSO), cells were stained with conjugated inhibitor of active Caspase-3 (FITC-DEVD-FMK) and measured by FACS. (A) Left graph showing arithmetic means \pm SEM (n=3) of the percentage gated A549 cells. (B) Right graph showing arithmetic means \pm SEM (n=3) of the percentage gated DU145 cells. *** (p<0,001) indicates a significant difference to respective value of untreated control (unpaired t-test).

evident upon treatment with 500 nM 3-R-POD for 24h. Very similar pro-apoptotic responses were also observed in DU145 cells treated with 500 nM 3-R-POD (Fig. 2A, B). In line with these findings, a significant activation of caspase 3 was observed in both cell treated for 24h with 500 nM 3-R-POD (Fig. 3A, B). These *in vitro* data fully support the notion that the strong anti cancer activity of this compound reported in prostate- and non-small cell lung-cancer animal efficacy studies [9] correlates with the potent pro-apoptotic responses demonstrated in both cell lines.

3.2. 3-R-POD Regulates BCL-2- and c-Fos- Gene Transcription in A549 Cancer Cells

To elucidate molecular targets that may be implicated in 3-R-POD-governed anti-cancer activity, we further tested whether this compound alters gene transcription of BCL-2, a member of the BCL-2 family of proteins that regulates anti-apoptotic responses [11]. As shown in Fig. 4A, RT-PCR analysis revealed a significant down-regulation of BCL-2 gene transcripts in A549 NSCLC cells treated with 500nM 3-R-POD. This effect was evident after a treatment of 8 hours, while in earlier time points gene transcription was not modified. We further addressed whether 3-R-POD may regulate c-Fos gene expression. In contrast to the classic concept that c-Fos is acting as an oncogene, several reports indicated that c-Fos expression may foster pro-apoptotic responses in various tumors, including ovarian and hepatocellular carcinomas [12-14]. Interestingly, RT-PCR analysis revealed that treatment of A549 cells with 500 nM 3-R-POD induced a dramatic increase in c-Fos gene transcription after 2 to 8 hours (Fig. 4B).

3.3. 3-R-POD Activates p38 MAPK in A549 Cancer Cells

p38 MAPK is a mediator of apoptosis in response to a number of cellular signals (for a recent review see [15]). Various transcriptional and posttranscriptional mechanisms, which can involve the regulation of apoptotic and/or survival pathways, control the p38 MAPK-initiated pro-apoptotic responses. Accordingly, we further addressed the involvement of p38 MAPK in 3-R-POD-induced apoptosis in A549 lung cancer cells. As shown in Fig. 5A, a significant increase in p38 MAPK phosphorylation became evident after long term (8 hours) treatment of A549 cells with 3-R-POD.




Fig. (4). Time dependent effects of 3-R-POD on BCL-2 and c-Fos gene transcription in A549 lung cancer cells. A549 cells were stimulated with 500 nM 3-R-POD for the indicated time periods. BCL-2 (A) and c-Fos (B) mRNA levels were analysed by quantitative real-time PCR. Bars indicate the mean values of $2^{-\Delta\Delta Ct}$ using GAPDH as housekeeping gene \pm SEM from n=4 independent experiments. *** (p<0,001) indicates a significant difference to respective value of untreated control (unpaired t-test).




Fig. (5). Time dependent effects of 3-R-POD on p38 MAPK activity in A549 lung cancer cells. A549 cells were stimulated with 500 nM 3-R-POD for the indicated time periods. Following cell lysis, equal amounts of total lysates were immunoblotted with a specific antibody. Protein abundance was analyzed by densitometry. A. Intensity of phospho-p38 MAPK (Thr180/Tyr182), normalized to the intensity of the corresponding total p38 MAPK band. Bars indicate the relative fold increases as arithmetic means \pm SEM from $n=4$ independent experiments. ** indicates $p<0,01$ to respective value of untreated control (unpaired t-test). Below, representative original western blots showing the protein abundance of phosphorylated (p, Thr180/Tyr182) p38 MAPK, total MAPK and respective GAPDH as loading control. B. Intensity of phospho-FAK (Tyr397) normalized to the intensity of the corresponding total FAK band. Bars indicate the relative fold increases as arithmetic means \pm SEM from $n=6$ independent experiments. * indicates $p<0,05$, *** indicates $p<0,001$ to respective value of untreated control (unpaired t-test). Below representative original western blots showing the protein abundance of phosphorylated (p, Tyr397) FAK, total FAK and respective GAPDH as loading control.

3.4. 3-R-POD Down-regulates FAK Phosphorylation in A549 Cancer Cells

We further analyzed the capacity of 3-R-POD to regulate the activity of key signaling kinases implicated in cell survival and growth. For this we addressed the phosphorylation state of the Focal Adhesion Kinase FAK. As previously described [16], FAK regulates maintenance of normal cell survival, while disruption of FAK signaling controls various cellular outcomes including loss of substrate adhesion and apoptotic responses [16]. As shown in Fig. 5B, the ratio of phosphorylated FAK to the total protein content dropped significantly after long term (8 hours) incubation of the cells with 500 nM 3-R-POD, indicating inactivation of this key signaling kinase.

4. DISCUSSION

Na^+/K^+ ATPase is an emerging target in anti-cancer therapy, currently exploited in the development of novel therapeutics (reviewed in [2-6, 8]). Analysis of the molecular mechanisms mediating the anti-cancer properties of known Na^+/K^+ ATPase inhibitors (e.g. sugar-containing, plant-derived inhibitors such as digoxin), defined effects of these inhibitors in specific proliferation, survival, metabolism, angiogenesis and cell attachment pathways in cancer cells; these included effects on multiple molecular targets such as Myc, NF- κ B, HIF-1, p53 and others (reviewed in [2-6, 8]).

Previous studies in our laboratory have focused on the characterization of the anti-cancer activity of a series of non-sugar containing, steroidal Na^+/K^+ ATPase inhibitors originally developed as novel cardiotherapeutic agents [17-21]. These studies reasonably correlated the potency of the observed Na^+/K^+ ATPase inhibition with anti-cancer activity in multiple cells [9]. Specifically, it

was shown that the presence of a methylene or a methoxyimino functionality at C6 in combination with a (*R*)-pyrrolidin-3-yloxyimino group at C3 were instrumental for the anticancer activity of this class of compounds. Thus, the 3-[(*R*)-3-pyrrolidinyl] oxime derivative (3-R-POD, Fig. 1A) was identified as a potent Na^+/K^+ ATPase inhibitor with strong anti-cancer action in prostate and lung cancer cells and their corresponding xenografts *in vivo* [9]. In the current study, we present novel mechanistic insights underlying the potent anti-cancer activity reported for 3-R-POD. Specifically, we show that treatment of DU145 prostate- and A549 NSLC- tumor cell lines with 500nM 3-R-POD induced: a) strong pro-apoptotic responses and b) significant activation of caspase 3. Furthermore and in line with these findings, the steroidal inhibitor was shown to: c) down-regulate BCL-2- and up-regulate c-Fos-gene transcription and d) inhibit FAK- and trigger p38 MAPK-activity in A549 NSCLC- cells. Although additional studies will be required to determine whether the reported effects on signalling pathways are directly linked to 3-R-POD-dependent Na^+/K^+ ATPase inhibition, our findings clearly expand the current knowledge in the field by linking, for the first time, several gene products and proteins involved in cell survival and apoptosis with Na^+/K^+ ATPase inhibition.

Apoptotic cell death is regulated by complex interactions between pro-survival and pro-apoptotic members of the B-cell lymphoma-2 (BCL-2) protein family [22]. BCL-2 apoptosis-related family members are involved in the regulation of cellular fate as a response to antineoplastic agents and treatments in tumors (for recent reviews see [23, 24]). BCL-2 family proteins have been considered as drug targets in tumor biology [25], while BCL-2 antagonists have been recently proposed for cancer therapy [26]. The downregulation of BCL-2 gene transcription upon treatment of

A549 NSCLC cells by 3-R-POD is in line with these reports and support the notion that the steroidal Na⁺/K⁺ ATPase inhibitor may be a potent agent implicated in BCL-2 apoptosis-related induction of cell death.

Although the classic concept for c-Fos activity is based on its oncogenic properties, several studies in recent years indicate that c-Fos may also have tumor-suppressor activity. Indeed, c-Fos was shown to act as a proapoptotic agent in tumor necrosis factor-related apoptosis-inducing ligand (TRAIL)-induced apoptosis in prostate cancer cells [27]. In addition, c-Fos has been proposed to act as a mediator of the c-myc-induced apoptotic signaling in hepatocellular carcinomas *via* the p38 mitogen-activated protein kinase pathway [13]. Moreover, it has been reported that loss of c-Fos expression was associated with tumor progression in epithelial ovarian carcinomas [14]. In another study, fenretinide (HPR) that exhibits preventive and therapeutic activity against ovarian tumors was reported to cause induction of c-Fos gene expression in ovarian carcinoma cells [12]. This study established a relationship between c-Fos induction and apoptosis by the finding of an increased apoptotic rate in cells overexpressing c-Fos [12]. Our results showing strong induction of c-Fos gene transcription in 3-R-POD-treated A549 NSCLC cells that undergo potent apoptosis fully support the postulated relationship between c-Fos induction and apoptosis. Although further studies are now required to identify the molecular link of c-Fos up-regulation with apoptotic signaling in lung cancer cells, our findings showing potent p38 MAPK activation imply potential involvement of c-Fos regulated MAPK apoptotic signaling in 3-R-POD-induced effects.

The regulation of p38 MAPK apoptotic function is controlled by different signaling pathways [28]. Classically, p38 MAPK is activated by the small Rho family GTPases Rac1 and Cdc42 [29] and can mediate apoptosis or cell survival, by mechanisms involving the regulation of the expression and/or activity of different members of the Bcl-2 family. Indeed, Bax a pro-apoptotic member of the Bcl-2 family can be regulated by p38 MAPK. For example, in cardiomyocytes and MEFs-derived cell lines, p38 MAPK controls apoptosis, induced by different stimuli, through up-regulation of the pro-apoptotic proteins Fas and Bax and down-regulation of the activity of ERKs and Akt pro-survival pathways [30-31]. In lung cancer cells, p38 MAPK was shown to mediate isothiocyanate-induced strong pro-apoptotic response [32], while a critical role of the p38-mediated signaling pathway in the determination of lung cancer cell's sensitivity to EF24, a fluorinated synthetic curcumin analog was recently reported [33]. Our findings showing that the steroid inhibitor 3-R-POD activates p38 MAPK signaling and down-regulates of BCL-2 gene transcription in A549 lung cancer cells are in line with these reports and provide additional experimental evidence for the involvement of the p38 MAPK pathway in apoptosis regulation in response to Na⁺/K⁺ ATPase inhibition in lung cancer cells.

FAK is a nonreceptor protein-tyrosine kinase mainly localized within focal adhesions. A wide variety of extracellular signals including growth factors, steroid hormones, cytokines and neuropeptides have been reported to phosphorylate and thus activate FAK [34-37]. FAK signaling in turn is further activating downstream molecules such as PI-3K/Akt [38, 16] and paxillin [39]. Subsequent studies over the past years have established that FAK is important in maintenance of normal cell survival and cell adhesion, while inhibition of FAK activity and signaling results in loss of substrate adhesion and apoptosis (for recent reviews see [40, 38, 16]). Thus FAK may represent a potential target molecule for drug discovery in human cancers (for recent reviews see [41, 42]). Our findings showing 3-R-POD-induced inhibition of FAK phosphorylation are in line with the observed potent apoptotic responses of prostate- and lung-cancer cells upon stimulation with the steroidal Na⁺/K⁺ ATPase inhibitor 3-R-POD and further support

the notion that inhibition of FAK activity and signaling comprises important anti-tumorigenic potential.

In conclusion, this study implicates for the first time a series of important regulators of cell survival and apoptosis with Na⁺/K⁺ ATPase inhibition by 3-R-POD in prostate and lung tumor cells. Moreover, our results, when taken into consideration with previous observations showing higher selectivity and therapeutic index versus digoxin and potent anti-tumor efficacy in prostate and lung xenografts [9], clearly support the favourable properties of 3-R-POD and warrant its further anti-cancer development. This is in line with previous observations by Gobbini and co-workers [20] showing that a family of steroidal inhibitors comprising 3-R-POD has 5-20 times higher cardiotherapeutic (inotropic) activity and significantly less toxicity (3-10 fold) in comparison to digoxin in guinea pig studies [20]. Future studies aiming to characterize additional effects of 3-R-POD in cells and assess its potential impact on non-pump related functions of the Na⁺/K⁺ ATPase are currently in progress.

AUTHORS' CONTRIBUTIONS

Design of Research/Study: K. Alevizopoulos, C. Stourmaras. Development of chemical methodology and synthesis: T. Calogeropoulou, K.C. Prousis. Performed research/study: S. Honisch, S. Alkahtani, M. Kounenidakis, S. Alarifi, G. Liu, H. Al-Yahya. Collection, analysis and interpretation of data: K. Dimas, A. Alkahtane, B. Al-Dahmash. Writing of the manuscript: F. Lang, K. Alevizopoulos, C. Stourmaras. Study supervision: F. Lang, C. Stourmaras

CONFLICT OF INTEREST

The authors declare that they have no competing interests towards any aspect of the work described in this paper.

ACKNOWLEDGEMENTS

Authors would like to thank Tanja Loch for the careful preparation of the manuscript. We are also grateful to Dr. E.A. Papakonstanti for noteworthy comments. This work was supported by grants from the University of Crete Research Committee (KA 3452 to CS), Medexis Biotech SA (to KA), Deutsche Forschungsgemeinschaft (SFB773 to FL) and the Deanship of Scientific Research at King Saud University (Research Group Project No. RGPVPP-018 to SA, SA, and CS).

LIST OF ABBREVIATIONS

3-R-POD	=	3-[(R)-3-pyrrolidinyl]oxime derivative
FAK	=	Focal Adhesion Kinase
Na ⁺ /K ⁺ ATPase	=	Sodium potassium pump
NSCLC	=	Non Small Cell Lung-Cancer
BCL-2	=	B-cell lymphoma-2
TRAIL	=	Tumor necrosis factor-related apoptosis-inducing ligand

REFERENCES

- Xie, Z.; Cai, T. Na⁺-K⁺-ATPase-mediated signal transduction: From protein interaction to cellular function. *Mol. Interv.* **2003**, *3* (3), 157-168.
- Akopyanz, N.S.; Broude, N.E.; Bekman, E.P.; Marzen, E.O.; Sverdllov, E.D. Tissue-specific expression of Na,K-ATPase beta-subunit. Does beta 2 expression correlate with tumorigenesis? *FEBS Lett.* **1991**, *289* (1), 8-10.
- Espineda, C.E.; Chang, J.H.; Twiss, J.; Rajasekaran, S.A.; Rajasekaran, A.K. Repression of Na,K-ATPase beta1-subunit by the transcription factor snail in carcinoma. *Mol. Biol. Cell*, **2004**, *15* (3), 1364-1373.
- Lefranc, F.; Mijatovic, T.; Kondo, Y.; Sauvage, S.; Roland, I.; Debeir, O.; Krstic, D.; Vasic, V.; Gailly, P.; Kondo, S.; Blanco, G.;

- Kiss, R. Targeting the alpha 1 subunit of the sodium pump to combat glioblastoma cells. *Neurosurgery*, **2008**, *62* (1), 211-221; Discussion 221-212.
- [5] Mathieu, V.; Pirker, C.; Martin de Lassalle, E.; Vernier, M.; Mijatovic, T.; DeNeve, N.; Gaussin, J.F.; Dehoux, M.; Lefranc, F.; Berger, W.; Kiss, R. The sodium pump alpha1 sub-unit: a disease progression-related target for metastatic melanoma treatment. *J. Cell. Mol. Med.*, **2009**, *13* (9B), 3960-3972.
- [6] Mijatovic, T.; Roland, I.; Van Quaquebeke, E.; Nilsson, B.; Mathieu, A.; Van Vynckt, F.; Darro, F.; Blanco, G.; Facchini, V.; Kiss, R. The alpha1 subunit of the sodium pump could represent a novel target to combat non-small cell lung cancers. *J. Pathol.*, **2007**, *212* (2), 170-179.
- [7] Mijatovic, T.; Van Quaquebeke, E.; Delest, B.; Debeir, O.; Darro, F.; Kiss, R. Cardiotonic steroids on the road to anti-cancer therapy. *Biochim. Biophys. Acta*, **2007**, *1776* (1), 32-57.
- [8] Sakai, H.; Suzuki, T.; Maeda, M.; Takahashi, Y.; Horikawa, N.; Minamimura, T.; Tsukada, K.; Takeguchi, N. Up-regulation of Na(+),K(+)-ATPase alpha 3-isoform and down-regulation of the alpha1-isoform in human colorectal cancer. *FEBS Lett.*, **2004**, *563* (1-3), 151-154.
- [9] Dimas, K.; Papadopoulou, N.; Baskakis, C.; Prousis, K.C.; Tsakos, M.; Alkahtani, S.; Honisch, S.; Lang, F.; Calogeropoulou, T.; Alevizopoulos, K.; Stourmaras, C. Steroidal Cardiac Na⁺/K⁺ ATPase Inhibitors Exhibit Strong Anti-Cancer Potential *in vitro* and in prostate and lung cancer xenografts *in vivo*. *Anticancer Agents Med. Chem.*, **2014**, *14*(5), 762-70.
- [10] Pfaffl, M. W. A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res.*, **2001**, *29* (9), e45.
- [11] Shortt, J.; Johnstone, R. W. Oncogenes in cell survival and cell death. *Cold Spring Harb. Perspect Biol.*, **2012**, *4* (12).
- [12] Appierto, V.; Villani, M.G.; Cavadini, E.; Lotan, R.; Vinson, C.; Formelli, F. Involvement of c-Fos in fenretinide-induced apoptosis in human ovarian carcinoma cells. *Cell Death Differ.*, **2004**, *11* (3), 270-279.
- [13] Kalra, N.; Kumar, V. c-Fos is a mediator of the c-myc-induced apoptotic signaling in serum-deprived hepatoma cells *via* the p38 mitogen-activated protein kinase pathway. *J. Biol. Chem.*, **2004**, *279* (24), 25313-25319.
- [14] Mahner, S.; Baasch, C.; Schwarz, J.; Hein, S.; Wolber, L.; Janicke, F.; Milde-Langosch, K. C-Fos expression is a molecular predictor of progression and survival in epithelial ovarian carcinoma. *Br. J. Cancer*, **2008**, *99* (8), 1269-1275.
- [15] Wagner, E.F.; Nebreda, A.R. Signal integration by JNK and p38 MAPK pathways in cancer development. *Nat. Rev. Cancer*, **2009**, *9* (8), 537-549.
- [16] Lu, Q.; Rounds, S. Focal adhesion kinase and endothelial cell apoptosis. *Microvasc. Res.*, **2012**, *83* (1), 56-63.
- [17] Cerri, A. F. G.; Benicchio, A.; Bianchi, G.; Ferrari, P.; Gobbini, M.; Micheletti, R.; Pozzi, M.; Scotti, P.E. *Azaheterocyclyl derivatives of androstanes and androstenes as medicaments for cardiovascular disorders. WO Patent 2007118830 A1 2007.*
- [18] De Munari, S.; Cerri, A.; Gobbini, M.; Almirante, N.; Banfi, L.; Carzana, G.; Ferrari, P.; Marazzi, G.; Micheletti, R.; Schiavone, A.; Sputore, S.; Torri, M.; Zappavigna, M. P.; Melloni, P. Structure-based design and synthesis of novel potent Na⁺,K⁺ -ATPase inhibitors derived from a 5alpha,14alpha-androstane scaffold as positive inotropic compounds. *J. Med. Chem.*, **2003**, *46* (17), 3644-3654.
- [19] Gobbini, M.; Armaroli, S.; Banfi, L.; Benicchio, A.; Carzana, G.; Fedrizzi, G.; Ferrari, P.; Giacalone, G.; Giubileo, M.; Marazzi, G.; Micheletti, R.; Moro, B.; Pozzi, M.; Scotti, P.E.; Torri, M.; Cerri, A. Novel analogues of istaroxime, a potent inhibitor of Na⁺,K⁺-ATPase: Synthesis and structure-activity relationship. *J. Med. Chem.*, **2008**, *51* (15), 4601-4608.
- [20] Gobbini, M.; Armaroli, S.; Banfi, L.; Benicchio, A.; Carzana, G.; Ferrari, P.; Giacalone, G.; Marazzi, G.; Moro, B.; Micheletti, R.; Sputore, S.; Torri, M.; Zappavigna, M.P.; Cerri, A. Novel analogues of Istaroxime, a potent inhibitor of Na(+),K(+)-ATPase: Synthesis, structure-activity relationship and 3D-quantitative structure-activity relationship of derivatives at position 6 on the androstane scaffold. *Bioorg. Med. Chem.*, **2010**, *18* (12), 4275-4299.
- [21] Micheletti, R.; Mattered, G.G.; Rocchetti, M.; Schiavone, A.; Loi, M.F.; Zaza, A.; Gagnol, R.J.; De Munari, S.; Melloni, P.; Carminati, P.; Bianchi, G.; Ferrari, P. Pharmacological profile of the novel inotropic agent (E,Z)-3-((2-aminoethoxy)imino)androstane-6,17-dione hydrochloride (PST2744). *J. Pharmacol. Exp. Ther.*, **2002**, *303* (2), 592-600.
- [22] Kelly, P. N.; Strasser, A. The role of Bcl-2 and its pro-survival relatives in tumorigenesis and cancer therapy. *Cell Death Differ.*, **2011**, *18* (9), 1414-1424.
- [23] Frenzel, A.; Grespi, F.; Chmielewski, W.; Villunger, A. Bcl2 family proteins in carcinogenesis and the treatment of cancer. *Apoptosis*, **2009**, *14* (4), 584-596.
- [24] Kontos, C.K.; Christodoulou, M. I.; Scorilas, A. Apoptosis-related BCL2-family members: Key players in chemotherapy. *Anticancer Agents Med. Chem.*, **2014**, *14* (3), 353-374.
- [25] Christodoulou, M.I.; Kontos, C.K.; Halabalaki, M.; Skaltsounis, A. L.; Scorilas, A. Nature Promises New Anticancer Agents: Interplay with the Apoptosis-related BCL2 Gene Family. *Anticancer Agents Med. Chem.*, **2014**, *14* (3), 375-399.
- [26] Lessene, G.; Czabotar, P.E.; Colman, P.M. BCL-2 family antagonists for cancer therapy. *Nat. Rev. Drug Discov.*, **2008**, *7* (12), 989-1000.
- [27] Zhang, X.; Zhang, L.; Huang, X.; Otu, H.; Libermann, T. A.; DeWolf, W. C.; Khosravi-Far, R.; Olumi, A. F. c-Fos as a proapoptotic agent in TRAIL-induced apoptosis in prostate cancer cells. *Cancer Res.*, **2007**, *67* (19), 9425-9434.
- [28] Wada, T.; Penninger, J.M. Mitogen-activated protein kinases in apoptosis regulation. *Oncogene*, **2004**, *23* (16), 2838-2849.
- [29] Nobes, C.D.; Hall, A. Rho, rac and cdc42 GTPases: regulators of actin structures, cell adhesion and motility. *Biochem. Soc. Trans.*, **1995**, *23* (3), 456-459.
- [30] Porras, A.; Zuluaga, S.; Black, E.; Valladares, A.; Alvarez, A.M.; Ambrosino, C.; Benito, M.; Nebreda, A.R. P38 alpha mitogen-activated protein kinase sensitizes cells to apoptosis induced by different stimuli. *Mol. Biol. Cell.*, **2004**, *15* (2), 922-933.
- [31] Zuluaga, S.; Alvarez-Barrientos, A.; Gutierrez-Uzquiza, A.; Benito, M.; Nebreda, A.R.; Porras, A. Negative regulation of Akt activity by p38alpha MAP kinase in cardiomyocytes involves membrane localization of PP2A through interaction with caveolin-1. *Cell. Signal.*, **2007**, *19* (1), 62-74.
- [32] Yan, H.; Zhu, Y.; Liu, B.; Wu, H.; Li, Y.; Wu, X.; Zhou, Q.; Xu, K. Mitogen-activated protein kinase mediates the apoptosis of highly metastatic human non-small cell lung cancer cells induced by isothiocyanates. *Br. J. Nutr.*, **2011**, *106* (12), 1779-1791.
- [33] Thomas, S.L.; Zhao, J.; Li, Z.; Lou, B.; Du, Y.; Purcell, J.; Snyder, J. P.; Khuri, F. R.; Liotta, D.; Fu, H. Activation of the p38 pathway by a novel monoketone curcumin analog, EF24, suggests a potential combination strategy. *Biochem. Pharmacol.*, **2010**, *80* (9), 1309-1316.
- [34] Gu, S.; Kounenidakis, M.; Schmidt, E.M.; Deshpande, D.; Alkahtani, S.; Alarifi, S.; Foller, M.; Alevizopoulos, K.; Lang, F.; Stourmaras, C. Rapid activation of FAK/mTOR/p70S6K/PAK1-signaling controls the early testosterone-induced actin reorganization in colon cancer cells. *Cell. Signal.*, **2013**, *25* (1), 66-73.
- [35] Kallergi, G.; Agelaki, S.; Markomanolaki, H.; Georgoulas, V.; Stourmaras, C. Activation of FAK/PI3K/Rac1 signaling controls actin reorganization and inhibits cell motility in human cancer cells. *Cell. Physiol. Biochem.*, **2007**, *20* (6), 977-986.
- [36] Koukouritaki, S.B.; Gravanis, A.; Stourmaras, C. Tyrosine phosphorylation of focal adhesion kinase and paxillin regulates the signaling mechanism of the rapid nongenomic action of dexamethasone on actin cytoskeleton. *Mol. Med.*, **1999**, *5* (11), 731-742.
- [37] Papakonstanti, E.A.; Kampa, M.; Castanas, E.; Stourmaras, C.A. rapid, nongenomic, signaling pathway regulates the actin reorganization induced by activation of membrane testosterone receptors. *Mol. Endocrinol.*, **2003**, *17* (5), 870-881.
- [38] Hanks, S.K.; Ryzhova, L.; Shin, N.Y.; Brabek, J. Focal adhesion kinase signaling activities and their implications in the control of cell survival and motility. *Front. Biosci.*, **2003**, *8*, d982-996.

- [39] Jimenez, C.; Portela, R.A.; Mellado, M.; Rodriguez-Frade, J.M.; Collard, J.; Serrano, A.; Martinez, A.C.; Avila, J.; Carrera, A.C. Role of the PI3K regulatory subunit in the control of actin organization and cell migration. *J. Cell. Biol.*, **2000**, *151* (2), 249-262.
- [40] Golubovskaya, V.M.; Cance, W.G. FAK and p53 protein interactions. *Anticancer Agents Med. Chem.*, **2011**, *11* (7), 617-619.
- [41] Han, E.K.; McGonigal, T. Role of focal adhesion kinase in human cancer: a potential target for drug discovery. *Anticancer Agents Med. Chem.*, **2007**, *7* (6), 681-684.
- [42] Yin, B. Focal adhesion kinase as a target in the treatment of hematological malignancies. *Leuk. Res.*, **2011**, *35* (10), 1416-1418.

Received: April 11, 2014

Revised: June 10, 2014

Accepted: June 16, 2014