




Determination of caffeine content in tea and soft drink

Caffeine:

- Caffeine, the common name for 1,3,7-trimethylxanthine.
- It belongs to a group of methylxanthene.



Sources of caffeine:

- Caffeine is a chemical that is **found naturally** in the **leaves** and **seeds** of various plants.
- Natural sources of caffeine include coffee beans, cocoa beans, kola nuts, tea leaves and fruits of more than 60 plants.
- **Tea** leaves contains **1.5% to 3.5%** caffeine.
- Roasted coffee beans contain 0.75% to 1.5% caffeine.
- Cocoa bean contains 0.03% to 1.7% caffeine.
- Caffeine can be **added** to energy drinks and some carbonated drinks and **drug** products.
- Various carbonated beverages contain caffeine in the amount 30 to 60 mg per 355 ml.





250 ml
Filter Coffee

90 mg

Range: 69–127 mg



250 ml
Black Tea

63 mg


Range: 26–116 mg



250 ml
Instant Coffee

79 mg

Range: 63–90 mg



100 g
Dark Chocolate Bar

71 mg

Range: 18–123mg



250 ml
Cocoa Beverage

6 mg

Range: 3–34 mg



355 ml
Cola Drink

40 mg

Range: 30–60 mg

The effect of caffeine:

- Caffeine's main effect on your body is to **make you feel more awake and alert for a while**, but it can also cause problems.
- Many studies confirm caffeine's (**if it consumed properly**) ability to enhance mood and, exercise performance, the speed at which information is processed, awareness, attention, and reaction time.
- **Non proper consuming of caffeine can make you** shaky, make it hard to fall asleep, your heart beat faster, raise your blood pressure, cause headaches, nervousness, In massive doses, caffeine is lethal.
- A **fatal dose of caffeine** is more than 10 grams (about 170 mg/kg body weight).

- **Caffeine is classified as a central nervous system stimulant :**


- 1- An increase in heart rate.

- 2- Constriction of blood vessels.

- 3- Relaxed air passages to improve breathing.

- 4- ease of muscle contraction.

Mechanism of action:



- **Adenosine** is a central nervous system **neuromodulator** that has specific receptors.
- When adenosine binds to its receptors, neural activity slows down, and you feel sleepy.
→ Adenosine thus facilitates sleep and dilates the blood vessels (**opposite action**).
- Caffeine acts as an **adenosine-receptor antagonist**. This means that it binds to these same receptors, but **without reducing neural activity**.
- Fewer receptors are thus available to the natural “braking” action of adenosine, and neural activity therefore speeds up.
- **Caffeine** also causes the pituitary gland to secrete hormones that in turn cause the adrenal glands to produce more **adrenalin** so it increases your attention level and gives your entire system an extra burst of energy.

Practical Part

Objective:

- Determination of caffeine content in tea and soft drink using direct absorption of caffeine at 270 nm.

Principle:

- Even though caffeine is soluble in water, it is **more soluble in chloroform**.
→ Therefore, caffeine can be **extracted by chloroform** from the aqueous mixture using **Liquid-liquid extraction** involves the distribution of a substance between two immiscible liquid phases.
- Caffeine absorb light at 270 nm directly.
- Note:
This method will give a **general estimation** of caffeine concentration, it will **not give an accurate concentration** of caffeine in the sample. (why?)

Method:

First: Sample preparation:

1. 10 ml of (soft drink samples or hot water extract of tea samples) is taken in separating funnels, and 10 ml of chloroform was added to each sample.
2. The separating funnel should be shaken vigorously for 5 min while shaking, open the cover from time to time to release any pressure within the funnel. Be sure funnel is pointing away from you before opening.
3. The solutions then allowed to separate for 10 min at room temperature.
4. Only the lower chloroform layer will be collected for further analysis in a test tube or flask.
5. This chloroform layer will be diluted with pure chloroform (as shown in the table) appropriately to read absorbance.
6. Absorbance at 270 nm against pure chloroform as blank.

**Chloroform
layer (which
must be
collected)**



Method:

Second: Preparation of caffeine standard :

Tubes	Caffeine standard (100µg/ml)	Sample	Chloroform
S1	0.1	--	2.9
S2	0.2	---	2.8
S3	0.3	----	2.7
S4	0.4	----	2.6
S5	0.5	----	2.5
S6	0.6	----	2.4
S7	0.7	----	2.3
S8	0.8	----	2.2
Sample	----	(try different dilutions)	

Results:

Tubes	Absorbance at 270 nm	Caffeine Concentration µg/ml
S1		
S2		
S3		
S4		
S5		
S6		
S7		
S8		
Soft drink Sample		
Tea sample		

Calculations:

- **Concentration ($\mu\text{g/ml}$)** = conc. from curve x dilution factor