

**MOCK EXAMINATION (442 PDS)**

- 1. The electrical burn of the commissure is 24 hours old. What will happen to it in 5-10 days?**
  - a. infection
  - b. complete healing
  - c. eschar slough
  - d. increased pain
  
- 2. What is the recommended dose of this anesthetic in mg/lb or mg/kg?**
  - a. 8 mg/lb or 16 mg/kg
  - b. 16 mg/lb or 8 mg/kg
  - c. 2 mg/lb or 4 mg/kg
  - d. 7 mg/lb or 7 mg/kg
  
- 3. In Xylocaine overdose, the initial clinical finding is:**
  - a. CNS excitation
  - b. CNS depression
  - c. Airway loss
  - d. Syncope
  
- 4. What are two clinical signs of mandibular body fracture?**
  - a. broken teeth, luxated teeth
  - b. frenum tear, chin laceration
  - c. mucosal degloving, avulsed teeth
  - d. sublingual hematoma, vertical gingival tears
  
- 5. What is the most Common oral manifestation of pediatric HIV infection?**
  - a. candidiasis
  - b. kaposi's sarcoma
  - c. parotitis
  - d. herpetic gingivostomatitis
  
- 6. Hemangiomas should be treated with:**
  - a. observation
  - b. steroid therapy
  - c. sclerosing agents followed by surgical excision
  - d. laser therapy
  
- 7. Treatment for herpetic gingivostomatitis in the child should include all the following except:**
  - a. hydration
  - b. antipyretics
  - c. topical analgesics / anesthetics
  - d. antibiotics

**8. Most odontogenic infections are caused by:**

- a. gram positive cocci
- b. gram negative cocci
- c. gram positive rods
- d. gram negative rods

**9. Which of the following characteristics is true of the most recent generation of dentin bonding agents?**

- a. bonding is accomplished directly to the smear layer
- b. they are not affected by dentinal tubule structure
- c. they achieve bond strengths no greater than 3-5 MPa (megapascals)
- d. they use hydrophilic wetting agents in the primer

**10. Which of the following statements regarding infraoccluded teeth is FALSE?**

- a. ankylosed maxillary teeth tend to "submerged" more than mandibular teeth
- b. ankylosed primary first molars tend to submerged less than second molars in the same arch
- c. multiple ankylosed teeth occur less frequently than single ankylosed teeth
- d. ankylosed primary mandibular first molars often exfoliate on schedule