

Definition

A function f in two variables x and y , is said to be homogeneous of degree n , if

$$f(tx, ty) = t^n f(x, y)$$

for any real number t .

For example:

$f(x, y) = \sqrt[3]{x^2 + y^2}$ is homogenous of degree $n = \frac{2}{3}$

$f(x, y) = \frac{5xy^4}{x^2 + 7y^2}$ is homogenous of degree $n = 3$

$f(x, y) = \frac{1}{x^2 + 7y^2}$ is homogenous of degree $n = -2$

$f(x, y) = 3 - \frac{2x}{y} + e^{\frac{x^2}{3y^2}}$ is homogenous of degree $n = 0$

$f(x, y) = -8x \sin(xy^{-1})$ is homogenous of degree $n = 1$

While $g(x, y) = 1 + x + xy^3$ is not homogeneous.

Here the function g is a polynomial.

Remark: a polynomial in two variables is homogeneous if all its terms are of the same degree.

For example $g(x, y) = x^3 - 5xy^2 - 4x^2y$

Homogeneous Differential Equation

A first order DE

$$\frac{dy}{dx} = f(x, y)$$

is said to be homogeneous DE if the function f is homogeneous of degree zero.

Equivalently, the DE $M(x, y)dx + N(x, y)dy = 0$

is homogeneous if both M and N are homogeneous functions of the same degree.

A homogeneous DE can be reduced to a separable DE using the substitution

$$y = ux \text{ or } x = vy$$

Example 1

Solve the DE $xydy + (x^2 - 3y^2)dx = 0$.

This is a homogeneous DE.

$$\text{Let } y = ux \Rightarrow dy = udx + xdu$$

$$\Rightarrow ux^2(udx + xdu) + (x^2 - u^2x^2)dx = 0$$

$$\Rightarrow ux^3 du = -x^2 dx$$

$$\Rightarrow u du = -\frac{1}{x} dx$$

$$\Rightarrow \frac{1}{2} u^2 = -\ln x + c_1$$

Thus, the solution of the DE is given by

$$y^2 = -2x^2 \ln x + cx^2$$

Example 2

Solve the following differential equation.

$$x \frac{dy}{dx} - y = \sqrt{x^2 + y^2}$$

Solution

This is homogeneous DE

Thus, let $y = ux \Rightarrow dy = udx + xdu$

Hence, we have

$$xudx + x^2 du = (xu + x\sqrt{1+u^2}) dx$$

$$\Rightarrow \frac{1}{\sqrt{1+u^2}} du = \frac{1}{x} dx$$

Example 3

Solve the initial value problem

$$x \frac{dy}{dx} = y + xe^{\frac{y}{x}}, \quad y(1) = 1.$$

Solution

The DE is homogeneous, thus, let

$$y = ux \Rightarrow dy = udx + xdu$$

Hence, we have

$$xudx + x^2 du = (xu + xe^u) dx$$

$$\Rightarrow e^{-u} du = \frac{1}{x} dx$$

$$\Rightarrow -e^{-u} = \ln |x| + c$$

$$\Rightarrow -e^{\frac{-y}{x}} = \ln |x| + c$$

Since $y(1) = 1 \Rightarrow c = -e^{-1}$

Hence , the solution is

$$\Rightarrow -e^{\frac{-y}{x}} = \ln |x| - e^{-1}$$

Homework

Solve the following DE

$$\frac{dy}{dx} + \frac{x+y}{x-y} = 0$$

Differential Equations with linear coefficients

Consider the first order DE

$$\frac{dy}{dx} = \frac{ax+by+c}{hx+ky+l},$$

where a, b, c, h, k, l are constants and $c, l \neq 0$.

If $\frac{a}{h} = \frac{b}{k}$, then the above differential equation can be reduced to a

separable DE using the substitution $u = ax + by$ or $v = hx + ky$.

If $\frac{a}{h} \neq \frac{b}{k}$, then it can be converted to a homogeneous DE as

follows:

Put $ax + by + c = 0$

$$hx + ky + l = 0$$

and solve these two equations simultaneously,
assume that $x = x_0$ and $y = y_0$

Now, let

$$x = X + x_0 \text{ and } y = Y + y_0$$

$$\Rightarrow \frac{dy}{dx} = \frac{dY}{dX}$$

which will reduce the equation to a homogeneous DE, then it can be reduced to a separable DE.

Example 1

Solve the DE: $\frac{dy}{dx} = \frac{2+x+y}{1-2x-2y}$

Here $a = b = 1$, $h = k = -2$. Hence $\frac{a}{h} = \frac{b}{k} = \frac{1}{-2}$, therefore put

$$u = x + y \Rightarrow \frac{dy}{dx} = \frac{du}{dx} - 1$$

Hence the DE becomes

$$\frac{du}{dx} - 1 = \frac{2+u}{1-2u} \Rightarrow \frac{du}{dx} = \frac{3-u}{1-2u}$$

Which is separable DE.

Example 2

Solve the DE: $\frac{dy}{dx} = \frac{3+x+y}{1-x+y}$

Here $a = b = 1$, $h = -1$, $k = 1$. Hence $\frac{a}{h} \neq \frac{b}{k}$, therefore solve the two equations $3 + x + y = 0$ and $1 - x + y = 0$ to obtain $x = -1$, $y = -2$.

Now let $x = X + (-1)$ and $y = Y + (-2) \Rightarrow \frac{dy}{dx} = \frac{dY}{dX}$

Hence the DE becomes

$$\frac{dY}{dX} = \frac{X + Y}{-X + Y}$$

Which is a homogeneous DE.