



Department of Statistics & Operations Research
College of Science, King Saud University



STAT 145
Final Examination
Second Semester 1431 – 1432 H

		اسم الطالب
	رقم التحضير	الرقم الجامعي
	اسم الدكتور	رقم الشعبة

- Mobile Telephones are not allowed in the classrooms.
- Time allowed is **3 Hours**.
- Answer all questions.
- Choose the nearest number to your answer.
- For each question, put the code (**Capital Letters**) of the correct answer in the following table beneath the question number. **Do not use pencil or red pen.**

1	2	3	4	5	6	7	8	9	10
B	D	A	C	A	C	B	C	B	D
11	12	13	14	15	16	17	18	19	20
D	D	B	A	B	A	A	B	C	A
21	22	23	24	25	26	27	28	29	30
C	D	C	B	B	D	D	A	C	C
31	32	33	34	35	36	37	38	39	40
B	A	D	D	C	A	B	C	C	A
41	42	43	44	45	46	47	48	49	50
B	D	D	B	A	A	D	A	C	B

Term Marks	Final Exam. Marks	Total Marks

»»»

Following are the weights (in kg) for a sample of 6 children.

13, 20, 18, 12, 15, and 12.

(1) The mean of the data is:

A)	12	B)	<u>15</u>	C)	10	D)	18
----	----	----	-----------	----	----	----	----

(2) The median of the data is:

A)	17	B)	12	C)	10	D)	<u>14</u>
----	----	----	----	----	----	----	-----------

(3) The mode of the data is:

A)	<u>12</u>	B)	20	C)	15	D)	2
----	-----------	----	----	----	----	----	---

(4) The variance of the data is:

A)	3.347	B)	3.055	C)	<u>11.200</u>	D)	9.333
----	-------	----	-------	----	---------------	----	-------

(5) The coefficient of variation (C.V.) of the data is:

A)	<u>22.3%</u>	B)	17.4%	C)	74.7%	D)	62.22%
----	--------------	----	-------	----	-------	----	--------

 »»»

Temperatures recorded at 2 pm for 5 days of a year, for a city are:

7, 4, 0, -5, and 40.

(6) The range of temperatures is:

A)	33	B)	40	C)	<u>45</u>	D)	5
----	----	----	----	----	-----------	----	---

(7) The most suitable measure of centre for the data is:

A)	Mean	B)	<u>Median</u>	C)	Mode	D)	Range
----	------	----	---------------	----	------	----	-------

 »»»

Let A and B denote two events defined on the same sample space with $P(A) = 0.6$, $P(B) = 0.4$, and $P(A \cup B) = 0.74$, then:

(8) The events A and B are:

A)	independent	B)	mutually exclusive	C)	<u>dependent</u>	D)	impossible
----	-------------	----	--------------------	----	------------------	----	------------

(9) The $P(\bar{A} \cap \bar{B})$ is:

A)	0.18	B)	<u>0.26</u>	C)	0.50	D)	1.00
----	------	----	-------------	----	------	----	------

 »»»

Consider the following cumulative frequency distribution table for the ages of all workers in a certain factory.

Age	Cumulative frequency
26 - 35	10
36 - 45	40
46 - 55	50

(10) Percentage of workers in the age group 36 - 45 is:

A)	40%	B)	80%	C)	30%	D)	<u>60%</u>
----	-----	----	-----	----	-----	----	------------

(11) Number of workers having age 36 or more is:

A)	90	B)	50	C)	10	D)	<u>40</u>
----	----	----	----	----	----	----	-----------

(12) The true class limits for the first class are:

A)	26 - 35	B)	21.5 - 35.5	C)	25.5 - 34.5	D)	<u>25.5 - 35.5</u>
----	---------	----	-------------	----	-------------	----	--------------------

»»»

Let A and B be two independent events. Suppose that $P(A) = 0.6$ and $P(B) = 0.3$ then

(13) $P(\bar{A} \cap B)$ is equal to:

A)	0.08	B)	<u>0.12</u>	C)	0.20	D)	0.42
----	------	----	-------------	----	------	----	------

(14) $P(A \cup B)$ is equal to:

A)	<u>0.72</u>	B)	0.90	C)	0.10	D)	0.7
----	-------------	----	------	----	------	----	-----

»»»

Suppose that a town has 20% of men known to have a certain disease. A certain medical test is applied to randomly selected 500 men. The following data is obtained.

Test	Disease		Total
	Present	Absent	
Positive	82	80	162
Negative	38	300	338
Total	120	380	500

Let an individual be selected at random from the sample.

(15) The probability that the selected person has the disease is:

A)	0.20	B)	<u>0.24</u>	C)	0.68	D)	0.32
----	------	----	-------------	----	------	----	------

(16) The probability that the test gives a false negative result is:

A)	<u>0.32</u>	B)	0.68	C)	0.21	D)	0.79
----	-------------	----	------	----	------	----	------

(17) The sensitivity of the test is:

A)	<u>0.68</u>	B)	0.16	C)	0.51	D)	0.79
----	-------------	----	------	----	------	----	------

(18) Suppose that 20% of men in the town have the disease, the predictive probability negative for the test is:

A)	0.37	B)	<u>0.91</u>	C)	0.09	D)	0.89
----	------	----	-------------	----	------	----	------

»»»

In a large population of people, 34% have blood type A+. We choose randomly 8 persons from this population. Let X = the number of persons having blood type A+ among the 8.

(19) The values of the parameters of the distribution of X are:

A)	3 and 0.34	B)	8, and 0.66	C)	8 and 0.34	D)	8 and 34
----	------------	----	-------------	----	------------	----	----------

(20) The probability that there is exactly one person with blood type A+ is:

A)	<u>0.1484</u>	B)	0.0028	C)	0.3400	D)	0.0185
----	---------------	----	--------	----	--------	----	--------

(21) The probability that there is at least one person with blood type A+, is :

A)	0.1484	B)	0.1844	C)	<u>0.9640</u>	D)	0.0360
----	--------	----	--------	----	---------------	----	--------

»»»

The number of serious surgical operations that are performed in a hospital during a day follows a Poisson distribution with an average of 5 persons per day, then:

(22) The probability that no operations is performed in the next day is:

A)	0.99996	B)	0.08972	C)	0.54210	D)	<u>0.0067</u>
----	---------	----	---------	----	---------	----	---------------

(23) The probability that 5 operations are performed in the next day is:

A)	0.2145	B)	0.8521	C)	<u>0.175</u>	D)	0.5124
----	--------	----	--------	----	--------------	----	--------

(24) The average number of operations that are performed in two days is:

A)	20	B)	<u>10</u>	C)	5	D)	30
----	----	----	-----------	----	---	----	----

»»»

In a population of people, X = the body mass index (in kg/m^2) is normally distributed with mean $\mu = 25$ and standard deviation $\sigma = 2$. For a randomly chosen person,

(25) $P(24 < X < 26) =$

A)	0.6915	B)	<u>0.3830</u>	C)	0.2085	D)	1
----	--------	----	---------------	----	--------	----	---

(26) $P(X = 21) =$

A)	0.9772	B)	0.0228	C)	1	D)	<u>0</u>
----	--------	----	--------	----	---	----	----------

(27) The value of k such that $P(X > k) = 0.2578$, is:

A)	0.257	B)	25	C)	-0.65	D)	<u>26.3</u>
----	-------	----	----	----	-------	----	-------------

»»»

A sample of size 100 is taken from a population having a proportion $p_1 = 0.8$. Another independent sample of size 400 is taken from a population having a proportion $p_2 = 0.5$.

(28) The sampling distribution for the difference in sample proportions has a mean equal to:

A)	<u>0.3</u>	B)	1.3	C)	0	D)	0.8
----	------------	----	-----	----	---	----	-----

(29) The sampling distribution for the difference in sample proportions has a standard error equal to:

A)	0.015	B)	0.0022	C)	<u>0.047</u>	D)	0.1239
----	-------	----	--------	----	--------------	----	--------

(30) $P(\hat{p}_1 - \hat{p}_2 < 0.2) = :$

A)	0.4423	B)	0.993	C)	<u>0.0166</u>	D)	0.2415
----	--------	----	-------	----	---------------	----	--------

»»»

Suppose it has been established that for a certain type of clients, the average length of a home visit by a public health nurse is 45 minutes with a standard deviation of 15 minutes, and that for a second type of clients, the average home visit time is 30 minutes with a standard deviation of 20 minutes. If a nurse randomly visits 35 clients from the first population and 40 from the second population, then

(31) The mean of the difference between two sample means is:

A)	5	B)	<u>15</u>	C)	20	D)	35
----	---	----	-----------	----	----	----	----

(32) The standard error of the difference between two sample means is:

A)	<u>4.0532</u>	B)	16.4286	C)	8.2143	D)	0.5241
----	---------------	----	---------	----	--------	----	--------

(33) The probability that the average length of home visit will differ between the two groups by 20 or more is:

A)	0.8907	B)	0.4215	C)	0.5	D)	<u>0.1093</u>
----	--------	----	--------	----	-----	----	---------------

»»»

A researcher wishes to determine if vitamin E supplements could increase cognitive ability among elderly women. In 1999 the researcher recruits a sample of elderly women age 75-80. At the time of the enrollment into the study, the women were randomized to either take Vitamin E, or a placebo for six months. At the end of the six month period, the women were given a cognition test. Higher scores on this test indicate better cognition. The mean of the test scores of 81 women who took vitamin E supplements was $\bar{X}_1 = 27$, while the mean of the test scores of the 90 women who took placebo supplements was $\bar{X}_2 = 24$. Assuming the two populations follow approximately two different normal distributions with standard deviations, $\sigma_1 = 6.9$ and $\sigma_2 = 6.2$, respectively.

(34) The point estimate for the difference between the two population means ($\mu_1 - \mu_2$):

A)	27	B)	24	C)	6.2	D)	<u>3</u>
----	----	----	----	----	-----	----	----------

(35) The standard error for the difference between the two sample means is:

A)	6.9	B)	6.2	C)	<u>1.007</u>	D)	3
----	-----	----	-----	----	--------------	----	---

(36) A lower limit of a 95% confidence interval for the difference between the two population means is:

A)	<u>1.0263</u>	B)	4.9745	C)	5.9120	D)	1.2354
----	---------------	----	--------	----	--------	----	--------



»»»

Six healthy three year old female sheep were injected with the antibiotic Gentamicin, at a dosage of 10 mg/kg body weight. Their blood serum concentrations (mg/ml) of Gentamicin after injection were 33; 26; 34; 31; 23; 25, the summary statistics for these data are

n	mean	Standard deviation	SE of mean
6	28.67	4.59	1.87

Assuming the data follows approximately a normal distribution,

(37) At the 90% level of confidence, the reliability coefficient is:

A)	2.33	B)	<u>2.015</u>	C)	3.215	D)	1.96
----	------	----	--------------	----	-------	----	------

(38) The 90% confidence interval for the population mean score on this test is:

A)	(27.412, 30.145)	B)	(24.48, 29.10)	C)	<u>(24.902, 32.438)</u>	D)	(32.48, 39.55)
----	------------------	----	----------------	----	-------------------------	----	----------------

(39) The value of the test statistic, for testing the hypothesis $H_0 : \mu = 30$ vs $H_1 : \mu < 30$ is:

A)	-2.2587	B)	2.5812	C)	<u>-0.7112</u>	D)	3.3412
----	---------	----	--------	----	----------------	----	--------

(40) At the 5% significance level, the critical region is :

A)	<u>$(-\infty, -2.015)$</u>	B)	$(-2.015, 2.015)$	C)	$(2.015, \infty)$	D)	$(2.58, \infty)$
----	---------------------------------------	----	-------------------	----	-------------------	----	------------------

(41) At the 5% significance level, we are able to :

A)	Reject H_0	B)	<u>Can not reject H_0</u>	C)	Decision is not possible
----	--------------	----	--	----	--------------------------



»»»

A Biostatistician, found that among 2000 boys in the age group 7 to 12 years, 400 were overweight. On the basis of this study:

(42) The standard error of the sample proportion of the overweight boys in the age group 7 to 12 years is:

A)	0.0500	B)	0.0221	C)	0.6587	D)	<u>0.0089</u>
----	--------	----	--------	----	--------	----	---------------

(43) The 99% upper confidence limit for the population proportion of the overweight boys in the age group 7 to 12 years is:

A)	0.5000	B)	0.0221	C)	0.6587	D)	<u>0.223</u>
----	--------	----	--------	----	--------	----	--------------

(44) The value of the test statistic for testing the hypotheses the proportion of boys in the age group 7 to 12 years is 0.18, is:

A)	-2.2587	B)	<u>2.33</u>	C)	-0.7112	D)	3.3412
----	---------	----	-------------	----	---------	----	--------

(45) At the 5% significance level, can we conclude that more than 18% of boys in the age group 7 to 12 years are overweight:

A)	<u>Yes</u>	B)	No	C)	Decision is not possible
----	------------	----	----	----	--------------------------



»»»

A sample of 25 freshman nursing students made a mean score of 77 on a test designed to measure the attitude toward the dying patients. The sample standard deviation was 10. Assume that the data comes from a normal population.

(46) The statistical hypothesis for testing the hypothesis that the mean score is different than 80 is:

A)	$H_0 : \mu = 80$ vs $H_1 : \mu \neq 80$	B)	$H_0 : \mu = 80$ vs $H_1 : \mu < 80$
C)	$H_0 : \mu = 80$ vs $H_1 : \mu > 80$	D)	$H_0 : \mu = 77$ vs $H_1 : \mu < 77$

(47) The appropriate test statistic for testing the mean score is different than 80 is:

A)	$z = \frac{x - 80}{\sigma}$	B)	$z = \frac{\bar{x} - 80}{\sigma / \sqrt{n}}$
C)	$t = \frac{x - 80}{S / \sqrt{n}}$	D)	$t = \frac{\bar{x} - 80}{S / \sqrt{n}}$

(48) The value of the test statistic for the statistical hypothesis is:

A)	<u>-1.500</u>	B)	-2.025	C)	3.258	D)	0
----	---------------	----	--------	----	-------	----	---

(49) The test rejects, at 5% significance level, the hypothesis that the mean score is 80 if:

A)	$z > 1.96$	B)	$z < -1.96$ or $z > 1.96$
C)	<u>$t < -2.064$ or $t > 2.064$</u>	D)	$t > 1.71$

(50) At the 5% significance level we are able to :

A)	Reject H_0	B)	<u>Can not reject H_0</u>	C)	Decision is not possible
----	--------------	----	--	----	--------------------------