

Exercise 1 :

$$\text{a) } F'(x) = 4x^3 \ln(x) - 4x \ln(2x), \quad (1) + (1)$$

$$\text{b) } \int_0^5 \frac{dx}{\sqrt{1+x^4}} \approx \frac{1}{2}(3.284871) \approx 1.642435$$

k	x_k	$f(x_k)$	m	$mf(x_k)$
0	0	1	1	1
1	1	$\frac{1}{\sqrt{2}}$	2	$\sqrt{2} \approx 1.414213$
2	2	$\frac{1}{\sqrt{17}}$	2	$\frac{2}{\sqrt{17}} \approx 0.485071$
3	3	$\frac{1}{\sqrt{82}}$	2	$\frac{2}{\sqrt{82}} \approx 0.220863$
4	4	$\frac{1}{\sqrt{257}}$	2	$\frac{2}{\sqrt{257}} \approx 0.124756$
5	5	$\frac{1}{\sqrt{626}}$	1	$\frac{1}{\sqrt{626}} \approx 0.039968$
				3.284871

(1) correct formula
 (1.5) correct numbers
 (0.5) final answer

Exercise 2 :

a)

$$\begin{aligned} \int x 3^{2x^2} (3^{2x^2} + 1)^{-4} dx & \stackrel{u=3^{2x^2}+1}{=} \frac{1}{4 \ln 3} \int u^{-4} du \quad (2) \\ & = -\frac{1}{12 \ln 3} u^{-3} + c \\ & = -\frac{1}{12 \ln 3} (3^{2x^2} + 1)^{-3} + c. \quad (1) \end{aligned}$$

b)

$$\begin{aligned} \int \frac{(\log_2 x)^2 + \sqrt{x}}{x} dx & = \int \frac{\ln^2(x)}{x \ln^2 2} dx + \int x^{-\frac{1}{2}} dx \quad (1) \\ & = \frac{\ln^3(x)}{3 \ln^2 2} + 2\sqrt{x} + c. \quad (1)+(1) \end{aligned}$$

$$c) \int \frac{dx}{\sqrt{e^{4x} - 36}} \stackrel{6u=e^{2x}}{=} \frac{1}{12} \int \frac{du}{u\sqrt{u^2 - 1}} = \frac{1}{12} \sec^{-1}\left(\frac{e^{2x}}{6}\right) + c. \quad (1) + (1)$$

Exercise 3 :

a) By parts

$$\begin{aligned} \int \ln(x^2 + 1) dx & \stackrel{u=\ln(x^2+1), v'=1}{=} x \ln(x^2 + 1) - \int \frac{2x^2}{x^2 + 1} dx \quad (1.5) \\ & = x \ln(x^2 + 1) - 2x + 2 \tan^{-1}(x) + c. \quad (1.5) \end{aligned}$$

b)

$$\begin{aligned} \int \frac{dx}{x^3 \sqrt{x^2 - 1}} & \stackrel{x=\sec(\theta)}{=} \int \frac{\sec(\theta) \tan(\theta)}{\sec^3(\theta) \tan(\theta)} d\theta = \int \cos^2(\theta) d\theta \quad (1) \\ & = \frac{\theta}{2} + \frac{1}{2} \sin(\theta) \cos(\theta) + c \quad (1) \\ & = \frac{1}{2} \sec^{-1}(x) + \frac{\sqrt{x^2 - 1}}{2x^2} + c. \quad (1) \end{aligned}$$

$$c) \frac{f'(x)}{f(x)} = \frac{2}{x} + 3x^2 \ln(x^2 + 1) + \frac{2x^4 + 2x}{x^2 + 1} \quad (2)$$

$$f'(x) = \left(2x + 3x^4 \ln(x^2 + 1) + \frac{2x^3(x^3 + 1)}{x^2 + 1} \right) (x^2 + 1)^{(x^3 + 1)}. \quad (1)$$


Exercise 4 :

a)

$$\begin{aligned} \int \frac{3x - 1}{x^2 + 4x + 8} dx & = \frac{1}{2} \int \frac{3(2x + 4) - 14}{x^2 + 4x + 8} dx \quad (1) \\ & = \frac{3}{2} \ln|x^2 + 4x + 8| - 7 \int \frac{1}{(x + 2)^2 + 4} dx \quad (1) \\ & = \frac{3}{2} \ln|x^2 + 4x + 8| - \frac{7}{2} \tan^{-1}\left(\frac{x + 2}{2}\right) + c. \quad (1) \end{aligned}$$

b) The area of the region bounded by the curves $x = y^2$, $x + y = 6$, $y = -4$, $y = 2$ is

$$A = \int_{-3}^2 (6 - y - y^2) dy + \int_{-4}^{-3} (y^2 - 6 + y) dy = 23 + \frac{2}{3}. \quad (2)$$



c) $2x^2 = 8x \iff x(x - 4) = 0$ (0.5)

(i) $(x = 5), V = 2\pi \int_0^4 (5 - x)(8x - 2x^2) dx$, (1)

(ii) $(y = -1), V = \pi \int_0^4 ((1 + 8x)^2 - (1 + 2x^2)^2) dx$. (1.5)

Exercise 5 :

a)

$$SA = 2\pi \int_0^1 2y^3 \sqrt{1 + 36y^4} dy \quad (1)$$

$$\stackrel{u=1+36y^4}{=} \frac{\pi}{36} \int_1^{37} \sqrt{u} du \quad (1)$$

$$= \frac{\pi}{54} (\sqrt{37} - 1). \quad (1)$$

b)

$$A = \frac{1}{2} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} 1 - (1 - \cos(\theta))^2 d\theta \quad (1)$$

$$= \frac{1}{2} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \left(2 \cos(\theta) - \frac{1}{2} - \frac{1}{2} \cos(2\theta) \right) d\theta = 2 - \frac{\pi}{2}. \quad (2)$$

c)

$$L = \int_{-\pi}^{\pi} \sqrt{(1 + \cos \theta)^2 + \sin^2(\theta)} d\theta \quad (1)$$

$$= \int_{-\pi}^{\pi} 2 \cos\left(\frac{\theta}{2}\right) d\theta = 4 \sin\left(\frac{\theta}{2}\right) \Big|_{-\pi}^{\pi} = 8. \quad (2)$$