

# 1 $L^p$ Spaces

## 1 $L^p$ Spaces

$(X, \mathcal{A}, \mu)$  will be a fixed measure space.

### Definition 1.1

If  $0 < p < +\infty$ , we define the space  $\mathcal{L}^p(\mu)$  to be the set of all measurable functions  $f: X \rightarrow \bar{\mathbb{R}}$  such that  $\int_X |f|^p d\mu < \infty$ .

We Remark that the space  $\mathcal{L}^1(\mu)$  is the set of all functions which are integrable over  $X$  with respect to the measure  $\mu$ .

If  $\mu$  is the counting measure on a countable set  $X$ , then  $\int_X f(x) d\mu(x) = \sum_{x \in X} f(x)$ .

Then  $\mathcal{L}^p$  is usually denoted  $\ell^p$ , the set of sequences  $(x_n)_n$  such that  $\sum_{n=1}^{+\infty} |x_n|^p < +\infty$ .

### Definition 1.2

We define the relation  $\sim$  on  $\mathcal{L}^p(\mu)$  as follows: we write  $f \sim g$  if  $f = g$  a.e. on  $X$ .

### Proposition 1.3

The relation  $\sim$  on  $\mathcal{L}^p(\mu)$  is an equivalence relation.

### Proof .

It is obvious that  $f \sim f$  and that, if  $f \sim g$ , then  $g \sim f$ . Now, if  $f \sim g$  and  $g \sim h$ , then there exist  $A, B \in \mathcal{A}$  with  $\mu(A^c) = \mu(B^c) = 0$  such that  $f = g$  on  $A$  and  $g = h$  on  $B$ . This implies that  $\mu((A \cap B)^c) = 0$  and  $f = h$  on  $A \cap B$  and, hence,  $f \sim h$ .

The relation  $\sim$  defines equivalence classes. The equivalence class  $[f]$  of any  $f \in \mathcal{L}^p(\mu)$  is the set of all  $g \in \mathcal{L}^p(\mu)$  which are equivalents to  $f$ :  $[f] = \{g \in \mathcal{L}^p(\mu); g \sim f\}$ .

### Definition 1.4

We define  $L^p(\mu) = \mathcal{L}^p(\mu) / \sim = \{[f]; f \in \mathcal{L}^p(\mu)\}$ .

### Proposition 1.5

For  $p \geq 1$ , the space  $L^p(\mu)$  is an  $\mathbb{R}$  vector space.

**Proof .**

We shall use the trivial inequality  $|a + b|^p \leq 2^{p-1}(|a|^p + |b|^p)$ , for  $p \geq 1$  and  $a, b \in \mathbb{R}$ . For  $p = 1$  the statement is obvious. For  $p > 1$  the function  $y = x^p$ ;  $x > 0$  is convex since  $y'' \geq 0$ . Therefore  $\left(\frac{|a| + |b|}{2}\right)^p \leq \frac{|a|^p + |b|^p}{2}$ .

Assume that  $f, g$  are in  $L^p(\mu)$ . Then both functions  $f$  and  $g$  are finite a.e. on  $X$  and, hence,  $f + g$  is defined a.e. on  $X$ . If  $f + g$  is any measurable definition of  $f + g$ , then, using the above elementary inequality,  $|(f + g)(x)|^p \leq 2^{p-1}(|f(x)|^p + |g(x)|^p)$  for a.e.  $x \in X$  and, hence,

$$\int_X |f(x) + g(x)|^p d\mu(x) \leq 2^p \int_X |f(x)|^p d\mu(x) + 2^p \int_X |g(x)|^p d\mu(x) < +\infty.$$

Therefore  $f + g \in L^p(\mu)$ .

If  $f \in L^p(\mu)$  and  $\lambda \in \mathbb{R}$ , then  $\int_X |\lambda f(x)|^p d\mu(x) = |\lambda|^p \int_X |f(x)|^p d\mu(x) < +\infty$ .

Therefore,  $\lambda f \in L^p(\mu)$ .

### Definition 1.6

Let  $f: X \rightarrow \bar{\mathbb{R}}$  be measurable. We say that  $f$  is essentially bounded over  $X$  with respect to the measure  $\mu$  if there exists  $M < +\infty$  such that  $|f| \leq M$  a.e. on  $X$ .

### Proposition 1.7

Let  $f: X \rightarrow \bar{\mathbb{R}}$  be measurable. If  $f$  is essentially bounded over  $X$  with respect to the measure  $\mu$ , then there exists a smallest  $M$  with the property:  $|f| \leq M$  a.e. on  $X$ . This smallest  $M_0$  is characterized by:

- i)  $|f| \leq M_0$  a.e. on  $X$ ,
- ii)  $\mu(\{x \in X; |f(x)| > m\}) > 0$  for every  $m < M_0$ .

**Proof .**

We set  $A = \{M; |f| \leq M \text{ a.e. on } X\}$  and  $M_0 = \inf A$ . The set  $A$  is non-empty by assumption and is included in  $[0, +\infty[$  and, hence,  $M_0$  exists. We take a decreasing sequence  $(M_n)_n$  in  $A$  with  $\lim_{n \rightarrow +\infty} M_n = M_0$ . From  $M_n \in A$ , the set  $A_n = \{x \in X; |f(x)| > M_n\}$  is a null set for every  $n$  and, since  $\{x \in X; |f(x)| > M_0\} = \bigcup_{n=1}^{+\infty} A_n$ , we conclude that  $\{x \in X; |f(x)| > M_0\}$  is a null set. Therefore,  $|f| \leq M_0$  a.e. on  $X$ . If  $m < M_0$ , then  $m \notin A$  and, hence,  $\mu(\{x \in X; |f(x)| > m\}) > 0$ .  $\square$

### Definition 1.8

Let  $f: X \rightarrow \bar{\mathbb{R}}$  be a measurable function. If  $f$  is essentially bounded, then the smallest  $M$  with the property that  $|f| \leq M$  a.e. on  $X$  is called the essential supremum of  $f$  over  $X$  with respect to the measure  $\mu$  and it is denoted by  $\text{ess.sup}_X(f)$  or  $\|f\|_\infty$ .

The  $\|f\|_\infty$  is characterized by the properties:

1.  $|f| \leq \|f\|_\infty$  a.e. on  $X$ ,
2. for every  $m < \|f\|_\infty$ ,  $\mu(\{x \in X; |f(x)| > m\}) > 0$ .

**Definition 1.9**

We define  $L^\infty(\mu)$  to be the set of all measurable functions  $f: X \rightarrow \bar{\mathbb{R}}$  which are essentially bounded over  $X$  with respect to the measure  $\mu$ .

**Proposition 1.10**

The space  $L^\infty(\mu)$  is a linear space over  $\mathbb{R}$ .

**Proof .**

If  $f, g$  in  $L^\infty(\mu)$ , then there exist two subsets  $A_1, A_2 \in \mathcal{A}$  such that  $\mu(A_1^c) = \mu(A_2^c) = 0$  and  $|f| \leq \|f\|_\infty$  on  $A_1$  and  $|g| \leq \|g\|_\infty$  on  $A_2$ . If we set  $A = A_1 \cap A_2$ , then we have  $\mu(A^c) = 0$  and  $|f + g| \leq |f| + |g| \leq \|f\|_\infty + \|g\|_\infty$  on  $A$ . Hence  $f + g$  is essentially bounded over  $X$  with respect to the measure  $\mu$  and

$$\|f + g\|_\infty \leq \|f\|_\infty + \|g\|_\infty.$$

If  $f \in L^\infty(\mu)$  and  $\lambda \in \mathbb{R}$ , then there exists  $A \in \mathcal{A}$  with  $\mu(A^c) = 0$  such that  $|f| \leq \|f\|_\infty$  on  $A$ . Then  $|\lambda f| \leq |\lambda| \|f\|_\infty$  on  $A$ . Hence  $\lambda f$  is essentially bounded over  $X$  with respect to the measure  $\mu$  and  $\|\lambda f\|_\infty = |\lambda| \|f\|_\infty$ .

**Definition 1.11**

Let  $1 \leq p \leq +\infty$ . We define  $q = \frac{p}{p-1}$ , if  $1 < p < \infty$ ,  $q = \infty$  if  $p = 1$  and  $q = 1$ , if  $p = \infty$ . We say that  $q$  is the conjugate of  $p$  or the dual of  $p$ . Moreover,  $p, q$  are related by the symmetric equality

$$\frac{1}{p} + \frac{1}{q} = 1.$$

**Lemma 1.12**

Let  $p$  and  $q$  be two conjugate real numbers such that  $p > 1$ . Then for all  $a > 0; b > 0$ , we have

$$ab \leq \frac{a^p}{p} + \frac{b^q}{q}$$

**Proof .**

Note that  $\varphi(t) = \frac{t^p}{p} + \frac{1}{q} - t$  with  $t \geq 0$  has the only minimum at  $t = 1$ . It follows that  $t \leq \frac{t^p}{p} + \frac{1}{q}$ .

For  $t = ab^{-\frac{1}{p-1}}$  we have  $\frac{a^p b^{-q}}{p} + \frac{1}{q} \geq ab^{-\frac{1}{p-1}}$ ; and the result follows.  $\square$

**Theorem 1.13 (Hölder's inequalities)**

Let  $1 \leq p, q \leq +\infty$  and  $p, q$  be conjugate to each other. If  $f \in L^p(\mu)$  and  $g \in L^q(\mu)$ , then  $fg \in L^1(\mu)$  and

$$\int_X |fg(x)| d\mu(x) \leq \left( \int_X |f(x)|^p d\mu(x) \right)^{\frac{1}{p}} \left( \int_X |g(x)|^q d\mu(x) \right)^{\frac{1}{q}}$$

$$\int_X |fg(x)| d\mu(x) \leq \|g\|_\infty \int_X |f(x)| d\mu(x), \quad p = 1, q = +\infty.$$

**Proof .**

We start with the case  $1 < p, q < +\infty$ . If  $\int_X |f(x)|^p d\mu(x) = 0$  or if  $\int_X |g(x)|^q d\mu(x) = 0$ , then either  $f = 0$  a.e. on  $X$  or  $g = 0$  a.e. on  $X$  and the inequality is trivially true. So we assume that  $A = \int_X |f(x)|^p d\mu(x) > 0$  and  $B = \int_X |g(x)|^q d\mu(x) > 0$ . Applying lemma 1.12 with  $a = \frac{|f|}{A^{\frac{1}{p}}}$ ,  $b = \frac{|g|}{B^{\frac{1}{q}}}$ , we have that

$$\frac{|fg|}{A^{\frac{1}{p}} B^{\frac{1}{q}}} \leq \frac{1}{p} \frac{|f|^p}{A} + \frac{1}{q} \frac{|g|^q}{B}$$

a.e. on  $X$ . Integrating, we find

$$\frac{1}{A^{\frac{1}{p}} B^{\frac{1}{q}}} \int_X |fg(x)| d\mu(x) \leq \frac{1}{p} + \frac{1}{q} = 1$$

Then

$$\int_X |fg(x)| d\mu(x) \leq \left( \int_X |f(x)|^p d\mu(x) \right)^{\frac{1}{p}} \left( \int_X |g(x)|^q d\mu(x) \right)^{\frac{1}{q}}$$

Let now  $p = 1$  and  $q = +\infty$ . Since  $|g| \leq \|g\|_\infty$  a.e. on  $X$ , we have that  $|fg| \leq |f| \|g\|_\infty$  a.e. on  $X$ . Integrating, we find the inequality we want to prove.

**Theorem 1.14** (*Minkowski's inequality*)

Let  $1 \leq p < +\infty$ . If  $f, g$  in  $L^p(\mu)$ , then

$$\left( \int_X |f(x)+g(x)|^p d\mu(x) \right)^{\frac{1}{p}} \leq \left( \int_X |f(x)|^p d\mu(x) \right)^{\frac{1}{p}} + \left( \int_X |g(x)|^p d\mu(x) \right)^{\frac{1}{p}}, \quad 1 \leq p < +\infty.$$

**Proof .**

The case  $p = 1$  is trivial. Hence, we assume that  $1 < p < +\infty$ .

We write  $|f+g|^p \leq (|f|+|g|)|f+g|^{p-1} = |f||f+g|^{p-1} + |g||f+g|^{p-1}$  a.e. on  $X$  and, applying Hölder's inequality, we find

$$\begin{aligned} \int_X |f(x)+g(x)|^p d\mu(x) &\leq \left( \int_X |f(x)|^p d\mu(x) \right)^{\frac{1}{p}} \left( \int_X |f(x)+g(x)|^{(p-1)q} d\mu(x) \right)^{\frac{1}{q}} \\ &\quad + \left( \int_X |g(x)|^p d\mu(x) \right)^{\frac{1}{p}} \left( \int_X |f(x)+g(x)|^{(p-1)q} d\mu(x) \right)^{\frac{1}{q}} \\ &= \left( \int_X |f(x)|^p d\mu(x) \right)^{\frac{1}{p}} \left( \int_X |f(x)+g(x)|^p d\mu(x) \right)^{\frac{1}{q}} \end{aligned}$$

Simplifying, we get the inequality we want to prove.

**Corollary 1.15**

The mapping  $f \mapsto \|f\|_p = \left( \int_X |f(x)|^p \right)^{\frac{1}{p}}$  is a norm on  $L^p(\mu)$  and  $(L^p(\mu), \|\cdot\|_p)$  is a vector space.

## 2 Convergence of Functions

### Definition 2.1

Let  $A$  be an arbitrary non empty set and  $(f_n: A \rightarrow \bar{\mathbb{R}})_n$  be a sequence of functions defined on  $A$ .

1. We say that the sequence  $(f_n)_n$  converges pointwise on  $A$  to a function  $f: A \rightarrow \bar{\mathbb{R}}$  if  $\lim_{n \rightarrow +\infty} f_n(x) = f(x)$  for every  $x \in A$ .

In case  $f(x)$  is finite, this means that:  $\forall \varepsilon > 0, \exists N \in \mathbb{N}$  such that:  $|f_n(x) - f(x)| \leq \varepsilon; \forall n \geq N$ .

2. Let  $(X, \mathcal{A}, \mu)$  be a measure space. We say that the sequence  $(f_n)_n$  converges to  $f$  (pointwise) a.e. on  $A \in \mathcal{A}$  if there exists a set  $B \in \mathcal{A}, B \subset A$ , such that  $\mu(A \setminus B) = 0$  and  $(f_n)_n$  converges to  $f$  pointwise on  $B$ .

### Remark .

If  $(f_n)_n$  converges to both  $f$  and  $g$  a.e. on  $A$ , then  $f = g$  a.e. on  $A$ .

## 2.1 Convergence in $L^p$

### Definition 2.2

Let  $(f_n)_n$  be a sequence in  $L^p(\mu)$  and  $f \in L^p(\mu)$ . We say that  $(f_n)_n$  converges to  $f$  in  $L^p(\mu)$  if:  $\lim_{n \rightarrow +\infty} \|f_n - f\|_p = 0$ .

We say that  $(f_n)_n$  is Cauchy in  $L^p(\mu)$  if:  $\lim_{n, m \rightarrow +\infty} \|f_n - f_m\|_p = 0$ .

### Theorem 2.3

If  $(f_n)_n$  is Cauchy sequence in  $L^p(\mu)$ , then there exists  $f \in L^p(\mu)$  such that  $(f_n)_n$  converges to  $f$  in  $L^p(\mu)$ . (In other words  $L^p(\mu)$  is a Banach space.)

Moreover, there exists a subsequence  $(f_{n_k})_k$  which converges to  $f$  a.e. on  $X$ .

### Corollary 2.4

If  $(f_n)_n$  converges to  $f$  in  $L^p(\mu)$ , there is a subsequence  $(f_{n_k})_k$  which converges to  $f$  a.e. on  $X$ .

### Proof .

a) We consider the first case  $1 \leq p < +\infty$ .

Since each  $f_n$  is finite a.e. on  $X$ , there exists  $A \in \mathcal{A}$  such that  $\mu(A^c) = 0$  and all  $f_n$  are finite on  $A$ . Then for every  $k$ , there exists  $n_k$  such that  $\int_X |f_n(x) - f_m(x)|^p d\mu(x) < \frac{1}{2^{kp}}$  for every  $n, m \geq n_k$ . Since we may assume that each  $n_k$  is large enough, then we can take  $n_k < n_{k+1}$  for every  $k$ . Therefore,  $(f_{n_k})_k$  is a subsequence of  $(f_n)_n$ .

From the construction of  $n_k$  and from the fact that  $n_k < n_{k+1}$ , we get  $\int_X |f_{n_{k+1}}(x) - f_{n_k}(x)|^p d\mu(x) < \frac{1}{2^{kp}}$  for every  $k$ . We define the measurable function  $G$  by:  $G =$

$$\sum_{k=1}^{+\infty} |f_{n_{k+1}} - f_{n_k}|, \text{ on } A \text{ and } G = 0, \text{ on } A^c.$$

Let  $G_N = \sum_{k=1}^N |f_{n_{k+1}} - f_{n_k}|$  on  $A$  and  $G_N = 0$  on  $A^c$ , then  $\left(\int_X G_N^p(x) d\mu(x)\right)^{\frac{1}{p}} \leq \sum_{k=1}^N \left(\int_X |f_{n_{k+1}}(x) - f_{n_k}(x)|^p d\mu(x)\right)^{\frac{1}{p}} < 1$ , by Minkowski's inequality. Since  $(G_N)_N$  increases to  $G$  on  $X$ , we find that  $\int_X G^p(x) d\mu(x) \leq 1$  and, thus,  $G < +\infty$  a.e. on  $X$ .

It follows that the series  $\sum_{k=1}^{+\infty} (f_{n_{k+1}}(x) - f_{n_k}(x))$  converges for a.e.  $x \in A$ . Therefore,

there exists a  $B \in \mathcal{A}$ ,  $B \subset A$  such that  $\mu(A \setminus B) = 0$  and  $\sum_{k=1}^{+\infty} (f_{n_{k+1}}(x) - f_{n_k}(x))$  converges for every  $x \in B$ . We define the measurable function  $f$  by:  $f = f_{n_1} + \sum_{k=1}^{+\infty} (f_{n_{k+1}} - f_{n_k})$  on  $B$  and 0 on  $B^c$ .

On  $B$  we have  $f = f_{n_1} + \lim_{N \rightarrow +\infty} \sum_{k=1}^N (f_{n_{k+1}} - f_{n_k}) = \lim_{N \rightarrow +\infty} f_{n_N}$  and, hence,  $(f_{n_k})_k$  converges to  $f$  a.e. on  $X$ . We, also, have on  $B$

$$\begin{aligned} |f_{n_N} - f| &= \left| f_{n_N} - f_{n_1} - \sum_{k=1}^{+\infty} (f_{n_{k+1}} - f_{n_k}) \right| \\ &= \left| \sum_{k=1}^{N-1} (f_{n_{k+1}} - f_{n_k}) - \sum_{k=1}^{+\infty} (f_{n_{k+1}} - f_{n_k}) \right| \\ &\leq \sum_{k=N+1}^{+\infty} |f_{n_{k+1}} - f_{n_k}| \leq G \end{aligned}$$

for every  $N$  and, hence,  $|f_{n_N} - f|^p \leq G^p$  a.e. on  $X$  for every  $N$ . Since  $\int_X G^p(x) d\mu(x) < +\infty$  and  $\lim_{N \rightarrow +\infty} |f_{n_N} - f| = 0$  a.e. on  $X$ , we apply the Dominated Convergence Theorem we find that

$$\lim_{N \rightarrow +\infty} \int_X |f_{n_N}(x) - f(x)|^p d\mu(x) = 0$$

If  $n_k \rightarrow +\infty$ , we get

$$\begin{aligned} \lim_{k \rightarrow +\infty} \left( \int_X |f_k(x) - f(x)|^p d\mu(x) \right)^{\frac{1}{p}} &\leq \lim_{k \rightarrow +\infty} \left[ \left( \int_X |f_k(x) - f_{n_k}(x)|^p d\mu(x) \right)^{\frac{1}{p}} \right. \\ &\quad \left. + \left( \int_X |f_{n_k}(x) - f(x)|^p d\mu(x) \right)^{\frac{1}{p}} \right] = 0 \end{aligned}$$

and we conclude that  $(f_n)_n$  converges to  $f$  in  $L^p(\mu)$ .

b) Now, let  $p = +\infty$ . For each  $n, m$  we have a set  $A_{n,m} \in \mathcal{A}$  such that  $\mu(A_{n,m}^c) = 0$  and  $|f_n - f_m| \leq \|f_n - f_m\|_\infty$  on  $A_{n,m}$ .

Let  $A = \bigcap_{n,m \geq 1} A_{n,m}$ , then  $\mu(A^c) = 0$  and  $|f_n - f_m| \leq \|f_n - f_m\|_\infty$  on  $A$  for every  $n, m$ .

This gives that  $(f_n)_n$  is Cauchy sequence for the  $\|\cdot\|_\infty$  on  $A$  and, hence, there exists a mapping  $f$  such that  $(f_n)_n$  converges to  $f$  uniformly on  $A$ . Now,

$$\lim_{n \rightarrow +\infty} \text{ess.sup}_X(f_n - f) \leq \lim_{n \rightarrow +\infty} \text{Sup}_{x \in A} |f_n(x) - f(x)| = 0$$

## 2.2 Convergence in Measure

Let  $(X, \mathcal{A}, \mu)$  is a measure space.

### Definition 2.5

1. Let  $f, f_n: X \rightarrow \bar{\mathbb{R}}$  be measurable functions. We say that  $(f_n)_n$  converges to  $f$  in measure on  $A \in \mathcal{A}$  if all  $f, f_n$  are finite a.e. on  $A$  and for every  $\varepsilon > 0$ ;

$$\lim_{n \rightarrow +\infty} \mu(\{x \in A; |f_n(x) - f(x)| \geq \varepsilon\}) = 0.$$

2. We say that  $(f_n)_n$  is a Cauchy sequence in measure on  $A \in \mathcal{A}$  if all  $f_n$  are finite a.e. on  $A$  and for every  $\varepsilon > 0$

$$\lim_{n,m \rightarrow +\infty} \mu(\{x \in A; |f_n(x) - f_m(x)| \geq \varepsilon\}) = 0.$$

### Remarks .

1. The uniform convergence yields the convergence in measure
2. If we want to be able to write the values  $\mu(\{x \in A; |f_n(x) - f(x)| \geq \varepsilon\})$  and  $\mu(\{x \in A; |f_n(x) - f_m(x)| \geq \varepsilon\})$ , we first extend the functions  $|f_n - f|$  and  $|f_n - f_m|$  outside the set  $B \subset A$ , where all  $f, f_n$  are finites, as functions defined on  $X$  and measurables. Then, since  $\mu(A \setminus B) = 0$ , we get that the above values are equal to the values  $\mu(\{x \in B; |f_n(x) - f(x)| \geq \varepsilon\})$  and, respectively,  $\mu(\{x \in B; |f_n(x) - f_m(x)| \geq \varepsilon\})$ . Therefore, the actual extensions play no role and, hence, we may for simplicity extend all  $f, f_n$  as 0 on  $X \setminus B$ . Thus the replacement of all  $f, f_n$  by 0 on  $X \setminus B$  makes all functions finite everywhere on  $A$  and does not affect the fact that  $(f_n)_n$  converges to  $f$  in measure on  $A$  or that  $(f_n)_n$  is Cauchy in measure on  $A$ .
3. Let  $a, b > 0$  and  $A = \{x \in A; |f(x)| \geq a\}$ ,  $B = \{x \in A; |g(x)| \geq b\}$  and  $C = \{x \in A; |f(x) + g(x)| \geq a + b\}$ . If  $x \in C \setminus A$ , then  $|f(x)| < a$  and  $a + b \leq |f(x) + g(x)| \leq |f(x)| + |g(x)| < a + |g(x)|$ . It follows that  $\{x \in A; |f(x) + g(x)| \geq a + b\} \subset \{x \in A; |f(x)| \geq a\} \cup \{x \in A; |g(x)| \geq b\}$  and

$$\begin{aligned} \mu(\{x \in A; |f(x) + g(x)| \geq a + b\}) &\leq \mu(\{x \in A; |f(x)| \geq a\}) \\ &\quad + \mu(\{x \in A; |g(x)| \geq b\}). \end{aligned}$$

**Proposition 2.6**

If  $(f_n)_n$  converges to both  $f$  and  $g$  in measure on  $A$ , then  $f = g$  a.e. on  $A$ .

**Proof .**

We may assume that all  $f, g, f_n$  are finites on  $A$ . Applying the above remark we find that:

$$\begin{aligned} \mu(\{x \in A; |f(x) - g(x)| \geq \varepsilon\}) &\leq \mu(\{x \in A; |f_n(x) - f(x)| \geq \frac{\varepsilon}{2}\}) \\ &\quad + \mu(\{x \in A; |f_n(x) - g(x)| \geq \frac{\varepsilon}{2}\}) \xrightarrow{n \rightarrow +\infty} 0. \end{aligned}$$

This implies that  $\mu(\{x \in A; |f(x) - g(x)| \geq \varepsilon\}) = 0$  for every  $\varepsilon > 0$ . We now write

$$\{x \in A; f(x) \neq g(x)\} = \bigcup_{k=1}^{+\infty} \{x \in A; |f(x) - g(x)| \geq \frac{1}{k}\}.$$

Since each term in the union is a null set, we get  $\mu(\{x \in A; f(x) \neq g(x)\}) = 0$  and we conclude that  $f = g$  a.e. on  $A$ .

**Proposition 2.7**

If  $(f_n)_n$  converges to  $f$  and  $(g_n)_n$  converges to  $g$  in measure on  $A$  and let  $\alpha \in \mathbb{R}$ . Then

- a)  $(f_n + g_n)_n$  converges to  $f + g$  in measure on  $A$ .
- b)  $(\alpha f_n)_n$  converges to  $\alpha f$  in measure on  $A$ .
- c) If there exists  $M < +\infty$  such that  $|f_n| \leq M$  a.e. on  $A$ , then  $|f| \leq M$  a.e. on  $A$ .
- d) If there exists  $M < +\infty$  such that  $|f_n|, |g_n| \leq M$  a.e. on  $A$ , then  $(f_n g_n)_n$  converges to  $fg$  in measure on  $A$ .

**Proof .**

We may assume that all  $f, f_n$  are finites on  $A$ .

- a) We apply the remark 3

$$\begin{aligned} \mu(\{x \in A; |(f_n + g_n)(x) - (f + g)(x)| \geq \varepsilon\}) &\leq \mu(\{x \in A; |f_n(x) - f(x)| \geq \frac{\varepsilon}{2}\}) \\ &\quad + \mu(\{x \in A; |g_n(x) - g(x)| \geq \frac{\varepsilon}{2}\}) \xrightarrow{n \rightarrow +\infty} 0. \end{aligned}$$

- b) Also for  $\alpha \neq 0$ ,

$$\mu(\{x \in A; |\alpha f_n(x) - \alpha f(x)| \geq \varepsilon\}) = \mu(\{x \in A; |f_n(x) - f(x)| \geq \frac{\varepsilon}{|\alpha|}\}) \xrightarrow{n \rightarrow +\infty} 0.$$

c) For  $n$  large enough

$$\begin{aligned} \mu(\{x \in A; |f(x)| \geq M + \varepsilon\}) &\leq \mu(\{x \in A; |f_n(x)| \geq M + \frac{\varepsilon}{2}\}) \\ &\quad + \mu(\{x \in A; |f_n(x) - f(x)| \geq \frac{\varepsilon}{2}\}) \\ &= \mu(\{x \in A; |f_n(x) - f(x)| \geq \frac{\varepsilon}{2}\}) \xrightarrow{n \rightarrow +\infty} 0. \end{aligned}$$

Hence,  $\mu(\{x \in A; |f(x)| \geq M + \varepsilon\}) = 0$  for every  $\varepsilon > 0$ .

We have  $\{x \in A; |f(x)| > M\} = \bigcup_{k=1}^{+\infty} \{x \in A; |f(x)| \geq M + \frac{1}{k}\}$  and, since all sets of the union are null sets, then  $\mu(\{x \in A; |f(x)| > M\}) = 0$ . Hence,  $|f| \leq M$  a.e. on  $A$ .

d) Applying the result of c),

$$\begin{aligned} \mu(\{x \in A; |f_n(x)g_n(x) - f(x)g(x)| \geq \varepsilon\}) &\leq \mu(\{x \in A; |f_n(x)g_n(x) - f_n(x)g(x)| \geq \frac{\varepsilon}{2}\}) \\ &\quad + \mu(\{x \in A; |f_n(x)g(x) - f(x)g(x)| \geq \frac{\varepsilon}{2}\}) \\ &\leq \mu(\{x \in A; |g_n(x) - g(x)| \geq \frac{\varepsilon}{2M}\}) \\ &\quad + \mu(\{x \in A; |f_n(x) - f(x)| \geq \frac{\varepsilon}{2M}\}) \xrightarrow{n \rightarrow +\infty} 0. \end{aligned}$$

□

### Proposition 2.8

Let  $(f_n)_n$  be a sequence of measurable functions on a measure space  $(X, \mathcal{B}, \mu)$ . If  $\mu$  is finite and the sequence  $(f_n)_n$  converges almost everywhere to  $f$ , then the sequence  $(f_n)_n$  converges in measure to  $f$ .

### Proof .

Let  $\varepsilon > 0$ , we set

$$A_n(\varepsilon) = \{x; |f_n(x) - f(x)| \geq \varepsilon\}, \quad B_n(\varepsilon) = \bigcup_{k \geq n} A_k(\varepsilon)$$

and

$$B(\varepsilon) = \bigcap_{n \geq 1} B_n(\varepsilon) = \overline{\lim}_{n \rightarrow +\infty} A_n(\varepsilon)$$

If  $x \in B(\varepsilon)$ , then  $x$  belongs to an infinite of  $A_n(\varepsilon)$ . Then  $(f_n(x))_n$  can not converges to  $f(x)$  and then  $\mu(B(\varepsilon)) = 0$ . Moreover since  $\mu$  is finite  $\lim_{n \rightarrow +\infty} \mu(B_n(\varepsilon)) = 0$ , and since  $A_n(\varepsilon) \subset B_n(\varepsilon)$ , then  $\lim_{n \rightarrow +\infty} \mu(A_n(\varepsilon)) = 0$ .

### Theorem 2.9

If  $(f_n)_n$  is Cauchy in measure on  $A$ , then there exists  $f: X \rightarrow \bar{\mathbb{R}}$  such that  $(f_n)_n$  converges to  $f$  in measure on  $A$ . Moreover, there is a subsequence  $(f_{n_k})_k$  which converges to  $f$  a.e. on  $A$ .

**Corollary 2.10**

If  $(f_n)_n$  converges to  $f$  in measure on  $A$ , there is a subsequence  $(f_{n_k})_k$  which converges to  $f$  a.e. on  $A$ .

**Proof .**

As usual, we assume that all  $f_n$  are finites on  $A$ . We have, for all  $k$ ,  $\mu(\{x \in A; |f_n(x) - f_m(x)| \geq \frac{1}{2^k}\}) \xrightarrow{n,m \rightarrow +\infty} 0$ . Therefore, there exists  $n_k$  such that  $\mu(\{x \in A; |f_n(x) - f_m(x)| \geq \frac{1}{2^k}\}) < \frac{1}{2^k}$  for every  $n, m \geq n_k$ . Since we may assume that each  $n_k$  is as large as we like, we may inductively take  $n_k$  such that  $n_k < n_{k+1}$  for every  $k$ . Hence,  $(f_{n_k})_k$  is a subsequence of  $(f_n)_n$  and, from the construction of  $n_k$  and since  $n_k < n_{k+1}$ , we have that for every  $k$ ;

$$\mu(\{x \in A; |f_{n_{k+1}}(x) - f_{n_k}(x)| \geq \frac{1}{2^k}\}) < \frac{1}{2^k}$$

Let  $E_k = \{x \in A; |f_{n_{k+1}}(x) - f_{n_k}(x)| \geq \frac{1}{2^k}\}$  and, hence,  $\mu(E_k) < \frac{1}{2^k}$  for all  $k$ . We also

define the subsets of  $A$ :  $F_m = \bigcup_{k=m}^{+\infty} E_k$ ,  $F = \bigcap_{m=1}^{+\infty} F_m = \overline{\lim}_{k \rightarrow +\infty} E_k$ .

$\mu(F_m) \leq \sum_{k=m}^{+\infty} \mu(E_k) < \sum_{k=m}^{+\infty} \frac{1}{2^k} = \frac{1}{2^{m-1}}$  and, hence,  $\mu(F) \leq \mu(F_m) < \frac{1}{2^{m-1}}$  for every  $m$ .

This implies that  $\mu(F) = 0$ .

If  $x \in A \setminus F$ , then there exists  $m$  such that  $x \in A \setminus F_m$ , which implies that  $x \in A \setminus E_k$  for all  $k \geq m$ . Therefore,  $|f_{n_{k+1}}(x) - f_{n_k}(x)| < \frac{1}{2^k}$  for all  $k \geq m$ , such that

$\sum_{k=m}^{+\infty} |f_{n_{k+1}}(x) - f_{n_k}(x)| < \frac{1}{2^{m-1}}$ . Thus, the series  $\sum_{k=m}^{+\infty} (f_{n_{k+1}}(x) - f_{n_k}(x))$  converges and

we may define  $f: X \rightarrow \bar{\mathbb{R}}$  by:

$$f = f_{n_1}(x) + \sum_{k=1}^{+\infty} (f_{n_{k+1}} - f_{n_k}),$$

on  $A \setminus F$  and 0, on  $A^c \cup F$ .

$$f(x) = f_{n_1}(x) + \lim_{m \rightarrow +\infty} \sum_{k=1}^m (f_{n_{k+1}}(x) - f_{n_k}(x)) = \lim_{m \rightarrow +\infty} f_{n_m}(x)$$

for every  $x \in A \setminus F$  and since  $\mu(F) = 0$ , we get that  $(f_{n_k})_k$  converges to  $f$  a.e.

Now, on  $A \setminus F_m$ ; we have

$$\begin{aligned} |f_{n_m} - f| &= |f_{n_m} - f_{n_1} - \sum_{k=1}^{+\infty} (f_{n_{k+1}} - f_{n_k})| \\ &= \left| \sum_{k=1}^{m-1} (f_{n_{k+1}} - f_{n_k}) - \sum_{k=1}^{+\infty} (f_{n_{k+1}} - f_{n_k}) \right| \end{aligned}$$

$$= \sum_{k=m}^{+\infty} |f_{n_{k+1}} - f_{n_k}| < \frac{1}{2^{m-1}}.$$

Therefore,  $\{x \in A; |f_{n_m}(x) - f(x)| \geq \frac{1}{2^{m-1}}\} \subset F_m$  and, hence,

$$\mu(\{x \in A; |f_{n_m}(x) - f(x)| \geq \frac{1}{2^{m-1}}\}) \leq \mu(F_m) < \frac{1}{2^{m-1}}.$$

Take an arbitrary  $\varepsilon > 0$  and  $m_0$  large enough such that  $\frac{1}{2^{m-1}} \leq \varepsilon$ . If  $m \geq m_0$ ,  $\{x \in A; |f_{n_m}(x) - f(x)| \geq \varepsilon\} \subset \{x \in A; |f_{n_m}(x) - f(x)| \geq \frac{1}{2^{m-1}}\}$  and, hence,

$$\mu(\{x \in A; |f_{n_m}(x) - f(x)| \geq \varepsilon\}) < \frac{1}{2^{m-1}} \xrightarrow{m \rightarrow +\infty} 0.$$

This means that  $(f_{n_k})_k$  converges to  $f$  in measure on  $A$ . Since  $n_k \xrightarrow{k \rightarrow +\infty} +\infty$ , we have

$$\begin{aligned} \mu(\{x \in A; |f_k(x) - f(x)| \geq \varepsilon\}) &= \mu(\{x \in A; |f_k(x) - f_{n_k}(x)| \geq \frac{\varepsilon}{2}\}) \\ &\quad + \mu(\{x \in A; |f_{n_k}(x) - f(x)| \geq \frac{\varepsilon}{2}\}) \xrightarrow{k \rightarrow +\infty} 0. \end{aligned}$$

and we conclude that  $(f_n)_n$  converges to  $f$  in measure on  $A$ .

**Example .**

Consider the sequence  $f_1 = \chi_{]0,1[}$ ,  $f_2 = 2\chi_{]0,\frac{1}{2}[}$ ,  $f_3 = 2\chi_{]0,\frac{1}{2}[}$ , and for all  $n \in \mathbb{N}$ ,  $f_{\frac{n(n+1)}{2}+k+1} = n\chi_{]0,\frac{k}{n+1},\frac{k+1}{n+1}[}$ , for  $k = 0, \dots, n$ . If  $0 < \varepsilon \leq 1$ , the sequence of the values  $\mu(\{x \in ]0,1[; |f_n(x)| \geq \varepsilon\}) \xrightarrow{n \rightarrow +\infty} 0$ . Therefore,  $(f_n)_n$  converges to 0 in measure on  $]0,1[$ . But, as we have already seen, it is not true that  $(f_n)_n$  converges to 0 a.e. on  $]0,1[$ .