

**King Saud University**  
**College of Sciences**  
**Department of Mathematics**  
**MATH-244 (Linear Algebra); Final Exam; Semester 443**

**Max. Marks: 40****Time: 3 hours**

Name:	ID:	Section:	Signature:
-------	-----	----------	------------

**Note:** Attempt all the five questions. Scientific calculators are not allowed!

**Question 1** [Marks:  $10 \times 1$ ]:

Choose the correct answer:

(i) If the matrix  $\begin{bmatrix} 1 & 2 & -1 \\ 2 & 5 & -4 \\ 1 & 3 & h \end{bmatrix}$  is non-invertible, then  $h$  is equal to:  
 a) 5      b) -3      c) -5      d) 3.

(ii) Which of the following matrices cannot be a transition matrix?  
 a)  $\begin{bmatrix} 2 & -1 \\ 4 & 2 \end{bmatrix}$       b)  $\begin{bmatrix} 1 & 1 & 2 \\ 2 & 1 & 3 \\ 0 & 1 & 0 \end{bmatrix}$       c)  $\begin{bmatrix} 2 & 1 & 0 \\ 2 & 0 & 1 \\ 3 & 1 & 0 \end{bmatrix}$       d)  $\begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix}$ .

(iii) Let  $F$  denote the set of nontrivial solutions of homogenous linear system  $AX = 0$ , where the matrix of coefficients  $A$  is invertible and  $X \in \mathbb{R}^3$ . Then,  $F$  is equal to:  
 a)  $\mathbb{R}^3$       b)  $\{(0,0,0)\}$       c)  $\{\}$       d)  $\mathbb{R}^3 - \{(0,0,0)\}$ .

(iv) If  $W = \text{span} \{(1, -1, 0, 1), (-1, 1, 1, 0), (2, -2, 1, 3)\}$ , then  $\dim W$  is equal to:  
 a) 1      b) 2      c) 3      d) 4.

(v) If  $B = \{(2, -4), (3, -3)\}$  is an ordered basis of the vector space  $\mathbb{R}^2$ , then the coordinate vector  $[(1,1)]_B$  is equal to:  
 a)  $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$       b)  $\begin{bmatrix} -1 \\ 1 \end{bmatrix}$       c)  $\begin{bmatrix} \frac{1}{2} \\ 1 \end{bmatrix}$       d)  $\begin{bmatrix} -1 \\ 2 \end{bmatrix}$ .

(vi) If  $U = \begin{bmatrix} 2 & 5 \\ -1 & x \end{bmatrix}$  and  $V = \begin{bmatrix} -3 & 2x \\ 3 & -1 \end{bmatrix}$  are orthogonal matrices in  $M_2(\mathbb{R})$  with respect to the inner product  $\langle A, B \rangle = \text{trace}(AB^T)$ , then  $x$  is equal to:  
 a) 2      b) -2      c) 0      d) 1.

(vii) If  $\langle \cdot, \cdot \rangle$  is an inner product on  $\mathbb{R}^3$  such that  $\|u\|^2 = 2$ ,  $\|v\|^2 = 3$ ,  $\langle u, v \rangle = 1$ , then  $\langle 3u - v, 2u - 4v \rangle$  is equal to:  
 a)  $\sqrt{13}$       b) -14      c) 10      d) 38.

(viii) If the inner product on the vector space  $P_2$  of polynomials with degree  $\leq 2$  is defined by  $\langle p, q \rangle = aa_1 + 2bb_1 + cc_1$  for all  $p = a + bx + cx^2$ ,  $q = a_1 + b_1x + c_1x^2 \in P_2$  and  $\theta$  denote the angle between polynomials  $2 + x + x^2$  and  $-1 + x + 2x^2$ , then  $\cos \theta$  is equal to  
 a)  $\frac{2}{\sqrt{7}}$       b)  $\frac{2}{7}$       c) 0      d)  $\frac{6}{7}$ .

(ix) If the transformation  $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$  is defined by  $T(x, y) = (2x - y, -8x + 4y)$ . Then  $\ker(T)$  includes:  
 a) (5,10)      b) (10,2)      c) (-5,10)      d) (10,5).

(x) If the characteristic polynomial of a matrix  $A$  is  $q_A(\lambda) = \lambda^2 - 9$ , then the matrix  $A$  is:  
 a) not diagonalizable      b) diagonalizable      c)  $3 \times 3$       d) not invertible.

**Question 2** [Marks: 2+2+2]: Let the matrix  $A = \begin{bmatrix} 1 & -1 & 3 \\ 5 & -4 & -4 \\ 7 & -6 & 2 \end{bmatrix}$ . Then:

- a) Find a basis of the null space of A.
- b) Find a basis for the column space of A.
- c) Verify that  $\text{rank}(A) + \text{nullity}(A) = 3$ .

**Question 3** [Marks: 3+3]:

- a) Let the set  $B = \{u_1, u_2, u_3\}$  be a basis for a vector space  $V$ . If  $u, v \in V$  are linearly independent vectors. Then show that  $\{[u]_B, [v]_B\}$  is a linearly independent subset of  $\mathbb{R}^3$ .
- b) Find a basis for the Euclidean space  $\mathbb{R}^4$  that includes the vectors  $(0,0,0,1), (1,1,1,0), (0,1,1,0)$ .

**Question 4:** [Marks: 2+4]

- a) Let  $\{u, v, w\}$  be an orthonormal set of vectors in an inner product space. Use Pythagorean Theorem to evaluate  $\|u + v + w\|^2$ .
- b) Let the set  $B = \{u_1 = (1,1,1), u_2 = (0,1,1), u_3 = (0,0,1)\}$  be linearly independent in the Euclidean inner product space  $\mathbb{R}^3$ . Construct an orthonormal set  $C$  for  $\mathbb{R}^3$  by applying the Gram-Schmidt algorithm on  $B$  such that  $\text{span}(C) = \text{span}(B)$ .

**Question 5:** [Marks: (4+2) + (2+2+2)]

- a) Let  $A = \{v_1, v_2, v_3\}$  be a basis for vector space  $V$ ,  $B = \{w_1, w_2, w_3, w_4, w_5\}$  be a basis for vector space  $W$ . Let  $T : V \rightarrow W$  be the linear transformation such that:  $w_2 - 3w_4 + T(v_1) = 2w_1 + 7w_5; T(v_2) + w_4 = 2w_3 + w_5; w_3 + T(v_3) = 2w_2 + 4w_4 - w_5$ . Then:
  - (i) Find the transformation matrix  $[T]_A^B$  with respect to the ordered bases  $A$  and  $B$ .
  - (ii) Find the coordinate vector  $[T(v_1 + v_2 + v_3)]_B$ .
- b) Let  $A = \begin{bmatrix} 3 & 0 & 0 \\ -3 & 4 & 9 \\ 0 & 0 & 3 \end{bmatrix}$ . Then:
  - (i) Show that the matrix  $A$  is diagonalizable.
  - (ii) Find an invertible matrix  $P$  and a diagonal matrix  $D$  satisfying  $P^{-1}AP = D$ .
  - (iii) Find  $A^5$ .

\*\*\*!