

[Solution key]

King Saud University
College of Sciences
Department of Mathematics
Semester 462 / Final Exam / MATH-244 (Linear Algebra)

Max. Marks: 40**Time: 3 hours****Solution of Question 1: Correct choices:**

- (i) If square of a matrix A is zero matrix, then $I - A$ is equal to:
 a) 0 b) $(A-I)^{-1}$ c) ☒ $(A+I)^{-1}$ d) $A + I$ [Mark 1]
- (ii) If A is a square matrix of order 3 with $\det(A) = 2$, then $\det(\frac{1}{\det(A)} A^3) A^{-1}$ is equal to:
 a) $1/4$ b) ☒ $1/2$ c) $1/3$ d) $1/16$ [Mark 1]
- (iii) If the general solution of $AX = 0$ is $(-2r, 4r, r)$, $r \in \mathbb{R}$, and $(1, 0, -2)$ is a solution of $AX = B$, then the general solution of $AX = B$ is:
 a) ☒ $(1 - 2r, 4r, r - 2)$ b) $(-2r, 0, -2r)$ c) $(-2r, 4r, r)$ d) $(-2r - 1, 4r, r - 2)$ [Mark 1]
- (iv) A subset S of $\mathbb{R}^3$ is a basis of the vector space $\mathbb{R}^3$ if S is equal to:
 a) ☒ $\{(1, 0, 0), (0, 2, 1), (0, 6, 0)\}$ b) $\{(1, 1, 0), (2, 1, 0), (3, 2, 0)\}$ c) $\{(1, 1, 0), (0, 0, 0), (3, 2, 1)\}$ d) $\{(1, 1, 0), (0, 0, 1), (2, 2, 1)\}$ [Mark 1]
- (v) If $B = \{u_1 = (2, 1), u_2 = (4, 3)\}$ and $C = \{v_1 = (0, 1), v_2 = (6, 0)\}$ are ordered bases of $\mathbb{R}^2$, then the transition matrix $P_{C \rightarrow B}$ from C to B is equal to:
 a) $\begin{bmatrix} 1 & -1/2 \\ -2 & 3/2 \end{bmatrix}$ b) ☒ $\begin{bmatrix} -2 & 9 \\ 1 & -3 \end{bmatrix}$ c) $\begin{bmatrix} -2/3 & 3 \\ 1/3 & -1 \end{bmatrix}$ d) $\begin{bmatrix} 2 & 4 \\ 1 & 3 \end{bmatrix}$ [Mark 1]
- (vi) If B is a square matrix of order 3 with $\det(B) = 2$, then $\text{nullity}(B)$ is equal to:
 a) 2 b) 1 c) 3 d) ☒ 0 [Mark 1]
- (vii) If $\langle \cdot, \cdot \rangle$ is an inner product on $\mathbb{R}^n$ and $u, v \in \mathbb{R}^n$ such that $\|u\|^2 = 5$, $\|v\|^2 = 1$, $\langle u, v \rangle = -2$, then $\langle u + 2v, 5u - v \rangle$ is equal to:
 a) $\sqrt{5}$ b) ☒ 5 c) 9 d) 41 [Mark 1]
- (viii) If $S = \{A, I_2\} \subseteq M_{2 \times 2}(\mathbb{R})$, where A is a non-symmetric matrix, then S must be:
 a) linearly dependent b) a spanning set for $M_{2 \times 2}(\mathbb{R})$ c) ☒ linearly independent d) orthogonal [Mark 1]
- b) Let T be the transformation from the Euclidean space $\mathbb{R}^2$ to $\mathbb{R}$ given by $T(u) = \|u\|$ for all $u \in \mathbb{R}^2$, where $\|u\|$ is the Euclidean norm of u . Then, for $v, w \in \mathbb{R}^2$ and $k \in \mathbb{R}$, T satisfies:
 a) $T(u + v) = T(u) + T(v)$ b) ☒ $T(u + v) \leq T(u) + T(v)$ c) $T(0) > 0$ d) $T(ku) = kT(u)$ [Mark 1]
- (x) Zero is an eigenvalue of the matrix $\begin{bmatrix} 4 & 4 & 4 \\ 4 & 4 & 4 \\ 4 & 4 & 4 \end{bmatrix}$ with geometric multiplicity equal to:
 a) 1 b) ☒ 2 c) 3 d) 4 [Mark 1]

Question 2 [Marks 2 + 2 + 3]:

- (a) Find the square matrix A of order 3 such that $A^{-1}(A - I) = \begin{bmatrix} 1 & 1 & 1 \\ 2 & 1 & 1 \\ 1 & 1 & 2 \end{bmatrix}$ and evaluate $\det(A)$.

Solution: $I - A^{-1} = A^{-1}(A - I) = \begin{bmatrix} 1 & 1 & 1 \\ 2 & 1 & 1 \\ 1 & 1 & 2 \end{bmatrix} \Rightarrow A^{-1} = \begin{bmatrix} 0 & -1 & -1 \\ -2 & 0 & -1 \\ -1 & -1 & -1 \end{bmatrix} \Rightarrow \det(A^{-1}) = -1$, [Mark 1]

and $A = (A^{-1})^{-1} = \begin{bmatrix} 1 & 0 & -1 \\ 1 & 1 & -2 \\ -2 & -1 & 2 \end{bmatrix}$. [Mark 1]

- (b) Let $A = \begin{bmatrix} 1 & 0 & -1 \\ 1 & 1 & -2 \\ -2 & -1 & 2 \end{bmatrix}$ and $B = \begin{bmatrix} -2 & 1 & 1 \\ -1 & 1 & -2 \\ 1 & -1 & -2 \end{bmatrix}$. Find a matrix X that satisfies $XA = B$.

Solution: From Part (a), $A^{-1} = \begin{bmatrix} 0 & -1 & -1 \\ -2 & 0 & -1 \\ -1 & -1 & -1 \end{bmatrix}$; hence, $X = BA^{-1} = \begin{bmatrix} -3 & 1 & 0 \\ 0 & 3 & 2 \\ 4 & 1 & 2 \end{bmatrix}$. [Marks 0.5 + 1 + 0.5]

- (c) Solve the following system of linear equations:

$$\begin{aligned} x + y + z &= 1 \\ 2x + 2z &= 3 \\ 3x + 5y + 4z &= 2. \end{aligned}$$

Solution: The matrix of coefficient $A = \begin{bmatrix} 1 & 1 & 1 \\ 2 & 0 & 2 \\ 3 & 5 & 4 \end{bmatrix}$ has the inverse $A^{-1} = \frac{1}{-2} \begin{bmatrix} -10 & 1 & 2 \\ -2 & 1 & 0 \\ 10 & -2 & -2 \end{bmatrix}$. [Marks 1.5]

Hence, $\begin{bmatrix} x \\ y \\ z \end{bmatrix} = A^{-1} \begin{bmatrix} 1 \\ 3 \\ 2 \end{bmatrix} = \begin{bmatrix} \frac{3}{2} \\ -\frac{1}{2} \\ 0 \end{bmatrix}$. [Marks 1.5]

[The students may use any one of the methods included in the course MATH-244.]

Question 3 [Marks 3 + 3 + 2]:

Let $A = \begin{bmatrix} 1 & 0 & 1 \\ 1 & 2 & 3 \\ 2 & 0 & 2 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 2 & 2 \\ 0 & 0 & 0 \end{bmatrix}$. Then:

- (a) Find a basis and the dimension for each of the vector spaces $\text{row}(A)$, $\text{col}(A)$, and $N(A)$.

Solution: $RREF(A) = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{bmatrix}$. Hence, $\{(1,0,1), (0,1,1)\}$, $\{(1,1,2), (0,2,0)\}$, $\{(1,1,-1)\}$ are bases of $\text{row}(A)$, $\text{col}(A)$, $N(A)$, respectively, and so, $\dim(\text{row}(A)) = 2 = \dim(\text{col}(A))$, $\dim(N(A)) = 1$. [Marks 2]

- (b) Decide with justification whether the following statements are true or false:

(i) $\text{row}(A) = \text{row}(B)$

(ii) $\text{col}(A) = \text{col}(B)$

(iii) $N(A) = N(B)$.

Solution: $RREF(A) = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{bmatrix} = RREF(B) \Rightarrow \text{row}(A) = \text{row}(B)$ and $N(A) = N(B)$. [Marks 1 + 1]

But, $\text{col}(A) \neq \text{col}(B)$ since $(1,1,2) \notin \text{span}(\{(1,0,0), (0,2,0), (1,2,0)\})$. [Mark 1]

- (c) Find all square matrices Z of order 3 such that $AZ = O$.

Solution: From Part (a), $\{(1,1,-1)\}$ is a basis of the null space $N(A) = \{X \in \mathbb{R}^3 \mid AX = 0\}$. Hence,

$Z = \begin{bmatrix} a & b & c \\ a & b & c \\ -a & -b & -c \end{bmatrix}$ satisfies $AZ = aA \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix} + bA \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix} + cA \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix} = aO + bO + cO = O$, for all $a, b, c \in \mathbb{R}$. [Marks 0.5 + 1.5]

Question 4 [Marks 3 + (1 + 3)]:

- (a) Construct an orthonormal basis C of the Euclidean space $\mathbb{R}^3$ by applying the Gram-Schmidt algorithm on the given basis $B = \{v_1 = (1,1,0), v_2 = (1,0,1), v_3 = (0,1,1)\}$, and then find the coordinate vector of $v = (1,2,0) \in \mathbb{R}^3$ relative to the orthonormal basis C .

Solution: $u_1 = v_1 = (1,1,0)$; $u_2 = v_2 - \frac{\langle v_2, u_1 \rangle}{\|u_1\|^2} u_1 = (\frac{1}{2}, -\frac{1}{2}, 1)$; $u_3 = v_3 - \frac{\langle v_3, u_1 \rangle}{\|u_1\|^2} u_1 - \frac{\langle v_3, u_2 \rangle}{\|u_2\|^2} u_2 = (-\frac{2}{3}, \frac{2}{3}, \frac{2}{3})$.
Hence, $C = \{w_1 = \frac{1}{\sqrt{2}}(1,1,0), w_2 = \frac{1}{\sqrt{6}}(1,-1,2), w_3 = \frac{1}{\sqrt{3}}(-1,1,1)\}$ is the required orthonormal basis of $\mathbb{R}^3$.

Next, $\langle v, w_1 \rangle = \frac{3}{\sqrt{2}}$, $\langle v, w_2 \rangle = \frac{-1}{\sqrt{6}}$, and $\langle v, w_3 \rangle = \frac{1}{\sqrt{3}}$. Hence, $[v]_C = \begin{bmatrix} \frac{3}{\sqrt{2}} \\ \frac{-1}{\sqrt{6}} \\ \frac{1}{\sqrt{3}} \end{bmatrix}$. [Marks 1.5 + 0.5 + 1]

(b) Let $\mathcal{P}_2$ denote the vector space of real polynomials with degree ≤ 2 . Consider the linear transformation $T: \mathbb{R}^3 \rightarrow \mathcal{P}_2$ defined by: $T(1, 0, 0) = x^2 + 1$, $T(0, 1, 0) = 3x^2 + 2$, $T(0, 0, 1) = -x^2$. Then:

- (i) Compute $T(a, b, c)$, for all $(a, b, c) \in \mathbb{R}^3$.
- (ii) Find a basis for each of the vector spaces $Im(T)$ and $ker(T)$.

Solution: (i) $T(a, b, c) = aT(1, 0, 0) + bT(0, 1, 0) + cT(0, 0, 1) = (a + 3b - c)x^2 + a + 2b$. [Mark 1]

(ii) From Part (i), $Im(T) = \{(a + 3b - c)x^2 + (a + 2b)1 | (a, b, c) \in \mathbb{R}^3\} = span(\{x^2, 1\})$, [Mark 1]

and $ker(T) = \{(a, b, c) \in \mathbb{R}^3 | (a + 3b - c)x^2 + (a + 2b)1 = 0\}$

$= \{(a, b, c) \in \mathbb{R}^3 | a + 3b - c = 0, a + 2b = 0\}$

$= \{(a, b, c) \in \mathbb{R}^3 | a = -2b, b = c\}$

$= span(\{(-2, 1, 1)\})$.

Hence, $\{1, x^2\}$ and $\{(-2, 1, 1)\}$ are bases of $Im(T)$ and $ker(T)$, respectively.

[Mark 1]

[Mark 1]

Question 5 [Marks 3 + 2 + 3]: Let $A = \begin{bmatrix} 2 & 2 & -2 \\ 2 & 1 & -1 \\ 2 & 2 & -2 \end{bmatrix}$. Then:

- (a) Find the eigenvalues of A .

Solution: $\det(A - \lambda I) = \det \begin{pmatrix} 2-\lambda & 2 & -2 \\ 2 & 1-\lambda & -1 \\ 2 & 2 & -2-\lambda \end{pmatrix} = \lambda(\lambda + 1)(2 - \lambda) = 0$
 $\Rightarrow \lambda = -1, 0, 2$ are eigenvalues of A .

[Marks 1+1+1]

- (b) Find algebraic and geometric multiplicities of all the eigenvalues of A .

Solution: From Part (a), all eigenvalues of A are of same algebraic multiplicity 1. [Mark 0.5]

Next, $E_{-1} = span(\{(2, -1, 2)\})$, $E_0 = span(\{(0, 1, 1)\})$, and $E_2 = span(\{(1, 1, 1)\})$. Hence, all eigenvalues of A are of same geometric multiplicity 1. [Marks 1.5]

- (c) Is the matrix A diagonalizable? If yes, find a matrix P that diagonalizes A .

Solution: Since eigenvalues of A are different, A is diagonalizable. Next, from Part (b), the required matrix:

$$P = \begin{bmatrix} 2 & 0 & 1 \\ -1 & 1 & 1 \\ 2 & 1 & 1 \end{bmatrix}$$

[Marks 1.5 + 1.5]
