

## Example 1

A hypothesis test is to be performed to determine whether the mean waiting time during peak hours for customers in a supermarket has increased from the previous mean waiting time of 8.2 minutes. Previous experience indicates that the waiting time follows a normal distribution with standard deviation equal 3.8 minutes. To test the hypothesis, a random sample of 25 customers will be selected yields mean  $\bar{x} = 9.75$ .. **Answer the questions 1 to 8.**

**Question 1:**

The null and alternative hypotheses are...

(A)  $H_0 : \mu \geq 8.2$  &  $H_1 : \mu < 8.2$

(B)  $H_0 : \mu = 8.2$  &  $H_1 : \mu \neq 8.2$

(C)  $H_0 : \mu \leq 8.2$  &  $H_1 : \mu > 8.2$

(D)  $H_0 : \bar{X} \leq 8.2$  &  $H_1 : \bar{X} > 8.2$

**Question 2:**

This hypothesis test is classified as...

(A) Right-tailed

(B) Two-tailed

(C) Multi-tailed

(D) left-tailed

**Question 3:**

The appropriate test statistic is...

(A)  $Z = \frac{\bar{X} - \mu}{S/\sqrt{n}}$





(B)  $Z = \frac{\bar{X} - \mu}{\sigma/\sqrt{n}}$

(C)  $T = \frac{\bar{X} - \mu}{S/\sqrt{n}}$

(D)  $F = \frac{\bar{X} - \mu}{\sigma/\sqrt{n}}$

**Question 4:**

The critical region is best described by figure....



**Question 5:**

With significance level equal 0.10, the decision criterion for the hypothesis test in terms of the computed value of the test statistic is....

- |   |  |
|---|--|
| (A) Reject $H_0$ if $z_{stat} < -1.645$                       | (B) Reject $H_0$ if $z_{stat} > 1.96$  |
| (C) Reject $H_0$ if $Z_{stat} > 1.645$ or $Z_{stat} < -1.645$ | (D) Reject $H_0$ if $z_{stat} > 1.645$ |

**Question 6:**

The computed value of our test statistic is....

- |           |          |          |          |
|-----------|----------|----------|----------|
| (A) -2.04 | (B) 3.98 | (C) 2.04 | (D) 0.54 |
|-----------|----------|----------|----------|

**Solution:**

$$z_{stat} = \frac{9.75 - 8.2}{3.8 / \sqrt{25}} = 2.04$$

**Question 7:**

The decision would be to....

- |  |
|--|
| (A) Cannot be determined               |
| (B) Do not reject the null hypothesis. |
| (C) Reject the null hypothesis.        |
| (D) Reject the alternative hypothesis. |

**Question 8:**

Suppose that in fact the waiting time is ( $\mu_1 = 9.9$ ), then the decision has been made is...

- |                                      |                                     |
|--------------------------------------|-------------------------------------|
| (A) Committing Type I error          | (B) Committing Type II error        |
| (C) Correct decision( $1 - \alpha$ ) | (D) Correct decision( $1 - \beta$ ) |

**End of example 1**

## Example 2

It assumed from last experience that 75% of sports viewers are male. A famous sport newspaper reports that this proportion is greater than from 0.75. A random sample of 400 season ticket holders reveals that 352 are male. We wish to test the above hypothesis. **Answer the questions 1 to 9.**

### Question 1:

The null and alternative hypotheses are...

(A) $H_0: P \leq 0.75$ & $H_1: P > 0.75$	(B) $H_0: \pi < 0.75$ & $H_1: \pi \geq 0.75$
(C) $H_0: \pi \leq 0.75$ & $H_1: \pi > 0.75$	(D) $H_0: \pi = 0.75$ & $H_1: \pi \neq 0.75$

### Question 2:

This hypothesis test is classified as...

(A) Two-tailed	(B) Right-tailed
(C) Opposite-tailed	(D) left-tailed

### Question 3:

The appropriate test statistic is...

(A) $Z = \frac{P - \pi}{\sqrt{\pi(1-\pi)}/n}$	(B) $T = \frac{\bar{X} - \mu}{S/\sqrt{n}}$
(C) $Z = \frac{\bar{X} - \mu}{\sigma/\sqrt{n}}$	(D) $\chi^2 = \frac{P - \pi}{\sqrt{\pi(1-\pi)}/n}$


### Question 4:

With significance level equal 0.10, the decision criterion for the hypothesis test in terms of the computed value of the test statistic ( $Z_{stat}$ ) is....

(A) Reject $H_0$ if $z_{stat} < -1.645$	(B) Reject $H_0$ if $z_{stat} > 1.96$
(C) Reject $H_0$ if $Z_{stat} > 1.645$ or $Z_{stat} < -1.645$	(D) Reject $H_0$ if $z_{stat} > 1.645$

**Question 5:**

With level of significance 5%, the critical region is best described by figure....



**Question 6:**

The computed value of our test statistic is....

- (A) 0.01    (B) 5.99    (C) 0.23    (D) -0.01

**Solution:**

$$z_c = \frac{352/400 - 0.75}{\sqrt{(0.75)(0.25)/400}} = \frac{0.88 - 0.75}{0.0217} = \frac{0.88 - 0.75}{0.0217} = 5.99$$

**Question 7:**

The decision would be to....

- (A) Do not Reject the null hypothesis
- (B) Cannot be determined.
- (C) Reject the null hypothesis.
- (D) Reject the alternative hypothesis.

**Question 8:**

Suppose that in fact the true proportion is 0.85, then the decision has been made is...  $\alpha$

- |  |   |
|--|---|
| (A) Rejecting the true hypothesis( $\alpha$ ) type1 error                | (B) Do not Rejecting the false hypothesis ( $\beta$ ) type11 error. |
| (C) Do not rejecting the true hypothesis( $1 - \alpha$ )Correct decision | (D) Rejecting the false hypothesis( $1 - \beta$ )Correct decision   |

**Question 9:**

Suppose that in fact the true proportion is 0.74, then the decision has been made is...

(A) Rejecting the true hypothesis( $\alpha$ ) type1 error	(B)Do not Rejecting the false hypothesis ( $\beta$ ) type11 error.
(C) Do not rejecting the true hypothesis( $1-\alpha$ )Correct decision	(D) Rejecting the false hypothesis( $1-\beta$ )Correct decision

**End of example 2**

### Example 3

**Question 1:**

A 95% confidence interval is  $12 < \mu < 17$  null hypothesis is  $H_0: \mu = 10$   
 $H_1: \mu \neq 10$

What is the decision?

- |  |
|--|
| (A) Reject the null hypothesis.        |
| (B) Do not Reject the null hypothesis. |
| (C) Can not be determined              |
| (D) Reject the alternative hypothesis. |

**Question2:** A 95% confidence interval is  $0.85 \leq \pi \leq 0.91$ . The null hypothesis is  
 $H_0: \pi = 0.88$   
 $H_1: \pi \neq 0.88$

What is the decision?

- |   |
|---|
| (A) Reject the null hypothesis.               |
| (B) Do not Reject the null hypothesis.        |
| (C) Cannot be determined                      |
| (D) Do not Reject the alternative hypothesis. |

**End of example 3**