

**CHEM 108**

# **FUNDAMENTALS OF ORGANIC CHEMISTRY**

FOR B.Sc. PROGRAMS OF SCIENTIFIC COLLEGES

PRE-REQUISITES COURSE; CHEM 101

CREDIT HOURS; 4 (3+1)

Chemistry Department, College of Science, King Saud University

1  
Dr Mohamed El-Newehy

## **CHAPTER 10**

# **Fats, Lipids & Oils**

2  
Dr Mohamed El-Newehy

## Lipids

- Family of biomolecules.
- **Lipids** are biological molecules (biomolecules) that are insoluble in water but soluble in nonpolar solvents (nonpolar).
- They are **not defined** by a particular functional group, thus they have a **variety** of structures and functions.
- They are soluble in organic solvents but not in water (nonpolar).
- They contain many nonpolar C—C and C—H bonds and few polar bonds resulting in their water insolubility.




3

Dr Mohamed El-Newehy

## Lipids

- **Lipids** are biological molecules that are insoluble in water but soluble in nonpolar solvents (nonpolar).
- The word **lipid** comes from the Greek lipos, which means “fat.”
- **Lipids** are the waxy, greasy, or oily compounds found in plants and animals.
  - wax coating that protects plants
  - used as energy storage
  - structural components (cell membranes)
  - insulation against cold



4

Dr Mohamed El-Newehy

## Classification of Lipids

Lipids are divided into:

### Saponifiable (Hydrolyzable) lipids

- They contain esters,
- They can undergo saponification  
(hydrolysis under basic conditions)

#### Simple lipids

contain two components  
(fatty acid and an alcohol)

Triglycerides  
(Fats & oils)

Waxes

#### Complex lipids

contain more than two components  
(fatty acids, an alcohol, and other components)

phosphoglycerides

Sphingolipids.

### Nonsaponifiable (Nonhydrolyzable) lipids

- They do not contain ester groups,
- They cannot be saponified  
(cannot be cleaved into smaller molecules by aqueous hydrolysis)


steroids

prostaglandins

5  
Dr Mohamed El-Newehy

## Fatty Acids

- **Fatty acids** are long-chain unbranched carbon attached to a carboxyl group (-COOH).




Dr Mohamed El-Newehy

## Fatty Acids

### Characteristics of Fatty Acids:

- They are usually have **straight chains** (no branches) that are about **10 to 20 carbon atoms** in length.
- They usually have an **even number of carbon atoms** (counting the carboxyl carbon).
- The carbon chains may be;
  - **saturated** (all single bonds) or
  - **unsaturated** (containing double bonds).
- i.e. **Functional groups** are only the **carboxyl group** and the **double bonds**.
- The double bonds are usually in **cis configurations**:



Dr Mohamed El-Newehy 7

## Fatty Acids


### Saturated and Unsaturated Fatty Acids

- **Saturated fatty acids** have no double bonds in their long hydrocarbon chains.

**Stearic acid:**  $\text{CH}_3(\text{CH}_2)_{16}\text{COOH}$

Stearic Acid  
(m.p. 71°C)

They are solids at room temperature.



**Stearic acid** is found in palm oil, which is frequently used in handmade soap.




Dr Mohamed El-Newehy 8

## Fatty Acids

### Saturated and Unsaturated Fatty Acids

- **Unsaturated fatty acids** have 1 or more double bonds (generally *cis*) in their long hydrocarbon chains.
  - **Oleic acid** (one double bond), and
  - **Linoleic acid** (two double bonds)
  - All have 18 carbons in the chain, but their melting points are different
- **Oleic acid** is derived mainly from “olive” oil, sesame oil, Sunflower, shea butter, coconut oil, ... etc.
- **Linoleic acid** is found in soybean oil




Dr Mohamed El-Newehy

9

## Fatty Acids

### Essential Fatty Acids

- Most of the fatty acids we need can be synthesized in the body.
- Two fatty acids, **linoleic acid** and **linolenic acid**, both polyunsaturated fatty acids with 18-carbon chains, cannot be synthesized in the body and must be obtained from the diet.
- **These are essential fatty acids.**
- Both are found in plant and fish oils.
- In the body, they are used to produce hormone-like substances that regulate blood pressure, blood clotting, blood lipid levels, the immune response, and inflammatory reactions.



Dr Mohamed El-Newehy

10

## Fatty Acids


### Essential Fatty Acids

- o All fatty acids that bear the “omega” label are unsaturated, containing one or more double bonds.

**Omega-n acids**

n: the position of the first double bond

**Linoleic acid** is called an **omega-6 acid**, because of the position of the first C=C in the nonpolar chain.




Dr Mohamed El-Newehy

## Fatty Acids

### Essential Fatty Acids

**Linolenic acid** is called an **omega-3 acid**, because of the position of the first C=C in the nonpolar chain.



12

Dr Mohamed El-Newehy

## Fatty Acids

### Some Important Fatty Acids


#C's	Name	Formula	MP	Common Sources
<b>Saturated</b>				
14	Myristic acid	$\text{CH}_3(\text{CH}_2)_{12}\text{COOH}$	54°C	Butterfat, coconut oil, nutmeg oil
16	Palmitic acid	$\text{CH}_3(\text{CH}_2)_{14}\text{COOH}$	63°C	Lard, beef fat, butterfat, cottonseed oil
18	Stearic acid	$\text{CH}_3(\text{CH}_2)_{16}\text{COOH}$	70°C	Lard, beef fat, butterfat, cottonseed oil
20	Arachidic acid	$\text{CH}_3(\text{CH}_2)_{18}\text{COOH}$	76°C	Peanut oil
<b>Monounsaturated</b>				
16	Palmitoleic acid	$\text{CH}_3(\text{CH}_2)_5\text{CH}=\text{CH}(\text{CH}_2)_7\text{COOH}$	-1°C	Cod liver oil, butterfat
18	Oleic acid	$\text{CH}_3(\text{CH}_2)_7\text{CH}=\text{CH}(\text{CH}_2)_7\text{COOH}$	13°C	Lard, beef fat, olive oil, peanut oil
<b>Polyunsaturated</b>				
18	Linoleic acid	$\text{CH}_3(\text{CH}_2)_4(\text{CH}=\text{CHCH}_2)_2(\text{CH}_2)_6\text{COOH}$	-5°C	Cottonseed oil, soybean oil, corn oil, linseed oil
18	Linolenic acid	$\text{CH}_3\text{CH}_2(\text{CH}=\text{CHCH}_2)_3(\text{CH}_2)_6\text{COOH}$	-11°C	Linseed oil, corn oil
20	Arachidonic acid	$\text{CH}_3(\text{CH}_2)_4(\text{CH}=\text{CHCH}_2)_4(\text{CH}_2)_2\text{COOH}$	-50°C	Corn oil, linseed oil, animal tissues
20	Eicosapentaenoic acid	$\text{CH}_3\text{CH}_2(\text{CH}=\text{CHCH}_2)_5(\text{CH}_2)_2\text{COOH}$		Fish oil, seafoods
22	Docosahexaenoic acid	$\text{CH}_3\text{CH}_2(\text{CH}=\text{CHCH}_2)_6\text{CH}_2\text{COOH}$		Fish oil, seafoods

13  
Dr Mohamed El-Newehy

## Simple Lipids; Triglycerides

- Animal fats and vegetable oils are esters composed of three molecules of a fatty acid connected to a glycerol molecule, producing a structure called a *triglyceride* or a *triacylglycerol*:


### Esterification




Dr Mohamed El-Newehy

## Simple Lipids; Triglycerides

- Simple triacylglycerols have three identical fatty acid side chains.




- Mixed triacylglycerols have two or three different fatty acids.



Dr Mohamed El-Newehy

## Simple Lipids; Fats & Oils

- The fatty acids in a triglyceride molecule are usually not all the same;
- Natural triglycerides are often mixtures of different triglyceride molecules.



### Fats are



- triglycerides that are solids at room temp.
- usually derived from animals
- mostly saturated fatty acids



### Oils are


- triglycerides that are liquids at room temp.
- usually derived from plants or fish
- mostly unsaturated fatty acids

Dr Mohamed El-Newehy



## Simple Lipids; Fats & Oils

A comparison of saturated and unsaturated fatty acids in some foods.




17  
Dr Mohamed El-Newehy

## Simple Lipids; Fats & Oils

### Chemical Properties of Fats and Oils


#### Hydrolysis of Triglycerides

Triglycerides can be broken apart with water and an acid catalyst (*hydrolysis*), or by digestive enzymes called lipases:



#### Saponification of Triglycerides (Basic Hydrolysis)

Triglycerides react with strong bases (NaOH or KOH) to form the carboxylate salts of the fatty acids, called soaps:




18  
Dr Mohamed El-Newehy

## Simple Lipids; Fats & Oils

### Chemical Properties of Fats and Oils

#### Soaps

- NaOH produces a “hard” soap, commonly found in bar soaps;
- KOH produces a “soft” soap, such as those in shaving creams and liquid soaps.
- These salts combine two solubility characteristics:
  - a long, nonpolar, water-insoluble (*hydrophobic*) hydrocarbon “tail.”
  - a charged, water-soluble (*hydrophilic*) “head.”




Dr Mohamed El-Newehy

## Simple Lipids; Fats & Oils

### Chemical Properties of Fats and Oils

#### Soaps

In water, the “tails” become tangled, leaving the charged heads sticking out into the solution, forming a structure called a **micelle**.


20  
Dr Mohamed El-Newehy

## Simple Lipids; Fats & Oils

### Chemical Properties of Fats and Oils

#### Hydrogenation


- Alkenes are converted into alkanes with hydrogen gas ( $H_2$ ) and a catalyst (Pt, Ni, or some other metal).
- This process is used to convert unsaturated vegetable oils, which are liquids at room temp., to saturated fats, which are solids at room temp.




Dr Mohamed El-Newehy

## Simple Lipids; Waxes

- Waxes** are simple lipids contain a fatty acid joined to a long-chain (12-32 carbons) alcohol:



- Because of their long nonpolar C chains, waxes are very hydrophobic.



22

Dr Mohamed El-Newehy

## Simple Lipids; Waxes

- Waxes are insoluble in water, and not as easily hydrolyzed as fats and oils.
- Waxes often occur in nature as protective coatings on feathers, fur, skin, leaves, and fruits.
- Waxes are used commercially to make cosmetics, candles, ointments, and protective polishes.

Beeswax



Lanolin from wool



lotions


23

Dr Mohamed El-Newehy

## Complex Lipids; Phospholipids



- Phospholipids are lipids that contain a P atom.
- There are two common types:

### Phosphoglycerol



### Sphingolipids

Sphingolipids are complex lipids that contain sphingosine instead of glycerol.




Dr Mohamed El-Newehy

## Complex Lipids; Phospholipids

### Phosphoglycerols

- **Phosphoglycerols** are complex lipids that are *major components of cell membranes*.
- Structurally, they resemble a triacylglycerol, except the third fatty acid has been replaced with a **phosphodiester bonded to an alcohol**.




25  
Dr Mohamed El-Newehy


## Complex Lipids; Phospholipids

### Phosphoglycerols

- There are two types of phosphoglycerols;



- **Cephalins** are found in most cell membranes,
- They are particularly abundant in brain tissue.
- They are also found in blood platelets, and play a role in blood clotting.




- Lecithin** can act as an emulsifying agent:
  - important structural component of cell membranes.
  - play a role in the transport of lipids in the blood stream.
  - Commercially, lecithin extracted from soybeans is used as an emulsifying agent in margarine and candies to provide a smooth texture.

Dr Mohamed El-Newehy


## Complex Lipids; Phospholipids

### Sphingomyelins

- They do not contain a glycerol backbone, they have a sphingosine backbone instead.



- They **do not contain an ester**; their single fatty acid is bonded to the backbone by an **amide bond**.




- Sphingomyelins** are found brain and nerve tissue, and in the myelin sheath that protects nerves.

Dr Mohamed El-Newehy

## Complex Lipids; Phospholipids

### Glycolipids

- Glycolipids** are sphingolipids that contain carbohydrates (usually monosaccharides).
- They are also referred to as *cerebrosides* because of their abundance in brain tissue.




28

Dr Mohamed El-Newehy

## Cell Membrane

### Fluid mosaic model




The cell membrane as being composed of a lipid bilayer, in which the nonpolar tails of lipids point towards the “interior” of the bilayer, leaving the polar, hydrophilic portions pointing outwards.

**Semipermeable:** selected nutrients can enter and waste products can leave.

29  
Dr Mohamed El-Newehy

## Nonsaponifiable (Nonhydrolyzable) Lipids; Steroids

- **Steroids** are classified as lipids because they are soluble in nonpolar solvents.
- They are nonsaponifiable because the components are not held together by ester linkages.
- The basic steroid structure contains four fused rings:
- **Steroids** have
  - nucleus which is 4 carbon rings.
  - attached groups that make the different types of compounds.
  - no fatty acids.




steroid ring system

30  
Dr Mohamed El-Newehy

## Nonsaponifiable (Nonhydrolyzable) Lipids; Steroids

### Cholesterol

- **Cholesterol** is the most abundant steroid in the body.
- It is an essential component of cell membranes, and is a precursor for other steroids, such as the bile salts, sex hormones, vitamin D, and the adrenocorticoid hormones.
- There is apparently a correlation between high levels of cholesterol in the blood and atherosclerosis.



Cholesterol is obtained from meats, milk, and eggs.

Cholesterol is synthesized in the liver from fats, carbohydrates and proteins.


No cholesterol in vegetable and plants.

31  
Dr Mohamed El-Newehy

## Nonsaponifiable (Nonhydrolyzable) Lipids; Steroids

### Bile Salts

- **Bile** is a yellowish brown or green fluid produced in the liver and stored in the gall bladder.
- **Bile salts** act like soaps and other emulsifiers: they contain both polar and nonpolar regions, helping to break fats in foods into smaller pieces, allowing them to be hydrolyzed more easily.



32  
Dr Mohamed El-Newehy



## Nonsaponifiable (Nonhydrolyzable) Lipids; Steroids

### Steroid Hormones

- A **hormone** is a molecule that is synthesized in one part of an organism, which then elicits a response at a different site.

- Two types of steroids hormones:

#### 1. Sex hormones

Estrogens & progestins in females

Androgens in males

#### 2. Adrenal Cortical Steroids

33  
Dr Mohamed El-Newehy



## Nonsaponifiable (Nonhydrolyzable) Lipids; Steroids

### Steroid Hormones

#### Female Sex Hormones


##### Estrogens

The estrogens **estradiol** and **estrone** control development of secondary sex characteristics, *regulate the menstrual cycle*, and are made in the ovaries.



##### Progestins

The progestin **progesterone** is called the “pregnancy hormone”; it is responsible for the *preparation of the uterus for implantation of a fertilized egg*.



34  
Dr Mohamed El-Newehy

## Nonsaponifiable (Nonhydrolyzable) Lipids; Steroids



### Steroid Hormones

#### Male Sex Hormones

#### Androgens

Testosterone and Androsterone are androgens made in the testes.

They control the development of secondary sex characteristics in males.






35  
Dr Mohamed El-Newehy

## Nonsaponifiable (Nonhydrolyzable) Lipids; Steroids

### Steroid Hormones

#### Adrenocorticoid Hormones

- The **adrenocorticoid hormones** are produced in the adrenal glands (located on the top of the kidney).
- **Glucocorticoids** such as cortisol affect the metabolism of carbohydrates.
- **Cortisol and its derivatives, cortisone and prednisolone** (synthetic) are powerful anti-inflammatory drugs used to treat arthritis and asthma.



36  
Dr Mohamed El-Newehy


## Nonsaponifiable (Nonhydrolyzable) Lipids; Steroids

### Steroid Hormones

#### Adrenocorticoid Hormones

Mineralocorticoids regulate ion concentration (mainly  $\text{Na}^+$ ).


**Aldosterone** influences the absorption of  $\text{Na}^+$  and  $\text{Cl}^-$  in kidney tubules, thus regulating the retention of water in the body.



37  
Dr Mohamed El-Newehy

## Nonsaponifiable (Nonhydrolyzable) Lipids; Prostaglandins

**Prostaglandins** are **carboxylic acids** that contain a **five-membered ring** and have a wide range of biological activities.



- **Prostaglandins** are responsible for **inflammation**.
- **Prostaglandins** also decrease gastric secretions, inhibit blood platelet aggregation, stimulate uterine contractions, and relax smooth muscles.
- Aspirin and ibuprofen relieve pain and inflammation by **blocking the synthesis** of these molecules.

38  
Dr Mohamed El-Newehy


## Vitamins

- They are organic compounds required in small quantities for normal metabolism.
- They must be obtained from the diet (our cells cannot synthesize them).
- Vitamins are either **water soluble** or **fat soluble**.
- The four fat-soluble vitamins (**A**, **D**, **E**, and **K**) are lipids and nonpolar.
- They are found in fruits, vegetables, fish, liver, and dairy products.
- Excess vitamins are **stored in adipose cells** to be used when needed.

39  
Dr Mohamed El-Newehy

## Vitamins; Vitamin A


- It is found in *liver*, *fish*, and *dairy products*, and is made from  $\beta$ -carotene (the orange pigment in carrots).
- It is needed for **vision** and for healthy mucous membranes.
- Vitamin A **deficiency** causes night blindness and dry eyes and skin.



40  
Dr Mohamed El-Newehy

## Vitamins; Vitamin D


- Vitamin D can be synthesized from cholesterol.
- It can be obtained in the diet from many foods, especially milk, and helps regulate Ca and P metabolism.
- A deficiency of vitamin D causes rickets (bone malformation).



41  
Dr Mohamed El-Newehy


## Vitamins; Vitamin E

- Vitamin E is an antioxidant, protecting unsaturated side chains in fatty acids from unwanted oxidation.
- Deficiency of vitamin E causes numerous neurological problems, although it is rare.



## Vitamins; Vitamin K

- Vitamin K regulates the synthesis of clotting proteins (prothrombin), and deficiency of this leads to excessive or fatal bleeding.



Dr Mohamed El-Newehy