



KING SAUD UNIVERSITY
College of Science
Department of Mathematics

M-106

First Semester (1431/1432)

Solution First Mid-Exam

Name:	Number:
Name of Teacher:	Group No:

Max Marks: 20

Time: 90 minutes

Marks:

Multiple Choice (1-10)	
Question # 11	
Question # 12	
Question # 13	
Question # 14	
Total	

Multiple Choice

Q.No:	1	2	3	4	5	6	7	8	9	10
$\{a, b, c, d\}$	d	a	c	b	c	d	b	b	d	a

Q. No: 1 Using the definition of Riemann Sum as a definite integral,

$\lim_{\|P\| \rightarrow 0} \sum_k (\omega_k)^4 \Delta x_k$, on $[0, 1]$ is equal to:

- (a) 0 (b) ∞ (c) 1 (d) $\frac{1}{5}$

Q. No: 2 The value of the integral $\int_0^2 x\sqrt{4-x^2}dx$ is equal to:

- (a) $\frac{8}{3}$ (b) $\frac{3}{8}$ (c) $\frac{-3}{8}$ (d) $\frac{-8}{3}$

Q. No: 3 The number z that satisfies the conclusion of the Mean value Theorem for $f(x) = x$ on $[\alpha, \beta]$ (where $\alpha < \beta$ are two constants) is:

- (a) α (b) $\beta + 1$ (c) $\frac{\alpha+\beta}{2}$ (d) β

Q. No: 4 The average value of $f(x) = |x - 1|$ on $[0, 2]$ is equal to:

- (a) 1 (b) $\frac{1}{2}$ (c) 0 (d) 2

Q. No: 5 If $F(x) = \int_1^{2x} f'(t) dt$, then $F'(x)$ is equal to:

- (a) $2f(2x) - f(1)$ (b) $2f(2x)$ (c) $2f'(2x)$ (d) $f'(2x)$

Q. No: 6 The derivative of the function $f(x) = \sec^{-1}(e^x)$ is equal to:

- (a) $\frac{1}{e^x\sqrt{e^x-1}}$ (b) $\frac{1}{\sqrt{e^x-1}}$ (c) $\frac{1}{\sqrt{e^{2x}+1}}$ (d) $\frac{1}{\sqrt{e^{2x}-1}}$

Q. No: 7 The value of the integral $\int_0^1 4^x dx$ is equal to:

- (a) $\frac{4}{\ln 4}$ (b) $\frac{3}{2\ln 2}$ (c) $4 \ln 4$ (d) $3 \ln 4$

Q. No: 8 To evaluate $\int \frac{x^3}{\sqrt{1-x^8}} dx$, we use the change of variable:

- (a) $u = x^2$ (b) $u = x^4$ (c) $u = x^8$ (d) $u = x^3$

Q. No: 9 The value of the integral $\int \frac{\sin x}{\sqrt{4-\cos^2 x}} dx$ is equal to:

- (a) $\sin^{-1}\left(\frac{\cos x}{2}\right) + c$ (b) $\sin^{-1}(\cos x) + c$ (c) $-\cos^{-1}\left(\frac{\sin x}{2}\right) + c$ (d) $\cos^{-1}\left(\frac{\cos x}{2}\right) + c$

Q. No: 10 The value of the integral $\int \frac{1}{\sqrt{4+x^2}} dx$ is equal to:

(a) $\sinh^{-1} \left(\frac{x}{2} \right) + c$ (b) $\sin^{-1} \left(\frac{x}{2} \right) + c$ (c) $\frac{1}{2} \sinh^{-1} \left(\frac{x}{2} \right) + c$ (d) $\frac{1}{2} \sin^{-1} \left(\frac{x}{2} \right) + c$

Full Questions

Question No: 11 Approximate the integral $\int_0^4 \sqrt{1+x^3} dx$ using the **Simpson's rule** with $n = 4$. [3]

Solution:

Let $f(x) = \sqrt{1+x^3}$.

$\Delta x = \frac{4}{4} = 1$

$x_0 = 0, x_1 = 1, x_2 = 2, x_3 = 3$ and $x_4 = 4$. (1)

$$\begin{aligned} \int_0^4 \sqrt{1+x^3} dx &\approx \frac{4-0}{3 \times 4} \{f(0) + 4f(1) + 2f(2) + 4f(3) + f(4)\} & (1) \\ &= \frac{1}{3} \{1 + 4(1.4142) + 2(3) + 4(5.2915) + 8.0623\} \\ &= \frac{1}{3} \{1 + 5.6568 + 6 + 21.166 + 8.0623\} \\ &= \frac{1}{3} \{41.885\} \approx 13.962 & (1) \end{aligned}$$

Question No: 12 If $y = (1+x^2)^{\sin x}$, then find y' . [2]

Solution:

$\ln y = \sin x \ln (1+x^2)$ (1)

$\frac{y'}{y} = \cos x \ln (1+x^2) + \frac{2x \sin x}{1+x^2}$

So $y' = \left[\cos x \ln (1+x^2) + \frac{2x \sin x}{1+x^2} \right] (1+x^2)^{\sin x}$ (1).

Question No: 13 Evaluate the integral $\int \frac{x+1}{\sqrt{4-x^2}} dx$. [2]

Solution:

$$\begin{aligned} \int \frac{x+1}{\sqrt{4-x^2}} dx &= \int \frac{x}{\sqrt{4-x^2}} dx + \int \frac{1}{\sqrt{4-x^2}} dx \\ &= -\sqrt{4-x^2} + \sin^{-1}\left(\frac{x}{2}\right) \end{aligned} \quad (1+1)$$

Question No: 14 Evaluate the integral $\int \frac{1}{x\sqrt{1-x^4}} dx$. [3]

Solution:

$$\text{Let } u = x^2 \Rightarrow du = 2x dx \quad (1)$$

$$\begin{aligned} \int \frac{1}{x\sqrt{1-x^4}} dx &= \frac{1}{2} \int \frac{1}{u\sqrt{1-u^2}} du \quad (1) \\ &= -\frac{1}{2} \operatorname{sech}^{-1}(u) + c \\ &= -\frac{1}{2} \operatorname{sech}^{-1}(x^2) + c \quad (1) \end{aligned}$$