


**CHEM 244**

# **PRINCIPLES OF ORGANIC CHEMISTRY I**

FOR CHEMICAL ENGINEERING' STUDENTS, COLLEGE OF ENGINEERING

PRE-REQUISITES COURSE; CHEM 101

CREDIT HOURS; 2 (2+0)



**Dr. Mohamed El-Newehy**

Chemistry Department, College of Science, King Saud University

<http://fac.ksu.edu.sa/melnewehy/home>

## **CHAPTER 4**

# **AROMATIC COMPOUNDS**


## THE STRUCTURE OF BENZENE RING

- **Benzene**,  $C_6H_6$ , is the parent hydrocarbon of the especially stable compounds known as **aromatic compounds**.
- The carbon-to-hydrogen ratio in **benzene**,  $C_6H_6$ , suggests a **highly unsaturated** structure.
- It does not undergo the typical addition reactions of alkenes or alkynes.
- Instead, **benzene** reacts *mainly* by **substitution reactions**.

## THE STRUCTURE OF BENZENE RING

### THE KEKULÉ STRUCTURE OF BENZENE


- He suggested that **six carbon atoms are located at the corners** of a **regular hexagon**, with one hydrogen atom attached to each carbon atom.
- To give each carbon atom a valence of 4, he suggested that **single and double bonds alternate around the ring** (what we now call a **conjugated system of double bonds**).
- *All of the* carbon–carbon **bond lengths are identical**: 1.39 Å, intermediate between typical single (1.54 Å) and double (1.34 Å) carbon–carbon bond lengths.



## THE STRUCTURE OF BENZENE RING

### ORBITAL MODEL FOR BENZENE

- Each carbon is therefore  $sp^2$ -hybridized,
- It also explains its hexagonal shape, with H – C – C and C – C – C angles of  $120^\circ$ .




An orbital representation of the bonding in benzene. Sigma ( $\sigma$ ) bonds are formed by the end-on overlap of  $sp^2$  orbitals.

In addition, each carbon contributes one electron to the pi ( $\pi$ ) system by lateral overlap of its  $p$  orbital with the  $p$  orbitals of its two neighbors.

### SYMBOLS FOR BENZENE

- Two symbols are used to represent benzene.
  - One is the Kekulé structure, and
  - The other is a hexagon with an inscribed circle, to represent the idea of a delocalized pi electron cloud.



Kekulé



delocalized pi cloud

## AROMATIC CHARACTER (AROMATICITY)






○ **Aromatic character (Aromaticity)** is associated with several structural requirements.

- 1) **Cyclic structure.**
- 2) Cyclic structure contains what looks like a continuous system of **alternating double and single bonds.**
- 3) Aromatic compounds must be **planar.**
- 4) Aromaticity is possible only if it **obeys Hückel's rule.**

*The number of  $\pi$  electrons in the compound =  $(4n + 2)$*

Where ( $n = 0, 1, 2, 3$ , and so on).

## AROMATIC CHARACTER (AROMATICITY)

$n$	$4n + 2$	Structure and name of aromatic compound				
1	6					

Benzene










Pyridine

Pyrrole

Furan


Thiophene

### Examples


				
$4n+2 = 8$	10	2	2	4
$n = 1.5$	2	0	0	0.5
				
$4n+2 = 4$	4	6	4	
$n = 0.5$	0.5	1	0.5	

## NOMENCLATURE OF AROMATIC COMPOUNDS

- **Monosubstituted benzenes** that do not have common names accepted by IUPAC are named as derivatives of benzene.




- **Monosubstituted benzenes** that have common names are accepted by IUPAC (parent compounds).



Dr Mohamed El-Newehy 9

## NOMENCLATURE OF AROMATIC COMPOUNDS


- When **two substituents** are present, *three isomeric structures are possible*.
- They are designated by the prefixes; *ortho- (o-)*, *meta- (m-)* and *para- (p-)*.
- If substituent X is attached to carbon 1; *o- groups* are on *carbons 2 and 6*, *m- groups* are on *carbons 3 and 5*, and *p- groups* are on *carbon 4*.




Dr Mohamed El-Newehy 10

## NOMENCLATURE OF AROMATIC COMPOUNDS


### ○ Examples;



- The prefixes; *ortho*- (*o*-), *meta*- (*m*-) and *para*- (*p*-) are used when the two substituents are not identical.




- When **more than two substituents** are present, their positions are designated by **numbering the ring**.



Dr Mohamed El-Newehy 11

## NOMENCLATURE OF AROMATIC COMPOUNDS

- Aromatic hydrocarbons, as a class, are called **arenes**.
- The symbol **Ar** is used for an **aryl group**, (symbol **R** is used for an alkyl group).
- Therefore, the formula **Ar - R** would represent any **arylalkane**.
- Two groups with special names occur frequently in aromatic compounds; the **phenyl group** and the **benzyl group**.




Dr Mohamed El-Newehy 12


## REACTIONS OF BENZENE

### A) ELECTROPHILIC AROMATIC SUBSTITUTION


#### 1) Halogenation




#### 4) Alkylation (Friedel-Crafts)




#### 2) Nitration



#### 5) Acylation (Friedel-Crafts)



#### 3) Sulfonation




Dr Mohamed El-Newehy 13

## REACTIONS OF BENZENE

### A) ELECTROPHILIC AROMATIC SUBSTITUTION

#### The Mechanism of Electrophilic Aromatic Substitution

We can generalize this two-step mechanism for all the electrophilic aromatic substitutions.




Dr Mohamed El-Newehy 14

## REACTIONS OF BENZENE

## A) ELECTROPHILIC AROMATIC SUBSTITUTION

## The Mechanism of Electrophilic Aromatic Substitution

## 1) Halogenation

15  
Dr Mohamed El-Newehy


## REACTIONS OF BENZENE

## A) ELECTROPHILIC AROMATIC SUBSTITUTION

## The Mechanism of Electrophilic Aromatic Substitution

## 2) Nitration

In aromatic nitration reactions, the **sulfuric acid catalyst** protonates the **nitric acid**, which then loses water to generate the **nitronium ion** ( $\text{NO}_2^+$ ), which contains a positively charged nitrogen atom.

16  
Dr Mohamed El-Newehy





## REACTIONS OF BENZENE

### A) ELECTROPHILIC AROMATIC SUBSTITUTION

#### The Mechanism of Electrophilic Aromatic Substitution

#### 3) Sulfonation

- We use either concentrated or *fuming sulfuric acid*, and the electrophile may be sulfur trioxide,  $\text{SO}_3$ , or *protonated sulfur trioxide*,  $^+\text{SO}_3\text{H}$ .
- Sulfuric acid provides catalyst as following:



17  
Dr Mohamed El-Newehy


## REACTIONS OF BENZENE

### A) ELECTROPHILIC AROMATIC SUBSTITUTION

#### The Mechanism of Electrophilic Aromatic Substitution

#### 4) Alkylation (Friedel-Crafts)

The *electrophile is a carbocation*, which can be formed either by removing a halide ion from an *alkyl halide* with a *Lewis acid catalyst* (for example,  $\text{AlCl}_3$ ).



18  
Dr Mohamed El-Newehy

## REACTIONS OF BENZENE


### A) ELECTROPHILIC AROMATIC SUBSTITUTION

#### The Mechanism of Electrophilic Aromatic Substitution

##### 5) Acylation (Friedel-Crafts)

The *electrophile is an acyl cation* generated from an acid derivative, usually an *acyl halide*.

The reaction provides a useful general route to aromatic ketones.




19  
Dr Mohamed El-Newehy


### DISUBSTITUTED BENZENES: ORIENTATION

Substituents present on an aromatic ring determine the position taken by a new substituent.

- Nitration of toluene gives mainly a mixture of *o*- and *p*-nitrotoluene.



- On the other hand, nitration of nitrobenzene under similar conditions gives mainly the *meta* isomer.



20  
Dr Mohamed El-Newehy

## DISUBSTITUTED BENZENES: ORIENTATION & REACTIVITY


### Directing and Activating Effects of Common Functional Groups

	Substituent group	Name of group	
Ortho, Para-Directing	$-\text{NH}_2, -\text{NHR}, -\text{NR}_2$	amino	Activating
	$-\text{OH}, -\text{OCH}_3, -\text{OR}$	hydroxy, alkoxy	
	$-\text{NHC}(=\text{O})-\text{R}$	acylamino	
	$-\text{CH}_3, -\text{CH}_2\text{CH}_3, -\text{R}$	alkyl	
	$-\text{F}, -\text{Cl}, -\text{Br}, -\text{I}$	halo	
Meta-Directing	$-\text{C}(=\text{O})-\text{R}$	acyl, carboxy	Deactivating
	$-\text{C}(=\text{O})\text{NH}_2$	carboxamido, carboalkoxy	
	$-\text{SO}_3\text{H}$	sulfonic acid	
	$-\text{C}\equiv\text{N}$	cyano	
	$-\text{NO}_2$	nitro	

- Substituents that **release electrons** to the ring will **activate the ring** toward electrophilic substitution.
- Substituents that **withdraw electrons** from the ring will **deactivate the ring** toward electrophilic substitution.

## SIDE-CHAIN REACTIONS OF BENZENE-DERIVATIVES


### 1. Halogenation of an Alkyl Side Chain



## SIDE-CHAIN REACTIONS OF BENZENE-DERIVATIVES

### 2. Oxidation of an Alkyl Side Chain

- Conversion into a carboxyl group,  $\text{-COOH}$ , by treatment with hot potassium permanganate.
- Regardless the length of the alkyl chain, the product is always the same.



23  
Dr Mohamed El-Newehy

## GENERAL QUESTIONS

**PROBLEM 4.7** Draw the structure of

- |                                |                        |                          |                    |
|--------------------------------|------------------------|--------------------------|--------------------|
| a. o-dinitrobenzene            | b. m-nitrotoluene      | c. p-bromophenol         | d. 4-chloroaniline |
| e. p-vinylbenzaldehyde         | f. 1,4-dichlorobenzene | g. 1,3,5-trimethylphenol |                    |
| h. 4-ethyl-2,6-difluorotoluene |                        |                          |                    |

### EXAMPLE 4.2

Write out the steps in the mechanism for the nitration of benzene.

**PROBLEM 4.12** Write out the steps in the mechanism for the sulfonation of benzene.

24  
Dr Mohamed El-Newehy