

Fundamentals of Organic Chemistry

CHEM 109

For Students of Health Colleges

Credit hrs.: (2+1)

King Saud University

College of Science, Chemistry Department

CHEM 109

CHAPTER 4. ALCOHOLS, PHENOLS AND ETHERS

Alcohols, Phenols and Ethers

2

- **Alcohols, phenols and ethers** may be viewed as organic *derivatives of water*.
- **Alcohols and phenols** have a common functional group, *the hydroxyl group, -OH*.

| | | | | | |
|-------|---------|--------|-------|--------|---------|
| H-O-H | R-O-H | Ar-O-H | R-O-R | R-O-Ar | Ar-O-Ar |
| Water | Alcohol | Phenol | Ether | | |




- **Alcohols** are compounds whose molecules have a hydroxyl group attached to a *saturated* carbon atom.
- **Phenols** are compounds that have a hydroxyl group attached directly to a *benzene ring*.
- **Ethers** are compounds whose molecules have an oxygen atom bonded to **two** carbon atom.

Classification of Alcohols and Ethers

3

- Alcohols are classified as **primary (1°)**, **secondary (2°)**, or **tertiary (3°)**, depending on whether one, two, or three organic groups are connected to the hydroxyl-bearing carbon atom.




- Methyl alcohol**, which is not strictly covered by this classification, is usually grouped with the primary alcohols.
- Ethers are classified as
 - Symmetrical ethers**;
When the organic groups attached to the oxygen are **identical**.
 - Unsymmetrical ethers (mixed ethers)**;
When the organic groups attached to the oxygen are **different**.

Nomenclature of Alcohols

4

- The **common names** for the simplest alcohols consist of alkyl group attached to the hydroxyl function followed by the word alcohol: **Alkyl alcohol**.
- In the **IUPAC system**, alcohols are named according to the following rules.
 - Select the **longest continuous carbon chain that contains the -OH group**.
Drop the **-e** ending of the parent alkane and replace it by the suffix **-ol**: **Alkanol**
 - When **isomers are possible**, the chain is numbered so as to give the functional group (-OH) the **lowest possible number**.




Nomenclature of Alcohols


5

3. When alkyl side chains or other groups are present; they are named alphabetically and their positions are indicated by a number.

The position of the functional group (-OH) is always given the **lowest possible number** at the end of the name.



For cyclic alcohols, numbering always starts from the carbon bearing the -OH group.




Nomenclature of Alcohols

6

4. With Unsaturated Alcohols; If a molecule contains both an -OH group and a C=C or C-C triple bond, the -OH group takes preference before the double or triple bonds in getting the lower number.

The name should include (if possible) both the hydroxyl and the unsaturated groups, **even if this does not make the longest chain the parent hydrocarbon.**




Nomenclature of Alcohols

7

Alcohols with More Than One Hydroxyl Group

- Compounds with two adjacent alcohol groups are called **glycols**.
The most important example is ethylene glycol.
- Compounds with more than two hydroxyl groups are also known, and several, such as glycerol and sorbitol, are important commercial chemicals.




- Ethylene glycol** is used as the “permanent” antifreeze in automobile radiators and as a raw material in the manufacture of Dacron.
- Ethylene glycol** is completely miscible with water.
- Glycerol** is a syrupy, colorless, water-soluble, high-boiling liquid with a distinctly sweet taste. Its soothing qualities make it useful in shaving and toilet soaps and in cough drops and syrups.


Nomenclature of Phenols

8

- Phenols** are usually named as derivatives of the parent compounds.



- The hydroxyl group is named as a substituent when it occurs in the same molecule with carboxylic acid, aldehyde, or ketone functionalities, which have **priority in naming**.





Nomenclature of Ethers

9


Common Names

Ethers are usually named by giving the name of each alkyl or aryl group, in alphabetical order, followed by the word *ether*.

| | | | |
|--------------|---|-------------------------------|---|
| Methyl ether | $\text{CH}_3\text{—O—CH}_3$ | Ethyl methyl ether | $\text{CH}_3\text{—O—CH}_2\text{CH}_3$ |
| Ethyl ether | $\text{CH}_3\text{CH}_2\text{—O—CH}_2\text{CH}_3$ | Ethyl- <i>n</i> -propyl ether | $\text{CH}_3\text{CH}_2\text{—O—CH}_2\text{CH}_2\text{CH}_3$ |
| Vinyl ether | $\text{CH}_2=\text{CH—O—CH}=\text{CH}_2$ | <i>t</i> -Butyl methyl ether | $(\text{CH}_3)_3\text{C—O—CH}_3$ |
| Phenyl ether |  | Methyl phenyl ether (anisole) |  |

IUPAC system

- For ethers with more complex structures, it may be necessary to name the **-OR** group as an **alkoxy** group.
- In the IUPAC system, the smaller alkoxy group is named as a substituent.



Physical Properties of Alcohols and Ethers

10


Physical State

- The simplest **alcohol**, methanol, is a liquid at room temperature. In contrast, alkanes from methane to butane are gases.
- Phenol** is a colorless, crystalline, and low-melting solid and other phenols also are solids, .
- Ethers** are colorless compounds with characteristic, relatively pleasant odors.

Boiling Points

- Ethers** have lower boiling points (bps) than **alcohols** with an equal number of carbon atoms.
- Ether** has nearly the same b.p. as the corresponding hydrocarbon in which a $-\text{CH}_2-$ group replaces the ether's oxygen.

Because of their structures (no O-H bonds), ether molecules cannot form hydrogen bonds with one another.




| Compound | Formula | bp | mol wt | Water solubility (g/100 mL, 20°C) |
|---------------|---|-------|--------|-----------------------------------|
| 1-butanol | $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$ | 118°C | 74 | 7.9 |
| diethyl ether | $\text{CH}_3\text{CH}_2\text{—O—CH}_2\text{CH}_3$ | 35°C | 74 | 7.5 |
| pentane | $\text{CH}_3\text{CH}_2\text{—CH}_2\text{—CH}_2\text{CH}_3$ | 36°C | 72 | 0.03 |

Physical Properties of Alcohols and Ethers

11

Boiling Points

- Series of normal alcohols; The boiling points increase with increasing molecular weights.
- A comparison of boiling points among isomeric alcohols; The boiling points decrease as the number of alkyl branches from the carbinol group increases.



- Phenol and most other phenols have high boiling points.


Physical Properties of Alcohols and Ethers

12

Solubility

- The lower alcohols are completely miscible with water.
- As the number of carbons in the alcohol increases, the solubility in water decreases.
- Low-molecular-weight ethers, such as dimethyl ether, are quite soluble in water.

Ether molecules can form hydrogen bonds to water.



| Structure | Name | Mol.wt. | Bp (°C) | Solubility in H ₂ O At 20 °C |
|--|--------------------|---------|---------|---|
| CH ₃ CH ₂ CH ₃ | propane | 44 | -42 | insoluble |
| CH ₃ OCH ₃ | methyl ether | 46 | -24 | soluble |
| CH ₃ CH ₂ OH | ethanol | 46 | 78 | soluble |
| CH ₃ CH ₂ CH ₂ CH ₃ | n-butane | 58 | -0.5 | insoluble |
| CH ₃ CH ₂ OCH ₂ CH ₃ | ethyl methyl ether | 60 | 8 | soluble |
| CH ₃ CH ₂ CH ₂ OH | 1-propanol | 60 | 97 | soluble |
| CH ₃ (CH ₂) ₄ CH ₃ | n-pentane | 72 | 35 | insoluble |
| CH ₃ CH ₂ OCH ₂ CH ₂ CH ₃ | ethyl ether | 74 | 36 | 7.5 g/100 g |
| CH ₃ (CH ₂) ₃ CH ₂ OH | 1-butanol | 74 | 118 | 7.9 g/100 g |
| CH ₃ (CH ₂) ₆ CH ₃ | n-heptane | 100 | 98 | insoluble |
| CH ₃ (CH ₂) ₂ O(CH ₂) ₂ CH ₃ | n-propyl ether | 102 | 91 | 0.2 g/100 g |
| CH ₃ (CH ₂) ₅ CH ₂ OH | 1-hexanol | 102 | 157 | 0.6 g/100 g |

- Phenol and most other phenols are slightly soluble in water .

Hydrogen Bonding in Alcohols and Ethers


13

- The **boiling points** (bp's) of alcohols are much higher than those of ethers or hydrocarbons with similar molecular weights.

| | CH ₃ CH ₂ OH | CH ₃ OCH ₃ | CH ₃ CH ₂ CH ₃ |
|--------|------------------------------------|----------------------------------|---|
| mol wt | 46 | 46 | 44 |
| bp | +78.5°C | -24°C | -42°C |

Why? Because alcohols form hydrogen bonds with one another.

The O-H bond is polarized by the high electronegativity of the oxygen atom and places a partial positive charge on the hydrogen atom and a partial negative charge on the oxygen atom.




- Consequently, **alcohols** have relatively high boiling points because they must supply enough heat to break the hydrogen bonds before each molecule.
- Hydrogen bonds are weaker than ordinary covalent bonds.

Hydrogen Bonding in Alcohols and Ethers

14

- The lower molecular-weight alcohols and ethers can form H-bond with water molecules.
- This accounts for the complete miscibility of the lower alcohols and ethers with water.



- However, as the organic chain lengthens and the alcohol becomes relatively more hydrocarbon like, its water solubility decreases.

Table 7.1 Boiling Point and Water Solubility of Some Alcohols



| Name | Formula | bp, °C | Solubility in H ₂ O g/100 g at 20°C |
|------------|--|--------|--|
| methanol | CH ₃ OH | 65 | completely miscible |
| ethanol | CH ₃ CH ₂ OH | 78.5 | completely miscible |
| 1-propanol | CH ₃ CH ₂ CH ₂ OH | 97 | completely miscible |
| 1-butanol | CH ₃ CH ₂ CH ₂ CH ₂ OH | 117.7 | 7.9 |
| 1-pentanol | CH ₃ CH ₂ CH ₂ CH ₂ CH ₂ OH | 137.9 | 2.7 |
| 1-hexanol | CH ₃ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ OH | 155.8 | 0.59 |

The Acidity of Alcohols and Phenols

15

- Like water, alcohols and phenols are weak acids.


The hydroxyl group can act as a proton donor, and dissociation occurs in a manner similar to that for water



The Acidity of Alcohols and Phenols

16

- Phenols are stronger acids than alcohols mainly because the corresponding phenoxide ions are stabilized by resonance.




The negative charge of an *alkoxide ion* is concentrated on the oxygen atom, but the negative charge on a *phenoxide ion* can be delocalized to the ortho and para ring positions through resonance.

Because *phenoxide ions* are stabilized in this way, the equilibrium for their formation is more favorable than that for *alkoxide ions*

The Acidity of Alcohols and Phenols

17


- All **electron-withdrawing groups increase acidity** by stabilizing the conjugate base. **Electron-donating groups decrease acidity** because they destabilize the conjugate base.



The Acidity of Alcohols and Phenols

18


- Alkoxides**, the conjugate bases of alcohols, can be prepared by the reaction of an alcohol with sodium or potassium metal.



- Treatment of alcohols with sodium hydroxide **does not convert** them to their alkoxides.

This is because alkoxides are stronger bases than hydroxide ion, so the reaction goes in the reverse direction. Since alcohols are weaker acids than water, it is not possible to form the salt of an alcohol in aqueous alkaline solutions.

- Treatment of phenols with sodium hydroxide **converts** them to phenoxide ions.




Preparation of Alcohols


19

1. Hydration of Alkenes

a. Addition of water to a double bond in the presence of an *acid catalyst, H⁺*.




b. The addition follows *Markovnikov's rule*.



c. It is *not possible to prepare primary alcohols* except Ethanol.

2. Oxidation of Cycloalkenes

Alkenes react with alkaline potassium permanganate to form glycols.



Preparation of Alcohols



20

○ Nucleophilic Substitution of Alkyl Halide



○ Reduction of Ketones, and Aldehydes

Aldehydes and ketones are easily reduced to primary and secondary alcohols, respectively.






Preparation of Alcohols

21

○ Addition of Grignard's Reagent to Aldehydes and Ketones

General equations




Preparation of Phenols

22

○ The Alkali Fusion of Sulfonates

The alkali fusion of sulfonates involves the following steps;

1. **Sulfonation** of an aromatic ring.
2. **Melting (fusion)** of the aromatic sulfonic acid with sodium hydroxide to give a **phenoxide salt**.
3. **Acidification** of the phenoxide with HCl to produce the **phenol**.




Preparation of Ethers

23


1) Dehydration of Alcohols

- It takes place in the presence of acid catalysts (H_2SO_4 , H_3PO_4) (intermolecular reaction)




- Example;**

The most important commercial ether is diethyl ether. It is prepared from ethanol and sulfuric acid.



- When ethyl alcohol is dehydrated by **sulfuric acid at 180°C** , the dominant product is **ethylene**.



Preparation of Ethers

24

2) Williamson Synthesis

- This method has two steps;
 - An alcohol is converted to its alkoxide by treatment with a reactive metal (sodium or potassium).



- Displacement is carried out between the alkoxide and an alkyl halide.



- To obtain the best yields of mixed dialkyl ethers, we select a 1° rather than a 2° or 3° alkyl halide and react it with a sodium alkoxide
- To prepare an alkyl aryl ether, we must be careful not to pick a combination in which one of the reagents has a halogen directly attached to an aromatic ring.

Preparation of Ethers

25

○ Example 1; Preparation of *t*-butyl methyl ether, $(\text{CH}_3)_3\text{C-O-CH}_3$.


➤ In theory, this could be done by either of two reactions.

1. You could react sodium methoxide, $\text{CH}_3\text{O}^-\text{Na}^+$, with *t*-butyl chloride, $(\text{CH}_3)_3\text{C-Cl}$.

This combination leads to dehydrohalogenation to an alkene, an elimination reaction.

2. You could react sodium *t*-butoxide, $(\text{CH}_3)_3\text{C-O}^-\text{Na}^+$, with methyl chloride, CH_3Cl .

This route gives the desired ether by substitution.




Preparation of Ethers

26

Example 2; Assume you need to synthesize methyl phenyl ether (anisole), $\text{CH}_3\text{-O-C}_6\text{H}_5$, by the Williamson method.

➤ In theory, you could obtain anisole in either of two ways.



Reactions of Alcohols

29

A) Those that involve the rupture of the carbon-oxygen bond (C-OH).

1) Nucleophilic Substitution Reaction; The Reaction of Alcohols with Hydrogen Halides:


Alkyl Halides

Alcohols react with hydrogen halides (HCl, HBr and HI) to give alkyl halides.



2) Dehydration of Alcohols: Formation of Alkenes

Alcohols can be dehydrated by heating them with strong acid.




Reactions of Alcohols

30



B) Oxidation Reactions

Alcohols with at least one hydrogen attached to the hydroxyl-bearing carbon can be oxidized to carbonyl compounds.

- **Primary alcohols** give **aldehydes**, which may be further oxidized to **carboxylic acids**.



- **Primary alcohols**, oxidation can be stopped at aldehyde stage by special reagents, such as "pyridinium chlorochromate (PCC)".




Reactions of Alcohols


31

B) Oxidation Reactions

- Primary alcohols yield aldehydes when treated with *mild oxidizing agents* such as hot metallic copper or CrO_3 in pyridine.



- Primary alcohols; when treated with *stronger oxidizing agents*, such as chromic acid, $\text{H}_2\text{Cr}_2\text{O}_7$, or neutral potassium permanganate, KMnO_4 , the intermediate aldehydes formed initially are oxidized further to carboxylic acids.




Reactions of Alcohols

32

B) Oxidation Reactions

- Secondary alcohols, when treated with any of the oxidizing agents mentioned previously, yield ketones.




- Tertiary alcohols, having no hydrogen atom on hydroxyl-bearing carbon, do not undergo oxidation.

Reactions of Phenols

33

- Halogenation takes place *without catalyst*.




- The products depend on the solvent used.
 - In *aprotic solvents* (solvents that do not release protons) (CCl_4 , CS_2)-bromination gives a mixture of *o*- and *p*-bromophenol.
 - In *protic solvents* (solvents that can release protons) (H_2O)-halogenation gives a trisubstituted phenol is produced.

Reactions of Ethers

34

- Cleavage of Ethers by Hot Concentrated Acids

- When **ethers** are heated in concentrated acid solutions, the ether linkage is broken.



- The acids most often used in this reaction are HI, HBr, and HCl.
- If an excess of acid is present, the alcohol initially produced is converted into an alkyl halide by the reaction.



For example,

