

King Saud University  
 College of Sciences  
 Department of Mathematics

Final Examination Math-244 Semester I, (1441 H) Max. Time: 3h

**Note: Calculators are not allowed.** **2 pages**

**Question 1** [2+2+3]

a) Find the value of  $x$  so that  $\det A = 6$ , where  $A = \begin{pmatrix} 1 & 2 & 0 \\ -1 & 3x-6 & 1 \\ 1 & 0 & 2 \end{pmatrix}$ .

b) Let  $X_1 = \begin{pmatrix} 2 \\ 1 \end{pmatrix}$  and  $X_2 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$ . Find the matrix  $A$  of order 2 such that  $AX_1 = \begin{pmatrix} 3 \\ 1 \end{pmatrix}$  and  $AX_2 = \begin{pmatrix} 2 \\ 2 \end{pmatrix}$ .

c) Let  $m \in \mathbb{R}$  and consider the linear system:

$$\begin{cases} mx + y + 2z &= 3 \\ mx + my + 3z &= 5 \\ 3mx + (m+2)y + (m+6)z &= 2m+9. \end{cases}$$

Find the value(s) of  $m$  so that the system has infinitely many solutions.

**Question 2** [2+3+(2+2)]

a) Show that the matrix  $A = \begin{pmatrix} 1 & -1 & 2 \\ 3 & 4 & -5 \\ 4 & 3 & -3 \end{pmatrix}$  cannot be a transition matrix between two bases of any 3-dimensional vector space.

b) Find a basis for the solution space of the following homogeneous linear system:

$$\begin{cases} x - 3y + z &= 0 \\ 2x - 6y + 2z &= 0 \\ 3x - 9y + 3z &= 0. \end{cases}$$

c) Consider the bases  $B = \{u_1 = (1, 1, 0), u_2 = (1, 1, 1), u_3 = (1, 0, 1)\}$  and  $C = \{v_1 = (1, -1, 0), v_2 = (1, -1, 1), v_3 = (-1, 0, 1)\}$  for  $\mathbb{R}^3$ .

Find the matrices  ${}_C P_B$  and  ${}_B P_C$ . [ ${}_C P_B$  is the transition matrix from  $B$  to  $C$ .]

**Question 3** [3+(2+2)]

a) Let  $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$  be a linear transformation such that:  
 $T(x, y, z) = (ax + 2by + z, ax - by + 2z, 2ax + by + 3z)$ .  
Find the values of  $a$  and  $b$  so that  $(-5, 1, 3)$  is in the kernel of  $T$ .

b) Let  $T: \mathbb{R}^5 \rightarrow \mathbb{R}^3$  be the linear transformation defined by the formula:

$$T(x_1, x_2, x_3, x_4, x_5) = (x_1 + x_2, x_2 + x_3 + x_4, x_4 + x_5)$$

(i) Find the matrix  $A$  of  $T$  relative to the standard bases of  $\mathbb{R}^5$  and  $\mathbb{R}^3$ .  
(ii) Find the rank and the nullity of the matrix  $A$ .

**Question 4** [(1+1)+2+2]

Suppose  $T: V \rightarrow W$  is a linear transformation,  $B = \{v_1, v_2, v_3\}$  is a basis for  $V$  and  $C = \{w_1, w_2, w_3, w_4\}$  is a basis for  $W$ .

If  $T(v_1) + T(v_2) = w_1 - 3w_2 + w_4$ ,  $T(v_2) = 2w_1 + w_2 - w_3 + 3w_4$  and  $T(v_2) + T(v_3) = 2w_2 + 3w_3 - w_4$ .

Find:

(i)  $T(v_1)$  and  $T(v_3)$ .  
(ii) The matrix  $[T]_B^C$ .  
(iii)  $[T(v)]_C$ , where  $[v]_B = \begin{pmatrix} -3 \\ 4 \\ 1 \end{pmatrix}$ .

**Question 5** [(2+2)+2+2+3]

a) Let  $F$  be the subspace of the Euclidean space  $\mathbb{R}^4$  spanned by the following vectors:  $v_1 = (1, 1, -1, 0)$ ,  $v_2 = (0, 1, 1, 1)$ ,  $v_3 = (1, 0, 1, 1)$ ,  $v_4 = (0, -1, 2, 1)$ .

(i) Show that  $\{v_1, v_2, v_3\}$  is a basis for  $F$ .  
(ii) Use Gram-Schmidt algorithm to convert the basis  $\{v_1, v_2, v_3\}$  into an orthonormal basis for  $F$ .

b) Find  $a$  such that the vector  $\begin{pmatrix} 1 \\ -1 \end{pmatrix}$  is an eigenvector of the matrix  $A = \begin{pmatrix} 1 & -1 \\ 2 & a \end{pmatrix}$ .

c) If 3 is an eigenvalue of the matrix  $A = \begin{pmatrix} 2 & -1 \\ 1 & b \end{pmatrix}$ , find  $b$ .

d) Find  $A^{17}$ , where  $A = \begin{pmatrix} 3 & 5 & -6 \\ 0 & 0 & -5 \\ 0 & 0 & -3 \end{pmatrix}$ .