

421 SDS Sample Exam Questions

1. The width of maxillary central incisor may be estimated by:
 - A. Dividing the maximum bi zygomatic width by 16
 - B. adding 6 mm to the distance between the lateral extremities of the ala of the nose
 - C. Dividing the horizontal circumference of the cranium at a plane passing through the glabella and the maximum occipital point by 10
 - D. Adding 5 mm to the distance between the buccolingual centers of the hamular notches

2. To determine the profile of the face, the following points should be checked:
 - A. The forehead, the tip of the nose and point of the chin
 - B. the forehead, the base of the nose and the point of the chin
 - C. The forehead and the point of the chin
 - D. The forehead, the most prominent point of the nose and the tip of the chin

3. To provide femininity to your teeth selection, the following points should be considered:
 - A. Square teeth should be selected
 - B. curved labial surfaces should be selected instead of flat surfaces
 - C. Larger size teeth should be selected
 - D. Less number of teeth should be selected

4. One of the disadvantages of porcelain teeth is:
 - A. Porcelain teeth bond chemically with acrylic denture base
 - B. Porcelain teeth produce noise during function
 - C. Porcelain teeth are easily abraded by opposing teeth
 - D. Colour of porcelain teeth is not stable

5. When choosing the materials of posterior teeth replacement for RPD, the most accurate combination is:
 - A. Acrylic teeth should be selected to oppose upper porcelain teeth
 - B. porcelain teeth should be selected to oppose upper gold crowns
 - C. Gold occlusal coverage acrylic teeth should oppose upper porcelain teeth
 - D. Porcelain teeth should oppose upper porcelain fused to metal fixed partial denture

6. The length of edentulous space in distal extension cases opposed by complete denture is measured from:

- A. The distal surface of the canine to two thirds the retro molar pad
- B. the distal surface of the canine to the mesial end of retro molar pad
- C. The buccal frenum to the incline of the mandibular ramus
- D. The distal surface of the canine to the incline of the mandibular ramus

7. By determining the shape of the face, which of the following can be determined:

- A. The form of posterior teeth
- B. the form of anterior teeth
- C. The size of anterior teeth
- D. The size of posterior teeth

8. Regarding the anatomic teeth, which of the following is correct:

- A. They are indicated in cases of reduced residual ridges and poor neuromuscular control
- B. they have flat occlusal surfaces
- C. They transmit less destructive forces to the underlying ridges
- D. They provide good cutting abilities for mastication

9. During posterior teeth selection for complete denture cases, the following points should be considered:

- A. The size of teeth should be larger than natural teeth to provide better support to the cheek
- B. the length of the teeth is determined by the position of the corner of the mouth
- C. The length of teeth is determined by the available inter arch space
- D. Non-anatomic teeth is used with balanced occlusion scheme

10. When selecting lower anterior teeth for class III anterior relationship:

- A. Use the mandibular anterior teeth that corresponds to the upper anterior teeth in size, shape and shade
- B. use smaller teeth than the corresponding ones to overcome the reduced anterior space
- C. Use larger teeth than the corresponding ones to overcome the increased anterior space
- D. Use longer anterior teeth

Answers:

1.a

2.b

3.b

4.b

5.d

6. d

7.b

8.d

9.c

10. c