

Fundamentals of Organic Chemistry

CHEM 109

For Students of Health Colleges

Credit hrs.: (2+1)

King Saud University

College of Science, Chemistry Department

CHEM 109

CHAPTER 5. ALDEHYDES & KETONES

Learning Objectives







2

At the end of this chapter, students will be able to:

- Know the structural differences between aldehydes and ketones
- Know how to draw aldehydes and ketones
- know the common and IUPAC nomenclature of aldehydes and ketones
- Know the physical properties of aldehydes and ketones
- Know how to synthesize an aldehyde or a ketone.
- Know the different nucleophilic addition reactions at the carbonyl carbon.


Common Classes of Carbonyl Compounds

3

Class	General Formula	Class	General Formula
Ketones		Aldehydes	
Carboxylic acids		Acid Chlorides	
Esters		Amides	


The Carbonyl Group

4



- The carbon–oxygen double bond consists of a sigma bond and a pi bond.
- The carbon atom is sp^2 -hybridized.
- The three atoms attached to the carbonyl carbon lie in a plane with bond angles of 120° .
- The pi bond is formed by overlap of a p orbital on carbon with an oxygen p orbital.
- There are also two unshared electron pairs on the oxygen atom.
- The C=O bond distance is 1.24Å, shorter than the C–O distance in alcohols and ethers (1.43Å).


	<i>length</i>	<i>energy</i>
ketone C=O bond	1.23 Å	178 kcal/mol (745 kJ/mol)
alkene C=C bond	1.34 Å	146 kcal/mol (611 kJ/mol)




The Carbonyl Group

5

- Oxygen is much more electronegative than carbon. Therefore, the electrons in the C=O bond are attracted to the oxygen, producing a highly **polarized bond**.




- As a consequence of this polarization, *most carbonyl reactions involve nucleophilic attack at the carbonyl carbon*, often accompanied by addition of a proton to the oxygen (electron rich).



Structure of Aldehydes and Ketones

6

- Aldehydes and ketones** are characterized by the presence of the carbonyl group.
- Aldehydes** have at least one hydrogen atom attached to the carbonyl carbon atom. *The remaining group may be another hydrogen atom or any aliphatic or aromatic organic group.* The **-CH=O** group characteristic of aldehydes is often called a formyl group.
- In **ketones**, the carbonyl carbon atom is connected to two other carbon atoms.



IUPAC System


Nomenclature of Aldehydes

King Saud University


7

- **Aliphatic aldehydes** are named by dropping the suffix **-e** from the name of the hydrocarbon that has the same carbon skeleton as the aldehyde and replacing it with the suffix **-al**.

Alkane -e + al = Alkanal



- **Substituted aldehydes**, we number the chain starting with the aldehyde carbon.
 - **-CH=O group** is assigned the number **1 position**.
 - Aldehyde group has priority over a double bond or hydroxyl group.




IUPAC System

Nomenclature of Aldehydes


King Saud University

8

- **Cyclic aldehydes**, the suffix **-carbaldehyde** is used.



- **Aromatic aldehydes** are usually designated as derivatives of the simplest aromatic aldehyde, **benzaldehyde**.





Common Names

Nomenclature of Aldehydes



9

- Common names of ketones are formed by adding the word *ketone* to the names of the alkyl or aryl groups attached to the carbonyl carbon. **Alkyl ketone.**
- In still other cases, traditional names are used.






IUPAC Names

Nomenclature of Ketones



10

- In the IUPAC system, **the ending for ketones is -one.**
- The chain is numbered so that the **carbonyl carbon has the lowest possible number.**
- For **cyclic ketones**, numbering always starts from the C=O group.
- The prefix "**oxo**" is used when the ketone is not the principal functional group.




NOTES

Nomenclature of Aldehydes and Ketones



11

- In common names carbon atoms near the carbonyl group are often designated by Greek letters.
- The atom adjacent to the function is *alpha* (α), the next removed is *beta* (β) and so on. Since ketones have two sets of neighboring atoms, one set is labeled α , β etc., and the other α' , β' etc.



- The functional group priority order in nomenclature system is as following:
Acid and derivatives > aldehyde > ketone > alcohol > amine > alkene > alkyne > ether


Physical Properties of Aldehydes and Ketones




12

Boiling Points

- Carbonyl compounds boil at higher temperatures than hydrocarbons, but at lower temperatures than alcohols of comparable molecular weight.



- This is due to the intermolecular forces of attraction, called dipole-dipole interactions, which is stronger than van der Waals attractions but not as strong as hydrogen bonds.




Dipole-dipole attractions among carbonyl compounds

Physical Properties of Aldehydes and Ketones

13

Solubility



- Carbonyl compounds as aldehydes and ketones have a C=O bond, but no O-H bond, cannot form hydrogen bonds with themselves.
- The polarity of the carbonyl group also affects the solubility properties of aldehydes and ketones.
- Carbonyl compounds with low molecular weights are soluble in water as they can form **hydrogen bonds** with O-H or N-H compounds.



Preparation of Aldehydes and Ketones


14

1) Oxidation of Primary and Secondary Alcohols





2) Hydration of Alkynes

Hydration of terminal alkynes EXCEPT acetylene yields ketones (catalyzed by acid and mercuric).



3) Ozonolysis of Alkenes




B) Oxidation of Carbonyl Compounds



Reactions of Aldehydes and Ketones

17

- Oxidation of aldehydes gives a carboxylic acid with the same number of carbon atoms.
- Because the reaction occurs easily, many oxidizing agents, such as KMnO_4 , CrO_3 , Ag_2O and peracids will work.



- Example:**




C) Nucleophilic Addition Reactions

Reactions of Aldehydes and Ketones

18

- Nucleophiles attack the carbon atom of a carbon-oxygen double bond because that carbon has a partial positive charge.
- The overall reaction involves addition of a nucleophile and a proton across the pi bond of the carbonyl group (when carried out in alcohol or water).




C) Nucleophilic Addition Reactions

Reactions of Aldehydes and Ketones

19

1) Addition of Grignard Reagents: Formation of Alcohols

- Grignard reagents act as carbon nucleophiles toward carbonyl compounds.
- The reaction of a Grignard reagent with a carbonyl compound provides a useful route to alcohols.



- The type of carbonyl compound chosen determines the class of alcohol produced.


C) Nucleophilic Addition Reactions

Reactions of Aldehydes and Ketones


20

1) Addition of Grignard Reagents: Formation of Alcohols


- Formaldehyde gives primary alcohols.



- Other aldehydes give secondary alcohols



- Ketones give tertiary alcohols.




C) Nucleophilic Addition Reactions

Reactions of Aldehydes and Ketones



21

2) Addition of Hydrogen Cyanide: Formation of Cyanohydrins

- Hydrogen cyanide adds to the carbonyl group of aldehydes and ketones to form cyanohydrins, compounds with a hydroxyl and a cyano group attached to the same carbon.



Example




C) Nucleophilic Addition Reactions

Reactions of Aldehydes and Ketones


22

3) Addition of Alcohols: Formation of Hemiacetals and Acetals


- Alcohols add to the C=O bond, the OR group becoming attached to the carbon and the proton becoming attached to the oxygen.
- Aldehydes and ketones react with alcohols to form, first, [hemiacetals](#) and then, if excess alcohol is present, [acetals](#).



- Hemiacetals**; it contains both alcohol and ether functional groups on the same carbon atom.



- Acetals** have two ether functions at the same carbon atom.



Uses of Aldehydes and ketones



25

Aldehydes and ketones find application in different sectors such as pharmaceutical, food, fragrance, cosmetics because of their chemical properties.

- **Formaldehyde** is found in the gaseous form.
 - Formaldehyde with 40% solution in water forms formalin, helps in the preservation of biological specimens.
 - It is essential during industrial processes manufacture of polymeric products.
 - It acts as germicide, insecticide, and fungicides.
 - It helps in the testing of drugs. It is also used in photography.
 - It forms with ammonia a tricyclic condensation product called methenamine (hexamethylenetetramine).

Methenamine is used for the treatment of urinary tract infection: it decomposes at an acid pH to formaldehyde and ammonia, and the formed formaldehyde is bactericidal.

Uses of Aldehydes and ketones



26

The applications of aldehydes and ketones are used in different sectors such as pharmaceutical, food, fragrance, cosmetics industry.

- **Acetaldehyde**
 - It is used for the production of acetic acid and pyridine derivatives.
 - Its derivative, paraldehyde was once used as sedative and hypnotic drug.
- **Benzaldehyde**
 - It is an essential component for the production of perfumes, cosmetic products, and dyes.
 - is added to incorporate almond flavor into various food products.
 - It also acts as a bee repellent.
- **Anisaldehyde, vanillin or cinnamaldehyde** are mainly used in cosmetics, but also in drug formulation.
- **Glutaraldehyde** is used to disinfect medical and dental equipment.

Uses of Aldehydes and ketones

27

Therapeutic applications of aldehydes and ketones

Compound	Name in the Pharmacopoeia	Application
Aldehydes and analogues		
Formaldehyde	Formaldehydi solution (35 per centum)	antiseptic
Paraldehyde	Paraldehydum	sedative and hypnotic
Chloral hydrate	Chlorali hydras	sedative and hypnotic
Methenamine	Methenaminum	antiseptic
Vanillin	Vanillinum	flavouring additive

Uses of Aldehydes and ketones

28

Ketone behaves as an excellent solvent for certain types of plastics and synthetic fibers.

- **Acetone**
 - It act as a paint thinner and a nail paint remover.
 - It is also used for medicinal purposes such as chemical peeling procedure as well as acne treatments.
- **Butanone**
 - It is one of the common solvents.
 - It is used in textile production, varnishes production, paint remover production, paraffin wax production, plastic production, etc.
- **Cyclohexanone** is an important component in nylon production.
- **Acetophenone** is responsible for fragrances such as cherry, jasmine, honeysuckle, almond, strawberry, etc.