

**CHEM 108**

# **FUNDAMENTALS OF ORGANIC CHEMISTRY**

FOR B.Sc. PROGRAMS OF SCIENTIFIC COLLEGES

PRE-REQUISITES COURSE; CHEM 101

CREDIT HOURS; 4 (3+1)

Chemistry Department, College of Science, King Saud University

1  
Dr Mohamed El-Newehy

## **CHAPTER 5**

# **ALCOHOLS, PHENOLS AND ETHERS**

2  
Dr Mohamed El-Newehy

## Alcohols, Phenols and Ethers

- Alcohols, ethers and phenols have a common functional group, the hydroxyl group, -OH.


H-O-H Water	R-OH Alcohol	R-O-R Ethers	Ph-O-H Phenol
----------------	-----------------	-----------------	------------------

- Alcohols** are compounds whose molecules have a hydroxyl group attached to a *saturated* carbon atom.
- Phenols** are compounds that have a hydroxyl group attached directly to a *benzene ring*.
- Ethers** are compounds whose molecules have an oxygen atom bonded to **two** carbon atom.

Dr Mohamed El-Newehy

## Alcohols and Phenols


- Alcohols and phenols** may be viewed as organic *derivatives of water*.



- Alcohols** have the general formula R-OH, and structurally similar to water, but with one of the hydrogens replaced by an alkyl group.
- Phenols** have a hydroxyl group attached directly to an aromatic ring.


Dr Mohamed El-Newehy

## Alcohols



This is the functional group of an alcohol


- Alcohols can be viewed in two ways structurally:
  - as **hydroxyl derivatives** of alkanes
  - as **alkyl derivatives** of water.



Dr Mohamed El-Newehy

## Classification of Alcohols

- Alcohols** are classified as **primary (1°)**, **secondary (2°)**, or **tertiary (3°)**, depending on whether one, two, or three organic groups are connected to the hydroxyl-bearing carbon atom.




- Methyl alcohol**, which is not strictly covered by this classification, is usually grouped with the primary alcohols.

Dr Mohamed El-Newehy


## Nomenclature of Alcohols

- The **common names** for the simplest alcohols consist of alkyl group attached to the hydroxyl function followed by the word alcohol: *Alkyl alcohol*.
- In the **IUPAC system**, alcohols are named according to the following rules.
  - Select the **longest continuous carbon chain that contains the -OH group**.  
Drop the **-e** ending of the parent alkane and replace it by the suffix **-ol**: *Alkanol*
  - When **isomers are possible**, the chain is numbered so as to give the functional group (-OH) the **lowest possible number**.




Dr Mohamed El-Newehy

## Nomenclature of Alcohols



- When **alkyl side chains or other groups are present**, they are named alphabetically and their positions are indicated by a number.



The position of the functional group (-OH) is always given the **lowest possible number** at the end of the name.



Dr Mohamed El-Newehy





## Nomenclature of Alcohols

For cyclic alcohols, numbering always starts from the carbon bearing the -OH group.



4. With Unsaturated Alcohols; If a molecule contains both an -OH group and a C=C or C-C triple bond, the -OH group takes preference before the double or triple bonds in getting the lower number.

The name should include (if possible) both the hydroxyl and the unsaturated groups, even if this does not make the longest chain the parent hydrocarbon.







Dr Mohamed El-Newehy

## Nomenclature of Alcohols

### Alcohols with More Than One Hydroxyl Group

- Compounds with two adjacent alcohol groups are called **glycols**.  
The most important example is ethylene glycol.
- Compounds with more than two hydroxyl groups are also known, and several, such as glycerol and sorbitol, are important commercial chemicals.




- Ethylene glycol** is used as the "permanent" antifreeze in automobile radiators and as a raw material in the manufacture of polyester fiber: Dacron.
- Ethylene glycol** is completely miscible with water.
- Glycerol** is a syrupy, colorless, water-soluble, high-boiling liquid with a distinctly sweet taste. Its soothing qualities make it useful in shaving and toilet soaps and in cough drops and syrups.


Dr Mohamed El-Newehy

## Nomenclature of Phenols

- Phenols are usually named as derivatives of the parent compounds.



- The hydroxyl group is named as a substituent when it occurs in the same molecule with carboxylic acid, aldehyde, or ketone functionalities, which have **priority in naming**.



Dr Mohamed El-Neelhy

## Physical Properties of Alcohols

### Physical State


- The simplest alcohol, methanol, is a liquid at room temperature. In contrast, alkanes from methane to butane are gases.

### Solubility

- The lower alcohols are completely miscible with water.
- As the number of carbons in the alcohol increases, the solubility in water decreases.

### Boiling Points

- Series of normal alcohols; The boiling points increase with increase in molecular weights.
- A comparison of boiling points among isomeric alcohols; The boiling points decrease as the number of alkyl branches from the carbinol group increases.



Dr Mohamed El-Neelhy


## Hydrogen Bonding in Alcohols

- The **boiling points** (bp's) of alcohols are much higher than those of ethers or hydrocarbons with similar molecular weights.

	<chem>CH3CH2OH</chem>	<chem>CH3OCH3</chem>	<chem>CH3CH2CH3</chem>
mol wt	46	46	44
bp	+78.5°C	-24°C	-42°C

**Why?** Because alcohols form hydrogen bonds with one another.

The O-H bond is polarized by the high electronegativity of the oxygen atom and places a partial positive charge on the hydrogen atom and a partial negative charge on the oxygen atom.



Two or more alcohol molecules thus become loosely bonded to one another through hydrogen bonds.

Dr Mohamed El-Newehy

## Hydrogen Bonding in Alcohols

- Consequently, alcohols have relatively high boiling points because they must supply enough heat to break the hydrogen bonds before each molecule.
- Hydrogen bonds are weaker than ordinary covalent bonds.
- Water, of course, is also a hydrogen-bonded liquid.
- The **lower molecular-weight alcohols** can readily replace water molecules in the hydrogen bonded network.
- This accounts for the complete **miscibility of the lower alcohols with water**.
- However, as the organic chain lengthens and the alcohol becomes relatively more hydrocarbon like, its water solubility decreases.

Table 7.1 Boiling Point and Water Solubility of Some Alcohols

Name	Formula	bp, °C	Solubility in H <sub>2</sub> O g/100 g at 20°C
methanol	<chem>CH3OH</chem>	65	completely miscible
ethanol	<chem>CH3CH2OH</chem>	78.5	completely miscible
1-propanol	<chem>CH3CH2CH2OH</chem>	97	completely miscible
1-butanol	<chem>CH3CH2CH2CH2OH</chem>	117.7	7.9
1-pentanol	<chem>CH3CH2CH2CH2CH2OH</chem>	137.9	2.7
1-hexanol	<chem>CH3CH2CH2CH2CH2CH2OH</chem>	155.8	0.59

Dr Mohamed El-Newehy

## Physical Properties of Phenols


- **Phenol** is a colorless, crystalline, low-melting solid, with a high boiling point, that is moderately soluble in water.
- Most other phenols also are solids, with slight solubility in water and high boiling points.
- The most significant physical property that distinguishes alcohols from phenols is the acidity of phenols.

Dr Mohamed El-Newehy

## The Acidity of Alcohols and Phenols

- Like water, alcohols and phenols are weak acids.

*The hydroxyl group can act as a proton donor, and dissociation occurs in a manner similar to that for water*




Dr Mohamed El-Newehy



## The Acidity of Alcohols and Phenols

- Phenols are stronger acids than alcohols mainly because the corresponding phenoxide ions are stabilized by resonance.




The negative charge of an **alkoxide ion** is concentrated on the oxygen atom, but the negative charge on a phenoxide ion can be delocalized to the ortho and para ring positions through resonance.

Because **phenoxide ions** are stabilized in this way, the equilibrium for their formation is more favorable than that for alkoxide ions

Dr Mohamed El-Newehy

## The Acidity of Alcohols and Phenols


- All **electron-withdrawing groups increase acidity** by stabilizing the conjugate base. **Electron-donating groups decrease acidity** because they destabilize the conjugate base.



Dr Mohamed El-Newehy

## The Acidity of Alcohols and Phenols

- Alkoxides, the conjugate bases of alcohols, can be prepared by the reaction of an alcohol with sodium or potassium metal.




- Treatment of alcohols with sodium hydroxide does not convert them to their alkoxides.

*This is because alkoxides are stronger bases than hydroxide ion, so the reaction goes in the reverse direction.*

*Since alcohols are weaker acids than water, it is not possible to form the salt of an alcohol in aqueous alkaline solutions.*

- Treatment of phenols with sodium hydroxide converts them to phenoxide ions.





Dr Mohamed El-Newehy

## Preparation of Alcohols

### 1. From Alkenes

#### A. Acid-Catalyzed Hydration of Alkenes

- Addition of water to a double bond in the presence of an *acid catalyst*,  $\text{H}^+$ .



*It is not possible to prepare primary alcohols except Ethanol.*

Dr Mohamed El-Newehy


## Preparation of Alcohols; From Alkenes

## 1. From Alkenes

### A. Acid-Catalyzed Hydration of Alkenes

**2. The addition follows Markovnikov's rule.**

**Mechanism:** Water adds to alkenes in the presence of an acid catalyst following **Markovnikov's rule**, the reaction is reversible.




Dr Mohamed El-Newehi

## Preparation of Alcohols; From Alkenes

## 1. From Alkenes

## B. Oxidation of Alkenes and Cycloalkenes


Alkenes react with alkaline potassium permanganate to form glycols (compounds with two adjacent hydroxyl groups).





Dr Mohamed El-Neweh

## Preparation of Alcohols

### 2. Nucleophilic Substitution of Alkyl Halide



### 3. Reduction of Ketones and Aldehydes using $\text{LiAlH}_4$ or $\text{NaBH}_4$





Dr Mohamed El-Newehy

## Preparation of Alcohols

### 3. Reduction of Ketones and Aldehydes using $\text{LiAlH}_4$ or $\text{NaBH}_4$

Aldehydes and ketones are easily reduced to primary and secondary alcohols, respectively.




Dr Mohamed El-Newehy

## Preparation of Alcohols

### 4. Addition of Grignard's Reagent to Aldehydes and Ketones<sub>4</sub>

*General equations*



Dr Mohamed El-Newehy

## Reactions of Alcohols and Phenols




- **Alcohols** undergo two kinds of reactions:
  - Those that involve the breaking of the oxygen-hydrogen bond (CO-H).
  - Those that involve the rupture of the carbon-oxygen bond (C-OH).
- **Phenols** do not participate in reactions where the C-OH bond is broken.

Dr Mohamed El-Newehy

## Reactions of Alcohols

A) Those that involve the breaking of the oxygen-hydrogen bond (CO-H).

### 1) Reactions of Alcohols and Phenols as Acids: Salt Formation.



Dr Mohamed El-Newehy

## Reactions of Alcohols

**B) Those that involve the rupture of the carbon-oxygen bond (C-OH).**

### 1) The Reaction of Alcohols with Hydrogen Halides: Alkyl Halides.




*Alcohols react with hydrogen halides (HCl, HBr and HI) to give alkyl halides.*

- Reaction of Primary Alcohols with Hydrogen Halides: The S<sub>N</sub>2 Mechanism:


*Methyl and primary carbocations are the least stable, and they are not likely to be intermediates in reaction mechanism:*



Reactivity of the alcohol:



Reactivity of the H-X : parallels the acidity of HX



**increasing reactivity**


Dr Mohamed El-Neweh

## Reactions of Alcohols

B) Those that involve the rupture of the carbon-oxygen bond (C-OH).

### 2) Dehydration of Alcohols: Formation of Alkenes

Alcohols can be dehydrated by heating them with strong acid.




Dr Mohamed El-Newehy

## Reactions of Alcohols


### C) Oxidation Reactions

➤ **Oxidation** is the removal of H from a compound and/or the addition of O to a compound.



An oxidizing agent is the chemical reagent that does the oxidation.

➤ **Reduction** is the addition of H to a compound and/or the removal of O from a compound.




A reducing agent is a substance that does the reduction.

Dr Mohamed El-Newehy

## Reactions of Alcohols

### C) Oxidation Reactions

- Alcohols with at least one hydrogen attached to the hydroxyl-bearing carbon can be oxidized to carbonyl compounds.
  - Primary alcohols give aldehydes, which may be further oxidized to carboxylic acids.
  - Secondary alcohols give ketones.
  - Tertiary alcohols, having no hydrogen atom on hydroxyl-bearing carbon, do not undergo oxidation.




Dr Mohamed El-Newehy

## Reactions of Alcohols

### C) Oxidation Reactions

- Primary alcohols, oxidation can be stopped at aldehyde stage by special reagents, such as “pyridinium chlorochromate (PCC)”.




Dr Mohamed El-Newehy



## Reactions of Alcohols

### C) Oxidation Reactions

- Primary alcohols yield aldehydes when treated with mild oxidizing agents such as hot metallic copper or  $\text{CrO}_3$  in pyridine.




Dr Mohamed El-Newehy

## Reactions of Alcohols

### C) Oxidation Reactions

- Primary alcohols yield aldehydes when treated with stronger oxidizing agents, such as chromic acid,  $\text{H}_2\text{Cr}_2\text{O}_7$ , or neutral potassium permanganate,  $\text{KMnO}_4$ , the intermediate aldehydes formed initially are oxidized further to carboxylic acids.




Dr Mohamed El-Newehy

## Reactions of Alcohols

### C) Oxidation Reactions

- Secondary alcohols, when treated with any of the oxidizing agents mentioned previously, yield ketones.




Dr Mohamed El-Newehy

## Preparation of Phenols

### The Alkali Fusion of Sulfonates

The alkali fusion of sulfonates involves the following steps;


- Sulfonation of an aromatic ring.
- Melting (fusion) of the aromatic sulfonic acid with sodium hydroxide to give a phenoxide salt.
- Acidification of the phenoxide with HCl to produce the phenol.



Dr Mohamed El-Newehy

## Reactions of Phenols

Halogenation takes place without catalyst.



➤ The products depend on the solvent used.

- In *aprotic solvents* (solvents that do not release protons) ( $\text{CCl}_4$ ,  $\text{CS}_2$ )-bromination gives a mixture of *o*- and *p*-bromophenol.
- In *protic solvents* (solvents that can release protons) ( $\text{H}_2\text{O}$ )-halogenation gives a trisubstituted phenol is produced.

Dr Mohamed El-Newehy


## ETHERS

38


Dr Mohamed El-Newehy

## Structure of Ethers

- All **ethers** are compounds in which two organic groups are connected to a single oxygen atom.
- The **general formula for an ether** is  $R-O-R'$ , where R and R' may be identical or different, and they may be alkyl or aryl groups



- The geometry of simple ethers is similar to that of water.



- The ether is classified as

- **Symmetrical ethers;**

When the organic groups attached to the oxygen are **identical**.

- **Unsymmetrical ethers (mixed ethers);**

When the organic groups attached to the oxygen are **different**.

Dr Mohamed El-Newehy

## Nomenclature of Ethers


### Common Names

Ethers are usually named by giving the name of each alkyl or aryl group, in alphabetical order, followed by the word *ether*.

Methyl ether	$\text{CH}_3-\text{O}-\text{CH}_3$	Ethyl methyl ether	$\text{CH}_3-\text{O}-\text{CH}_2\text{CH}_3$
Ethyl ether	$\text{CH}_3\text{CH}_2-\text{O}-\text{CH}_2\text{CH}_3$	Ethyl- <i>n</i> -propyl ether	$\text{CH}_3\text{CH}_2-\text{O}-\text{CH}_2\text{CH}_2\text{CH}_3$
Vinyl ether	$\text{CH}_2=\text{CH}-\text{O}-\text{CH}=\text{CH}_2$	<i>t</i> -Butyl methyl ether	$(\text{CH}_3)_3\text{C}-\text{O}-\text{CH}_3$
Phenyl ether		Methyl phenyl ether (anisole)	

### IUPAC System

For ethers with more complex structures, it may be necessary to name the -OR group as an alkoxy group. In the IUPAC system, the smaller alkoxy group is named as a substituent.



Dr Mohamed El-Newehy


## Physical Properties of Ethers

### Physical State

Ethers are colorless compounds with characteristic, relatively pleasant odors.

### Boiling Points

- They have **lower boiling points (bp,s) than alcohols** with an equal number of carbon atoms.
- In fact, an ether has nearly the **same bp as the corresponding hydrocarbon** in which a -CH<sub>2</sub>- group replaces the ether's oxygen.
- Because of their structures (**no O-H bonds**), ether molecules cannot form hydrogen bonds with one another.




Compound	Formula	bp	mol wt	Water solubility (g/100 mL, 20°C)
1-butanol	CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> OH	118°C	74	7.9
diethyl ether	CH <sub>3</sub> CH <sub>2</sub> -O-CH <sub>2</sub> CH <sub>3</sub>	35°C	74	7.5
pentane	CH <sub>3</sub> CH <sub>2</sub> -CH <sub>2</sub> -CH <sub>2</sub> CH <sub>3</sub>	36°C	72	0.03

Dr Mohamed El-Newehy

## Physical Properties of Ethers

### Solubility

- Low-molecular-weight ethers, such as dimethyl ether, are quite soluble in water.
- Ether molecules can form hydrogen bonds to water.



Structure	Name	Mol.Wt.	Bp (°C)	Solubility in H <sub>2</sub> O At 20 °C
CH <sub>3</sub> CH <sub>2</sub> CH <sub>3</sub>	propane	44	-42	insoluble
CH <sub>3</sub> OCH <sub>3</sub>	methyl ether	46	-24	soluble
CH <sub>3</sub> CH <sub>2</sub> OH	ethanol	46	78	soluble
CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	n-butane	58	-0.5	insoluble
CH <sub>3</sub> CH <sub>2</sub> OCH <sub>2</sub> CH <sub>3</sub>	ethyl methyl ether	60	8	soluble
CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> OH	1-propanol	60	97	soluble
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>3</sub> CH <sub>3</sub>	n-pentane	72	35	insoluble
CH <sub>3</sub> CH <sub>2</sub> OCH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	ethyl ether	74	36	7.5 g/100 g
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>2</sub> CH <sub>2</sub> OH	1-butanol	74	118	7.9 g/100 g
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>5</sub> CH <sub>3</sub>	n-heptane	100	98	insoluble
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>2</sub> O(CH <sub>2</sub> ) <sub>2</sub> CH <sub>3</sub>	n-propyl ether	102	91	0.2 g/100 g
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>4</sub> CH <sub>2</sub> OH	1-hexanol	102	157	0.6 g/100 g

Dr Mohamed El-Newehy

## Preparation of Ethers

- There are *two general methods* for synthesizing ethers.

### 1) Dehydration of alcohols

It is used **commercially** and in the **laboratory** to make certain **symmetrical ethers**.

### 2) Williamson synthesis


General **laboratory** method used to prepare all kinds of ethers, **symmetrical** and **unsymmetrical**.

Dr Mohamed El-Newehy

## Preparation of Ethers


### 1) Dehydration of alcohols

It takes place in the presence of acid catalysts ( $\text{H}_2\text{SO}_4$ ,  $\text{H}_3\text{PO}_4$ ) (intermolecular reaction)




#### Example;

The most important commercial ether is diethyl ether. It is prepared from ethanol and sulfuric acid.



#### ○ To prepare ethyl ether

- Dissolve ethyl alcohol in sulfuric acid at ambient temperature.
- Heat the solution to  $140^\circ\text{C}$  while adding more alcohol.




Dr Mohamed El-Newehy

## Preparation of Ethers

### Scope and Limitations

- When ethyl alcohol is dehydrated by **sulfuric acid at 180° C**, the dominant product is **ethylene**.




Dr Mohamed El-Newehy

## Preparation of Ethers

### 2) Williamson Synthesis

- This method has two steps;
  - 1) An alcohol is converted to its alkoxide by treatment with a reactive metal (sodium or potassium).



- 2) Displacement is carried out between the alkoxide and an alkyl halide.



- To obtain the best yields of mixed dialkyl ethers, we select a 1° rather than a 2° or 3° alkyl halide and react it with a sodium alkoxide
- To prepare an alkyl aryl ether, we must be careful not to pick a combination in which one of the reagents has a halogen directly attached to an aromatic ring.

Dr Mohamed El-Newehy

## Preparation of Ethers

### 2) Williamson Synthesis

**Example 1;** Preparation of *t*-butyl methyl ether,  $(\text{CH}_3)_3\text{C}-\text{O}-\text{CH}_3$ .


➤ In theory, this could be done by either of two reactions.

1. You could react sodium methoxide,  $\text{CH}_3\text{O}^-\text{Na}^+$ , with *t*-butyl chloride,  $(\text{CH}_3)_3\text{C}-\text{Cl}$ .

*This combination leads to dehydrohalogenation to an alkene, an elimination reaction.*

2. You could react sodium *t*-butoxide,  $(\text{CH}_3)_3\text{C}-\text{O}^-\text{Na}^+$ , with methyl chloride,  $\text{CH}_3\text{Cl}$ .

*This route gives the desired ether by substitution.*




Dr Mohamed El-Newehy

## Preparation of Ethers

### 2) Williamson Synthesis

**Example 2;** Assume you need to synthesize methyl phenyl ether (anisole),  $\text{CH}_3-\text{O}-\text{C}_6\text{H}_5$ , by the Williamson method.

➤ In theory, you could obtain anisole in either of two ways.



Dr Mohamed El-Newehy




## Reactions of Ethers


- **Ethers** are quite stable compounds.
- The **ether** linkage does not react with bases, reducing agents, oxidizing agents, or active metals.
- **Ethers** react only under strongly acidic conditions.

### Cleavage of Ethers by Hot Concentrated Acids


- When **ethers** are heated in concentrated acid solutions, the ether linkage is broken.



- The acids most often used in this reaction are HI, HBr, and HCl.
- If an excess of *acid* is present, the alcohol initially produced is converted into an alkyl halide by the reaction.




For example,



Dr Mohamed El-Newehy

## Cyclic Ethers


- It is possible to obtain cyclic ethers by an *intramolecular reaction*.



- THF is so named because it is related to the unsaturated cyclic ether *furan*.



- When 1,5-pentanediol is heated in acid, the six-membered ring ether *tetrahydropyran* is obtained.



- Tetrahydropyran is so called because it is related to the unsaturated ether *pyran*.



Dr Mohamed El-Newehy


## Uses of Ethers

- Ether is used as a mild anesthetic and as a solvent in industries
- It is used as an antiseptic to prevent infection when an injection is administered.
- Dimethyl ether is used as refrigerant and as solvent at low temperature.
- Diethyl ether is a common ingredient as an anesthetic in surgery.
- Diethyl ether is common solvent for oils, gums, resins, etc....
- We use phenyl ether as a heat transfer medium because of its high boiling point.

Dr Mohamed El-Newehy


## Epoxides

- Epoxides are cyclic ethers in which the ether oxygen is part of a three membered ring.



An epoxide

- The simplest and most important epoxide is ethylene oxide.
- Ethylene oxide is prepared commercially by air oxidation of ethylene.



Dr Mohamed El-Newehy