

Fundamentals of Organic Chemistry

CHEM 108

King Saud University

College of Science, Chemistry Department

Carbohydrates



- The word **carbohydrate** can be expressed as **hydrates of carbon** because molecular formulas of these compounds.

Example;

Glucose has the molecular formula $C_6H_{12}O_6$, which might be written as $C_6(H_2O)_6$.

- **Carbohydrates** are **polyhydroxyaldehydes**, **polyhydroxyketones**, or substances that give such compounds on hydrolysis.
- The chemistry of **carbohydrates** is mainly the combined chemistry of two functional groups: the **hydroxyl group** and the **carbonyl group**.


Carbohydrates

3

- **Carbohydrates** are usually classified according to their structure as **monosaccharides**, **oligosaccharides**, or **polysaccharides**.

The term **saccharide** comes from Latin (*saccharum*, sugar) and refers to the sweet taste of some simple carbohydrates.

- **The three classes of carbohydrates are related to each other through hydrolysis.**



Carbohydrates



- **Monosaccharides** (or **simple sugars**) are carbohydrates that cannot be hydrolyzed to simpler compounds.
- **Oligosaccharides** (from the Greek *oligos*, few) contain at least two and generally no more than a few linked monosaccharide units.

They may be called **disaccharides**, **trisaccharides**, and **so on**, depending on the number of units, which may be the same or different.

Example; Maltose is a disaccharide made of two glucose units.

Sucrose is made of two different monosaccharide units: glucose and fructose.

- **Polysaccharides** contain many monosaccharide units - sometimes hundreds or even thousands. Usually, but not always, the units are identical.




Example;

Starch and cellulose, contain linked units of the same monosaccharide, glucose.

Monosaccharides

5


- **Monosaccharides** are classified according to:
 - The **number of carbon atoms** present (**triose, tetrose, pentose, hexose, and so on**).
 - Whether the carbonyl group is present as an aldehyde (**aldose**) or as a ketone (**ketose**).
- There are only two trioses: glyceraldehyde and dihydroxyacetone.
- Glyceraldehyde is the simplest aldose, and dihydroxyacetone is the simplest ketose.
- Each is related to glycerol in that each has a carbonyl group in place of one of the hydroxyl groups.



Monosaccharides

6


- Other **aldoses or ketoses** can be derived from glyceraldehyde or dihydroxyacetone by adding carbon atoms, each with a hydroxyl group.
- In **aldoses**, the chain is numbered from the aldehyde carbon.
- In most **ketoses**, the carbonyl group is located at C-2.



Monosaccharides


7

Chirality in Monosaccharides; Fischer Projection Formulas and D,L-Sugars




- He used a small capital d to represent the configuration of (+)-glyceraldehyde, with the hydroxyl group on the *right*; its enantiomer, with the hydroxyl group on the *left*, was designated L-(2)-glyceraldehyde.

The most oxidized carbon (CHO) was placed at the top.



Monosaccharides



Monosaccharides


9

- **Diastereomers;** when two or more stereoisomers of a compound have different configurations at one or more (but not all) of the equivalent (related) stereocenters and are not mirror images of each other.
- **Epimers;** A special name is given to diastereomers that differ in configuration *at only one stereogenic center*.
- Each pair has the same configurations at all **stereogenic centers** except one.
- **Examples;**
 - D-(-)-Erythrose and D-(-)-threose are not only diastereomers, they are epimers.
 - D-glucose and D-mannose are epimers (at C-2).
 - D-glucose and D-galactose are epimers (at C-4).
- **Dextrorotation (+)** and **levorotation (-)** are terms describing circular direction of rotating plane-polarized light.
 - **Dextrorotatory (+);** If the light rotates clockwise, (rotation to the right).
 - **Levorotatory (-);** if it rotates counterclockwise, (rotation to the left).

The Cyclic Hemiacetal Structures of Monosaccharides

10

- **Monosaccharides** exist mainly in cyclic, hemiacetal forms and not in the acyclic aldo- or keto-forms.




Manipulation of the Fischer projection formula of D-glucose to bring the C-5 hydroxyl group in position for cyclization to the hemiacetal form.

Haworth Projection

11

- The carbons are arranged clockwise numerically, with C-1 at the right.
- Substituents attached to the ring lie above or below the plane.




Haworth projection formulas for D-glucose

- **Carbons 1 through 5** are part of the ring structure, but carbon 6 (the $-\text{CH}_2\text{OH}$ group) is a substituent on the ring.
- **C-1 is special.**
C-1 is the hemiacetal carbon (it carries a hydroxyl group, and it is also connected to C-5 by an ether linkage).
- **C-2, C-3, and C-4** are secondary alcohol carbons.
- **C-6** is a primary alcohol carbon.
- **Hydroxyl groups on the right** in the Fischer projection are *down* in the Haworth projection (and conversely).
- **hydroxyl groups on the left** in the Fischer projection are *up* in the Haworth projection).
- For **D-sugars**, the terminal $-\text{CH}_2\text{OH}$ group is *up* in the Haworth projection; for **L-sugars**, it is down.

Anomeric Carbons; Mutarotation

12

- **Anomeric carbon;** The hemiacetal carbon, the carbon that forms the new stereogenic center.
- **Anomers;** Two monosaccharides that differ only in configuration at the anomeric center are (a special kind of epimers).
- **Anomers** are called α or β , depending on the position of the hydroxyl group.
- **For monosaccharides** in the D-series, the hydroxyl group is “down” in the α anomer and “up” in the β anomer.




- If D-glucose is crystallized from methanol, the pure α form is obtained.
- Crystallization from acetic acid gives the β form.
- The α and β forms of D-glucose are *diastereomers*.

Conformations of Pyranoses; Chair Conformation

13


- **At the anomeric carbon (C-1)**, where the hydroxyl group may be axial (in the α anomer) or equatorial (in the β anomer).



1) Reduction of Monosaccharides

14

- The **carbonyl group of aldoses and ketoses** can be reduced by various reagents to give polyols, called alditols.
- **Example;**
Catalytic hydrogenation or reduction with sodium borohydride (NaBH_4) converts D-glucose to D-glucitol (sorbitol).
- **Sorbitol** is used commercially as a sweetener and sugar substitute.



2) Oxidation of Monosaccharides

15

2.1. With Mild Oxidizing Agents


- The oxidation of aldoses is so easy that they react with such mild oxidizing agents as
 - Tollens' reagent (Ag^+ in aqueous ammonia),
 - Fehling's reagent (Cu^{2+} complexed with tartrate ion),
 - Benedict's reagent (Cu^{2+} complexed with citrate ion).
- These aldehyde groups can be easily oxidized to acids which are called aldonic acids.
- A carbohydrate that reacts with Ag^+ or Cu^{2+} is called a reducing sugar

2) Oxidation of Monosaccharides


16

2.1. With Mild Oxidizing Agents

- **Example;** D-glucose is easily oxidized to D-gluconic acid.



- With the copper reagents, the blue solution gives a red precipitate of cuprous oxide, Cu_2O .




2) Oxidation of Monosaccharides

17

2.2. With Strong Oxidizing Agents

- Stronger oxidizing agents, such as **aqueous nitric acid**.
- The aldehyde group and the primary alcohol group can be oxidized, producing dicarboxylic acids called aldaric acids.
- **Example;**


D-glucose gives D-glucaric acid.




Disaccharides




- **The most common oligosaccharides are disaccharides.**
- **In a disaccharide,**
two monosaccharides are linked by a glycosidic bond between the anomeric carbon of one monosaccharide unit and a hydroxyl group on the other unit.




- **Maltose** is the disaccharide obtained by the partial hydrolysis of starch.
- Further hydrolysis of maltose gives only **D-glucose**.
- **Maltose** must, therefore, consist of two linked glucose units.
- It turns out that the anomeric carbon of the left unit is linked to the C-4 hydroxyl group of the unit at the right as an acetal (glycoside).
- The configuration at the anomeric carbon of the left unit is α .



- **Cellobiose** is the disaccharide obtained by the partial hydrolysis of cellulose.
- Further hydrolysis of cellobiose gives only **D-glucose**.
- **Cellobiose** must therefore be an isomer of maltose.
- In fact, **cellobiose** differs from maltose only in having the β configuration at C-1 of the left glucose unit.





- **Lactose** is the major sugar in human and cow's milk (4% to 8% lactose).
- Hydrolysis of lactose gives equimolar amounts of **D-galactose and D-glucose**.
- The anomeric carbon of the galactose unit has the β configuration at C-1 and is linked to the hydroxyl group at C-4 of the glucose unit.



- **The most important commercial disaccharide is sucrose, ordinary table sugar.**
- **Sucrose** occurs in all photosynthetic plants, where it functions as an energy source.
- It is obtained commercially from sugar cane and sugar beets, in which it constitutes 14% to 20% of the plant juices.
- **Sucrose** is very water soluble (2 grams per milliliter at room temperature) because it is polar due to the presence of eight hydroxyl groups on its surface.
- Hydrolysis of sucrose gives equimolar amounts of **D-glucose and the ketose D-fructose.**


Polysaccharides

- **Polysaccharides** contain many linked monosaccharides and vary in chain length and molecular weight.
- Most **polysaccharides** give a single monosaccharide on complete hydrolysis.
- The monosaccharide units may be linked linearly, or the chains may be branched.



- **Starch** is the energy-storing carbohydrate of plants.
- It is a major component of cereals, potatoes, corn, and rice.
- **Starch** is made up of glucose units joined mainly by 1,4- α -glycosidic bonds, although the chains may have a number of branches attached through 1,6- α -glycosidic bonds.
- **Partial hydrolysis of starch gives maltose, and complete hydrolysis gives only D-glucose.**
- **Starch** can be separated by various techniques into two fractions: amylose and amylopectin.
- **In amylose**, which constitutes about 20% of starch, the glucose units (50 to 300) are in a continuous chain, with 1,4 linkages.
- **Amylopectin** is highly branched. Although each molecule may contain 300 to 5000 glucose units, chains with consecutive 1,4 links average only 25 to 30 units in length.
- These chains are connected at branch points by **1,6 linkages**.

- **Glycogen** is the energy-storing carbohydrate of animals.
- Like starch, it is made of 1,4- and 1,6-linked glucose units.
- **Glycogen** has a higher molecular weight than starch (perhaps 100,000 glucose units), and its structure is even more branched than that of amylopectin, with a branch every 8 to 12 glucose units.
- **Glycogen** is produced from glucose that is absorbed from the intestines into the blood; transported to the liver, muscles, and elsewhere; and then polymerized enzymatically.
- **Glycogen** helps maintain the glucose balance in the body by removing and storing excess glucose from ingested food and later supplying it to the blood when various cells need it for energy.



- **Cellulose** is an *unbranched* polymer of glucose joined by 1,4-β-glycosidic bonds.
- It consists of linear chains of **cellobiose units**.
- These linear molecules, containing an average of 5000 glucose units, aggregate to give fibrils bound together by hydrogen bonds between hydroxyls on adjacent chains.
- **Cellulose fibers** having considerable physical strength are built up from these fibrils, wound spirally in opposite directions around a central axis.
- Wood, cotton, hemp, linen, straw, and corncobs are mainly **cellulose**.