




PCR Troubleshooting


Polymerase Chain Reaction (PCR)=DNA Photocopier




Performing PCR steps :




Common Issues in PCR:




Low or no amplification



Non-specific band or primer dimer



Incorrect product size




Smeared Bands

1- No Band or Faint Band:

Causes Related to Cycling Times and Temp.

- Too Few cycles were used.
- Extension time was too short.
- **Incorrect annealing temperature.**
- Denaturation temperature was too low.




Causes Related to PCR Components

- **No enough template was in the reaction.**
- Primer concentration was too low.
- Impure primers, dNTPs, or water
- **PCR product has high GC content.**
- Primers were designed or synthesized incorrectly.
- **Not enough Mg^{2+} .**

2- Nonspecific Bands or Primer Dimer:

Causes Related to Cycling Times and Temp.

- **Annealing temperature** was too low.
- Too many cycles were used.
- Extension time was too long.




Causes Related to PCR Components

- **Too much primer** was added.
- **Too much Mg²⁺** was added.
- Primers were designed or synthesized incorrectly by user or manufacturer.

3- Incorrect PCR product size:

Causes Related to Cycling Times and Temp.

- **Incorrect annealing temperature.**




Causes Related to PCR Components

- Mispriming.
- **Improper Mg²⁺ concentration.**
- Impure primers, dNTPs, or water
- Primers were designed or synthesized incorrectly by user or manufacturer.

4- Smeared Band:

Causes Related to Cycling Times and Temp.

- Too many cycles were used.



Causes Related to PCR Components

- **Too much template was added.**
- Impure primers, dNTPs, or water.
- **Template contained an exonuclease or was degraded.**



Common PCR additive reagents:

1. Additives that benefit GC Rich templates:

→1-10% DMSO (Dimethylsulfoxid):

- GC rich (GC content >60%).
- Lowering the T_m.

→Q solution:

- High degree of secondary structure.
- GC-rich.
- Increases PCR specificity.




→PCRx Enhancer:

- For problematic and/or GC-rich templates.
- Higher primer specificity, broader magnesium concentration optima, broader annealing temperature optima.

2. Additives That Help PCR in the Presence of Inhibitors:

→400 ng/μl BSA (Bovine serum albumin).

→Non-ionic detergents: Ex: 0.1 to 1% Triton X.





Home Work:

- TERT is a gene code for telomerase, an enzyme that elongate the telomere and mutated in some cancers. Your aim is to amplify a region in its promoter using PCR technique.
1. Draw a flow chart that illustrate the steps of performing PCR .
 2. Design a primer using primer3plus tool.
 3. Check the specificity of the primer.

**