
# Introduction to Organic Chemistry
# CHEM 109

جامعة الملك سعود
King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

## *Chapter 6: Chemistry of Biomolecules*



جامعة الملك سعود
King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT


# Carbohydrates

# Chapter 6: Chemistry of Biomolecules

King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT


## 6.1. Carbohydrates

- **Carbohydrates** are polyhydroxy aldehydes or ketones, or compounds that can be hydrolyzed to produce such molecules.
- **Carbohydrates** typically follow the general chemical formula $C_x(H_2O)_y$. i.e. carbohydrate can be expressed as hydrates of carbon
- *Example;* Glucose has the molecular formula $C_6H_{12}O_6$, which might be written as $C_6(H_2O)_6$.




D-Glucose

**polyhydroxyaldehydes**




D-Fructose

**polyhydroxyketones**


### *6.1.1. Classification of Carbohydrates*

- **Monosaccharides** are the simplest carbohydrates and cannot be further hydrolyzed into smaller carbohydrate units.
- **Disaccharides** can be broken down by hydrolysis into two monosaccharide units.
- **Oligosaccharides** produce between two and ten monosaccharides upon hydrolysis.
- **Polysaccharides** yield more than ten monosaccharides when hydrolyzed.


COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

## 6.1.2. Monosaccharides


- Glyceraldehyde is the simplest aldose, and dihydroxyacetone is the simplest ketose.
- Each is related to glycerol in that each has a carbonyl group in place of one of the hydroxyl groups.




**Glyceraldehyde (an aldotriose)**





**Dihydroxyacetone (a ketotriose)**




D-(+)-Glyceraldehyde


and




L-(-)-Glyceraldehyde

### 6.1.2.1. Classification of Monosaccharides

- **Monosaccharides** are classified based on two main features:
  1. The number of carbon atoms they contain (triose, tetrose, pentose, and hexose).
  2. The type of carbonyl group present — either an aldehyde or a ketone group.




جامعة الملك سعود
King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

## 6.1.2. Monosaccharides

### 6.1.2.2. D- and L- Designations of Monosaccharides

- Chiral (asymmetric) carbon atom bonded to four different groups.
- The simplest carbohydrates are glyceraldehyde, which is chiral and assigned as L or D-configuration, and dihydroxyacetone, which is achiral.
- D-sugar if its highest-numbered chiral center has the same configuration as D-(+)-glyceraldehyde.
- L-sugar if its highest-numbered chiral center shares the same configuration as L-(–)-glyceraldehyde.





**A D-aldopentose** **An L-ketohexose**

جامعة الملك سعود
King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

## 6.1.2. Monosaccharides

### 6.1.2.3. Structural Formulas for Monosaccharides

- **Fischer projection**
  - Represents sugar stereochemistry in the open-chain form.
  - Horizontal lines → bonds projecting out of the plane (toward the viewer).
  - Vertical lines → bonds behind the plane (away from the viewer).
  - The –CHO group is placed at the top, and the –$CH_2OH$ group at the bottom.




King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

## 6.1.2. Monosaccharides

### 6.1.2.3. Structural Formulas for Monosaccharides

- The Ruff degradation and the Kiliani–Fischer synthesis allow us to place all of the aldoses into families or "family trees" based on their relation to D- or L-glyceraldehyde.
- Most, but not all, of the naturally occurring aldoses belong to the D family, with D-(+)-glucose being by far the most common.




King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

## 6.1.2. Monosaccharides

### 6.1.2.3. Structural Formulas for Monosaccharides

- **Haworth formulas**
  - Represent sugar stereochemistry in the cyclic form, which exists in equilibrium with the open-chain form.
  - The cyclic (hemiacetal) forms are produced by an intramolecular reaction between the –OH group at C5 and the aldehyde group at C1.
  - Cyclization creates a new chiral center at C1, explaining the existence of two cyclic forms.
  - These two forms are differing only in the configuration at C1 (the anomeric carbon).
    - α-anomer → the hemiacetal –OH is *trans* to the –$CH_2OH$ group.
    - β-anomer → the hemiacetal –OH is *cis* to the –$CH_2OH$ group.



جامعة الملك سعود
King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT


## 6.1.2. Monosaccharides

### 6.1.2.4. Isomerization

### Enantiomers

- **Stereogenic centers** are mostly carbon atoms (*asymmetric carbon*) that bind four different groups.
- **Stereoisomers = $2^n$** (n = number of stereogenic centers).

- **Epimers:** A special name is given to diastereomers that differ in configuration at only one stereogenic center as shown.




**D-glucose and D-mannose are epimers (at C-2).**

**D-glucose and D-galactose are epimers (at C-4).**


King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

## 6.1.2. Monosaccharides


### 6.1.2.5. Oxidation Reactions of Monosaccharides

### A) Benedict's or Tollens' Reagents: Reducing Sugars

- **Benedict's reagent:** an alkaline solution containing a cupric citrate complex ion.




- **Tollens' solution:** $[Ag(NH_3)_2\bar{O}H]$




COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

## 6.1.2. Monosaccharides

### 6.1.2.5. Oxidation Reactions of Monosaccharides

### A) Benedict's or Tollens' Reagents: Reducing Sugars

- They both oxidize and give positive tests with aldoses and ketoses even those that exist primarily as cyclic hemiacetals.
- Carbohydrates that contain hemiacetal groups give positive tests with Benedict's or Tollens' solutions, and they are called *reducing sugars*.
- Carbohydrates that contain only acetal groups do not give positive tests with Benedict's or Tollens' solutions, and they are called *nonreducing sugars*.




King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

## 6.1.2. Monosaccharides

### 6.1.2.5. Oxidation Reactions of Monosaccharides

### B) Bromine Water: The Synthesis of Aldonic Acids


- Bromine water is a general reagent that selectively oxidizes the −CHO group to a $-CO_2H$ group, thus converting an aldose to an **aldonic acid**.





Aldose → Aldonic acid

### C) Nitric Acid Oxidation: Aldaric Acids

- Dilute nitric acid is a stronger oxidizing agent than bromine water , it oxidizes both the −CHO group and the terminal $-CH_2OH$ group of an aldose to $-CO_2H$ groups, forming dicarboxylic acids are known as **aldaric acids**.




Aldose → Aldaric acid




D-Glucose → D-Glucaric acid


King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

## 6.1.2. Monosaccharides


### 6.1.2.5. Oxidation Reactions of Monosaccharides

### D) Periodate Oxidations: Oxidative Cleavage of Polyhydroxy Compounds

- Compounds with adjacent hydroxyl groups (vicinal diols) undergo oxidative cleavage by aqueous periodic acid ($HIO_4$), breaking the C–C bond to yield carbonyl compounds (aldehydes, ketones, or carboxylic acids).




- In chains with three or more contiguous –CHOH groups, the internal ones produce formic acid.
- Cleavage also occurs when an –OH group is adjacent to the carbonyl of an aldehyde or ketone (but not acids or esters).




جامعة الملك سعود
King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT


## 6.1.2. Monosaccharides

### 6.1.2.6. Reduction of Monosaccharides: Alditols

- Aldoses (and ketoses) can be reduced with sodium borohydride to give compounds called **alditols**.




- Example; Reduction of D-glucose, yields D-glucitol.




King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

## 6.1.3. Disaccharides

### 6.1.3.1. Sucrose


- Sucrose is the most widely occurring disaccharide, obtained commercially from sugarcane or sugar beets.
- It is a nonreducing sugar and has the molecular formula $C_{12}H_{22}O_{11}$.
- Acid-catalyzed hydrolysis of 1 mole of sucrose yields 1 mole of D-glucose and 1 mole of D-fructose.




(+)-Sucrose
α-D-Glucopyranosyl-(1→2)-β-D-fructofuranoside

### 6.1.3.2. Maltose

- Partial hydrolysis of starch gives Maltose.
- When 1 mol of maltose is subjected to acid-catalyzed hydrolysis, it yields 2 mol of D-(+)-glucose.
- Maltose is a reducing sugar; it gives positive tests with Benedict's, and Tollens' solutions.





(+)-maltose,
α-D-Glucopyranosyl-(1→4)- β- D-glucopyranose

جامعة الملك سعود
King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

## 6.1.3. Disaccharides

### 6.1.3.3. Cellobiose


- Partial hydrolysis of cellulose gives cellobiose.
- Cellobiose resembles maltose in every respect except one: the configuration of its 1→4 glycosidic linkage.
- Cellobiose is a reducing sugar that, on acid-catalyzed hydrolysis, yields two molar equivalents of D-glucose




Cellobiose,
β-D-Glucopyranosyl-(1→4)- β-D-glucopyranose

### 6.1.3.4. Lactose

- Lactose is a disaccharide present in the milk of humans, cows, and almost all other mammals.
- Lactose is a reducing sugar that hydrolyzes to yield D-glucose and D-galactose; the 1→4 glycosidic linkage is β.




Lactose,
β-D-Galactopyranosyl-(1→4)-β-D-glucopyranose

جامعة الملك سعود
King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

## 6.1.4. Polysaccharides

- Polysaccharides are polymers consisting of monosaccharides joined together by glycosidic linkages.
- Homopolysaccharides are polymers made of a single type of monosaccharide.
- Heteropolysaccharides are polymers made of more than one type of monosaccharide.

### 6.1.4.1. Starch


- Starch occurs as microscopic granules in roots, tubers, and seeds (e.g., corn, potatoes, wheat, rice).
- Heating starch with water swells the granules, forming a colloidal suspension.
- Starch is a nonreducing polysaccharide composed of amylose (10–20%) and amylopectin (80–90%).

King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT


## 6.1.4. Polysaccharides

### 6.1.4.1. Starch

- **Amylose** typically consists of more than 1000 D-glucopyranoside units connected in $\alpha$ linkages between C1 of one unit and C4 of the next.
- **Amylopectin** has a structure similar to amylose, with $\alpha(1\rightarrow4)$ glycosidic links in the linear chains, but its structure is branched by $\alpha(1\rightarrow6)$ glycosidic bonds.
- The branching occurs between C6 of one glucose unit and C1 of another.
- Branch points appear approximately every 20–25 glucose units.
- **Amylopectin** has a molecular weight of 1–6 million, consisting of hundreds of interconnected chains, each containing about 20–25 glucose units.




Partial structure of Amylose




Partial structure of Amylopectin

جامعة الملك سعود
King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT


## 6.1.4. Polysaccharides

### 6.1.4.2. Glycogen

- Glycogen has a structure very much like that of amylopectin; however, in glycogen the chains are much more highly branched.
- Glycogen has a very high molecular weight, and samples isolated under conditions minimizing hydrolysis show molecular weights as high as 100 million.

### 6.1.4.3. Cellulose

- Cellulose consists of D-glucopyranose units linked by β(1→4) glycosidic bonds forming very long, unbranched chains.
- The β-linkages make cellulose chains linear, unlike the helical α(1→4) chains in starch and glycogen.
- Hydroxyl (–OH) groups on adjacent chains form extensive hydrogen bonds, producing a rigid, insoluble, fibrous structure.
- The precise stereochemistry of D-glucose allows the formation of strong linear fibers ideal for plant cell walls.




Partial structure of Cellulose

جامعة الملك سعود
King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

# Lipids


جامعة الملك سعود
King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

## 6.2. Lipids


- **Lipids** are compounds of biological origin that dissolve in nonpolar solvents, such as chloroform and diethyl ether.
- The name lipid comes from the Greek word *lipos*, for fat.

### *6.2.1. Fatty Acids and Triacylglycerols*

- **Triacylglycerols (Triglycerides)** are the oils of plants and the fats of animal origin.
- **Triacylglycerols** are carboxylic acids of biological origin that are found as esters of glycerol.
- The groups R, R′, and R″ are usually long chain alkyl groups that may also contain one or more carbon–carbon double bonds and may all be different.
- **Triacylglycerols** include such common substances as peanut oil, soybean oil, corn oil, sunflower oil, butter, lard, and tallow.




Glycerol





A triacylglycerol

جامعة الملك سعود
King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

## 6.2.1. Fatty Acids and Triacylglycerols

- **Triacylglycerols** are known as oils and fats.
- **Fats** are solid at room temperature due to saturated fatty acids.
- **Oils** are liquid because of higher unsaturation.
- **Triacylglycerols** can be
  - **Simple triacylglycerols**; all three acyl groups are the same.
  - **Mixed triacylglycerols**; the acyl groups are different.
- Hydrolysis of fats or oils yields fatty acids that are typically unbranched and contain an even number of carbon atoms.





A triacylglycerol


King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

## 6.2.1. Fatty Acids and Triacylglycerols

- **Unsaturated fatty acids,**
  - Unsaturated fatty acids contain *cis* double bonds.
  - Many of them have two or three double bonds.
  - Such fats are called polyunsaturated fats or oils.
  - The first double bond usually occurs between C9 and C10
- **Saturated fatty acids** have relatively high melting points.
- **Omega-3 fatty acids** are those where the third to last carbon in the chain is part of a carbon–carbon double bond.




DHA, an omega-3-fatty acid
[(4Z,7Z,10Z,13Z,16Z,19Z)-4,7,10,13,16,19-docosahexaenoic acid]



King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

## 6.2.2. Hydrogenation of Triacylglycerols


- **Fats:** Triacylglycerols high in saturated fatty acids → *solid at room temperature, high melting points*.
- **Oils:** Triacylglycerols high in unsaturated/polyunsaturated fatty acids → *liquid at room temperature, lower melting points*.
- Catalytic hydrogenation converts unsaturated triacylglycerols into more saturated forms.
- Commercial solid fats are produced by partial hydrogenation of vegetable oils, while complete hydrogenation is avoided because fully saturated triacylglycerols are hard and brittle.



جامعة الملك سعود
King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

## 6.2.3. Reactions of the Carboxyl Group of Fatty Acids

- **Fatty acids** undergo reactions typical of carboxylic acids.
  - They react with $LiAlH_4$ to form alcohols,
  - They react with alcohols and mineral acid to form esters,
  - They react with bromine and phosphorus to form $\alpha$-halo acids,
  - They react with thionyl chloride to form acyl chlorides




جامعة الملك سعود
King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

# Amino Acids and Proteins


King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

## 6.3. Amino Acids and Proteins

- **Proteins** have many diverse functions; they are major components of the following biomolecules.
  - Enzymes and hormones which catalyze and regulate biological reactions.
  - Muscles and tendons which provide the body with means for movement.
  - Hemoglobin which carries oxygen to all parts of the body.
  - Antibodies they are integral parts of the immune system.
- **The 20 $\alpha$-amino acids** are the essential building blocks of proteins and peptides, forming polyamides through peptide bonds.




**An $\alpha$-amino acid**
***R* is a side chain at the $\alpha$ carbon that determines the identity of the amino acid (Table 24.1).**





**A portion of a protein molecule**
***Amide (peptide) linkages are shaded.***
***$R_1$–$R_5$ may be any of the possible side chains.***

King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

## 6.3.1. Amino acids

### 6.3.1.1. Structure and Names

- **All natural occurring amino acids** are chiral (except glycine which is achiral) and have the L-configuration (as related to glyceraldehyde) at the $\alpha$-carbon.




**An L-$\alpha$-amino acid [usually an (*S*)-$\alpha$-amino acid]** **L-Glyceraldehyde [(*S*)-glyceraldehyde]**

**Fischer projections for an L-$\alpha$-amino acid and L-glyceraldehyde**

- **Nomenclature and Classification of amino acids:**
  - The 20 amino acids are known by their common names (IUPAC nomenclature could be applied).
  - Abbreviated by **three letter** based on its common name, which is used when writing the formulas of peptides.
  - **One-letter** abbreviation used to describe the amino acid sequence in a protein.
  - **Example;** Glycine = Gly or (G); Alanine = Ala or (A); Valine = Val or (V), etc..

جامعة الملك سعود
King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

## 6.3.1. Amino acids

### 6.3.1.2. Classification of amino acids

- **Amino acids are classified according to the structure of R group:**

1. The acidity of R group: neutral or acidic or basic.
   - *Neutral R Group*: are those which containing one amino group and one carboxylic group.

     The majority of amino acids contain neutral R group.
   - *Acidic R group*: are those amino acids that contain more than one carboxylic group.

     Example; Aspartic, Glutamic --------"**hydrophilic"**
   - *Basic R group*: are those amino acids that contain more than one N atom.

     Example; Lysine, Arginine ---------"**hydrophilic"**
2. Polarity of the R group: Amino acid could be classified into polar (hydrophilic) or non-polar (hydrophobic).

جامعة الملك سعود
King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

## 6.3.1. Amino acids

### 6.3.1.2. Classification of amino acids

- **Essential and non-essential amino acid:**
  - *Essential amino acid:* amino acids, which cannot be synthesized by adult humans and therefore must be included in the diet in the form of proteins.


    Eight amino acids; Valine. Leucine, Isoleucine, Threonine, Methionine, Phenylalanine, Tryptophan, and Lysine.
  - *Non-essential amino acids:* amino acids, which can be synthesized in the body from other foods.


    Twelve amino acids; Glycine, Alanine, Serine, Cysteine, Proline, Tyrosine, Aspartic acid, Glutamic acid, Asparagine, Glutamine, Arginine, and Histidine.


King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

## 6.3.1. Amino acids

### 6.3.1.2. Classification of amino acids









| Structure of R | Name | Abbreviations[a] | $pK_{a_1}$ $\alpha$-$CO_2H$ | $pK_{a_2}$ $\alpha$-$NH_3^+$ | $pK_{a_3}$ R group | pI |
|---|---|---|---|---|---|---|
| **Neutral Amino Acids** | | | | | | |
| —H | Glycine | G or Gly | 2.3 | 9.6 | | 6.0 |
| —$CH_3$ | Alanine | A or Ala | 2.3 | 9.7 | | 6.0 |
| —$CH(CH_3)_2$ | Valine[b] | V or Val | 2.3 | 9.6 | | 6.0 |
| —$CH_2CH(CH_3)_2$ | Leucine[b] | L or Leu | 2.4 | 9.6 | | 6.0 |
| —$CHCH_2CH_3$ <br> $CH_3$ | Isoleucine[b] | I or Ile | 2.4 | 9.7 | | 6.1 |
| —$CH_2$  | Phenylalanine[b] | F or Phe | 1.8 | 9.1 | | 5.5 |
| —$CH_2CONH_2$ | Asparagine | N or Asn | 2.0 | 8.8 | | 5.4 |
| —$CH_2CH_2CONH_2$ | Glutamine | Q or Gln | 2.2 | 9.1 | | 5.7 |
| —$CH_2$  | Tryptophan[b] | W or Trp | 2.4 | 9.4 | | 5.9 |
|  (complete structure) | Proline | P or Pro | 2.0 | 10.6 | | 6.3 |

King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

## 6.3.1. Amino acids

### 6.3.1.2. Classification of amino acids





| Structure of R | Name | Abbreviations[a] | p$K_{a_1}$ $\alpha$-$CO_2H$ | p$K_{a_2}$ $\alpha$-$NH_3^+$ | p$K_{a_3}$ R group | p*I* |
|---|---|---|---|---|---|---|
| **Neutral Amino Acids** | | | | | | |
| —$CH_2OH$ | Serine | S or Ser | 2.2 | 9.2 | | 5.7 |
| —CHOH<br>$CH_3$ | Threonine[b] | T or Thr | 2.6 | 10.4 | | 6.5 |
| —$CH_2$—⟨benzene⟩—OH | Tyrosine | Y or Tyr | 2.2 | 9.1 | 10.1 | 5.7 |
| HOC(=O)—CH(NH—$CH_2$—$CH_2$)—CH(OH) | Hydroxyproline | Hyp | 1.9 | 9.7 | | 6.3 |
| —$CH_2SH$ | Cysteine | C or Cys | 1.7 | 10.8 | 8.3 | 5.0 |
| —$CH_2$—S<br>—$CH_2$—S | Cystine | Cys-Cys | 1.6<br>2.3 | 7.9<br>9.9 | | 5.1 |
| —$CH_2CH_2SCH_3$ | Methionine[b] | M or Met | 2.3 | 9.2 | | 5.8 |


(complete structure)

King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

## 6.3.1. Amino acids

### 6.3.1.2. Classification of amino acids




| Structure of R | Name | Abbreviations[a] | p$K_{a_1}$ $\alpha$-$CO_2H$ | p$K_{a_2}$ $\alpha$-$NH_3^+$ | p$K_{a_3}$ R group | p*I* |
|---|---|---|---|---|---|---|
| **R Contains an Acidic (Carboxyl) Group** | | | | | | |
| $-CH_2CO_2H$ | Aspartic acid | D or Asp | 2.1 | 9.8 | 3.9 | 3.0 |
| $-CH_2CH_2CO_2H$ | Glutamic acid | E or Glu | 2.2 | 9.7 | 4.3 | 3.2 |
| **R Contains a Basic Group** | | | | | | |
| $-CH_2CH_2CH_2CH_2NH_2$ | Lysine[b] | K or Lys | 2.2 | 9.0 | 10.5[c] | 9.8 |
| $-CH_2CH_2CH_2NH-C(=NH)-NH_2$ | Arginine | R or Arg | 2.2 | 9.0 | 12.5[c] | 10.8 |
|  | Histidine | H or His | 1.8 | 9.2 | 6.0[c] | 7.6 |


[a]Single-letter abbreviations are now the most commonly used form in current biochemical literature.
[b]An essential amino acid.
[c]p$K_a$ is of protonated amine of R group.

جامعة الملك سعود
King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

## 6.3.1. Amino acids

### 6.3.1.3. Amino acids as dipolar ions

- In the dry solid state amino acids exist as dipolar ions (zwittrions or inner salts).
- In aqueous solutions an equilibrium exists between the dipolar ion, the cationic and the anionic forms of the amino acid.
- The predominant form depends on the pH of the solution.
  - At low pH, the amino acid exists primarily in the cationic form.
  - At high pH, the amino acid exists primarily in the anionic form.
  - At some intermediate pH, called the P*I* (*isoelectric point*), the concentration of the dipolar ion is at maximum, and the concentrations of anionic and cationic forms are equal
- Each individual amino acid has a characteristic P*I*.




**Cationic form (predominant in strongly acidic solutions, e.g., at pH 0)** | **Dipolar ion** | **Anionic form (predominant in strongly basic solutions, e.g., at pH 14)**


King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

## 6.3.1. Amino acids


### 6.3.1.4. Synthesis of Amino acids

- There are many methods to prepare amino acids, two methods will be mentioned here.


1. **Direct Ammonolysis** of $\alpha$-halo-carboxylic acid with this method, yields tend to be poor.




2. **The Strecker Synthesis** (Reductive amination) treatment of an aldehyde with ammonia and hydrogen cyanide yields an $\alpha$-aminonitrile, which is hydrolyzed to the $\alpha$-amino acid.





The mechanism proceeds via an intermediate imine




King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT


## 6.3.2. Polypeptides and Proteins

- Enzymes polymerize amino acids by forming amide linkages.
- The polymer is called a peptide, and the amide linkages are called peptide bonds or peptide linkages.
- Each amino acid in the peptide is called an amino acid residue.
- Proteins can contain one or more peptide chains and others associated molecules or metal ions.




- **Polypeptides** are linear polymers.
- One end of a polypeptide chain in an amino acid residue has a free amino group -NH3, the other terminate in amino acid residue with free carboxylic group –CO2. These two groups are called the N-terminal and the C-terminal residues, respectively.
- **Polypeptides** are usually written with the N-terminal residue to the left and the C-terminal residue to the right.




King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

## 6.3.2. Polypeptides and Proteins

### 6.3.2.1. Hydrolysis of Polypeptides

- Hydrolysis of all amide linkages of a protein or polypeptide usually takes place by refluxing with 6M HCl for 24 hours.
- This hydrolysis results in the liberation of amino acids.
- The individual amino acids can be separated from each other using chromatographic methods.
- Example;

The complete hydrolysis of the tripeptide Glycylvalylpheneylalanine (Gly-Val-Phe) will give the three amino acids glycine, valine, and phenylalanine.




جامعة الملك سعود
King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

## 6.3.2. Polypeptides and Proteins

### 6.3.2.2. Primary Structure of Polypeptides and Proteins

- Primary structure; specific sequence of amino acid residues in a polypeptide or protein
- Determining the exact sequence is very important.
- First step: Identify the N-terminal and C-terminal residues.



جامعة الملك سعود
King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

# Nucleic Acids


جامعة الملك سعود
King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

## 6.4. Nucleic Acids

- Deoxyribonucleic acid (DNA) and ribonucleic acid (RNA) are molecules that carry genetic information in cells.
- DNA is the molecular archive of instructions for protein synthesis.
- RNA molecules transcribe and translate the information from DNA for the mechanics of protein synthesis.

### *6.4.1. The General Structure of Nucleic acids; Nucleotides and Nucleosides*

- Nucleic acids are linear polymers made up of nucleotide units.
- Hydrolysis of nucleic acids releases nucleotides, the basic building blocks of these molecules.



King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT


## 6.4.1. The General Structure of Nucleic acids; Nucleotides and Nucleosides

### 6.4.1.1. Nucleotides


- The Nucleotide Unit is composed of:
  1. A heterocyclic base from either the purine or pyrimidine family.

     **The heterocyclic base** of a nucleotide is attached through an *N*-glycosidic linkage to C1′ of the ribose or deoxyribose unit, and this linkage is always β.
  2. A five-carbon monosaccharide that is either d-ribose or deoxy-d-ribose.
  3. A phosphate ion.

     **The phosphate group** of a nucleotide is present as a phosphate ester and may be attached at C5′ or C3′.




A nucleotide




Adenylic acid


King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

## 6.4.1. The General Structure of Nucleic acids; Nucleotides and Nucleosides

### 6.4.1.1. Nucleotides

#### A) Heterocyclic Bases

- The nitrogenous base—either a purine or a pyrimidine—is linked to the 1′-carbon of the pentose sugar via replacement of the hydroxyl group.
- Uracil replaces thymine in an RNA nucleoside (or nucleotide).




King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

## 6.4.1. The General Structure of Nucleic acids; Nucleotides and Nucleosides


### 6.4.1.1. Nucleotides

#### B) A five-carbon monosaccharide

- The five-carbon monosaccharide in nucleic acids is either ribose or 2-deoxyribose, which are found in RNA and DNA, respectively.
- The key structural difference is at the C2′ carbon: ribose contains a hydroxyl (–OH) group, whereas in deoxyribose this group is replaced by hydrogen (–H).




D-ribose




2-deoxy-D-ribose


#### C) Phosphate group

- Phosphate groups link nucleotides through 3′–5′ phosphodiester bonds, forming the sugar–phosphate backbone of nucleic acids.
- The phosphate is attached to the 5′-carbon of one sugar and the 3′-hydroxyl of the next.

King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

## 6.4.1. The General Structure of Nucleic acids; Nucleotides and Nucleosides

### 6.4.1.2. Polynucleotides




COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT


## 6.4.1. The General Structure of Nucleic acids; Nucleotides and Nucleosides

### 6.4.1.3. Nucleosides

- Removal of the phosphate group of a nucleotide converts it to a compound known as **a nucleoside**.
- DNA-derived nucleosides consist of 2-deoxy-D-ribose and one of four bases: adenine, guanine, cytosine, or thymine.
- RNA nucleosides contain D-ribose and uracil instead of thymine, which is the main structural difference from DNA.




- RNA nucleosides contain D-ribose and uracil instead of thymine, which is the main structural difference from DNA.




King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

## 6.4.1. The General Structure of Nucleic acids; Nucleotides and Nucleosides

### 6.4.1.4. The structural difference between DNA and RNA


| | DNA | RNA |
|---|---|---|
| Sugar | 2-deoxy-D-ribose | D-ribose |
| A heterocyclic base | Adenine (A), Thymine (T), Guanine (G), Cytosine (C) | Adenine (A), Uracil (U), Guanine (G), Cytosine (C) |
| Number of strands | double-stranded (double helix) | single-stranded |



King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

## 6.4.1. The General Structure of Nucleic acids; Nucleotides and Nucleosides

### 6.4.1.5. DNA and RNA – Distinctions Summary



جامعة الملك سعود
King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT

# Practice Problems

King Saud University
COLLEGE OF SCIENCES
CHEMISTRY DEPARTMENT


## 6.5. Practice Problems

### A. Carbohydrates

**1** Classify the following carbohydrates:

| Compound | Classification |
|---|---|
| (a) Glucose | _____ |
| (b) Sucrose | _____ |
| (c) Starch | _____ |

**2** Identify the anomer (α or β) of the following D-glucopyranose:



A. α-Anomer B. β-Anomer
C. Both α and β D. Neither

**3** Which of the following is a ketose and a hexose?

A. Glucose C. Ribose
B. Fructose D. Glyceraldehyde

### B. Lipids


**1** What is the major product of complete hydrolysis of a triacylglycerol?

A. Glycerol and three fatty acids B. Three alcohols only
C. One fatty acid and two glycerols D. Three fatty acids only

**2** Classify the following fatty acids as saturated or unsaturated:

(a) $CH_3-(CH_2)_{16}-COOH$ → _____
(b) $CH_3-(CH_2)_7-CH=CH-(CH_2)_7-COOH$ → _____


**3** Identify the labeled parts of the triacylglycerol:




A. A = Glycerol backbone, B = Ester linkage
B. A = Fatty acid chain, B = Carbonyl group
C. A = Glycerol backbone, B = Fatty acid chain
D. A = Ester linkage, B = Glycerol backbone

### C. Amino Acids

**1** Identify the amino acid represented by the structure:




A. Glycine B. Alanine
C. Valine D. Serine

**2** Arrange the following amino acids by increasing side-chain hydrophobicity:

| Glycine (Gly) | Alanine (Ala) | Valine (Val) |
|---|---|---|

A. Val < Ala < Gly B. Gly < Ala < Val
C. Ala < Gly < Val D. Gly < Val < Ala


**3** What is the product of condensation between two amino acids?



A. Ester B. Peptide bond C. Glycosidic bond D. Disulfide bond

### D. Nucleic Acids

**1** Identify the labeled parts of the nucleotide:




A. A = Phosphate, B = Pentose (ribose), C = Adenine
B. A = Base, B = Deoxyribose, C = Phosphate
C. A = Phosphate, B = Base, C = Ribose
D. A = Pentose, B = Phosphate, C = Adenine

**2** Which base is found in DNA but not in RNA?

A. Uracil B. Thymine C. Cytosine D. Adenine

**3** Which of the following statements describes DNA and RNA?

(1) DNA: _____
_____
(2) RNA: _____
_____

A. (1) Usually single-stranded; (2) Contains thymine
B. (1) Double-stranded helix; (2) Contains uracil
C. (1) Contains uracil; (2) Contains thymine
D. (1) Ribose sugar; (2) Deoxyribose sugar