# Limits **Definitions**

Precise Definition : We say  $\lim_{x \to a} f(x) = L$  if for every  $\varepsilon > 0$  there is a  $\delta > 0$  such that whenever  $0 < |x - a| < \delta \text{ then } |f(x) - L| < \varepsilon.$ 

"Working" Definition : We say  $\lim_{x\to a}f(x)=L$  if we can make f(x) as close to L as we want by taking x sufficiently close to a (on either side of a) without letting x = a.

**Right hand limit :**  $\lim_{x \to a^+} f(x) = L$ . This has the same definition as the limit except it requires x > a.

**Left hand limit :**  $\lim_{x \to a} f(x) = L$ . This has the same There is a similar definition for  $\lim_{x \to a} f(x) = -\infty$ definition as the limit except it requires x < a.

**Limit at Infinity :** We say  $\lim_{x \to \infty} f(x) = L$  if we can make f(x) as close to L as we want by taking xlarge enough and positive.

There is a similar definition for  $\lim_{x \to -\infty} f(x) = L$ except we require x large and negative.

**Infinite Limit**: We say  $\lim f(x) = \infty$  if we can make f(x) arbitrarily large (and positive) by taking xsufficiently close to a (on either side of a) without letting x = a.

except we make f(x) arbitrarily large and negative.

## Relationship between the limit and one-sided limits

$$\lim_{x\to a} f(x) = L \quad \Rightarrow \quad \lim_{x\to a^+} f(x) = \lim_{x\to a^-} f(x) = L \qquad \qquad \lim_{x\to a^+} f(x) = \lim_{x\to a^-} f(x) = L \quad \Rightarrow \quad \lim_{x\to a} f(x) = L$$
 
$$\lim_{x\to a^+} f(x) \neq \lim_{x\to a^+} f(x) \Rightarrow \lim_{x\to a} f(x) \text{ Does Not Exist}$$

Assume  $\lim_{x \to a} f(x)$  and  $\lim_{x \to a} g(x)$  both exist and c is any number then,

1. 
$$\lim_{x \to a} [cf(x)] = c \lim_{x \to a} f(x)$$

2. 
$$\lim_{x \to a} [f(x) \pm g(x)] = \lim_{x \to a} f(x) \pm \lim_{x \to a} g(x)$$

3. 
$$\lim_{x \to a} [f(x)g(x)] = \lim_{x \to a} f(x) \lim_{x \to a} g(x)$$

$$\text{4.} \lim_{x \to a} \left[ \frac{f(x)}{g(x)} \right] = \lim_{x \to a} \frac{f(x)}{g(x)} \text{ provided } \lim_{x \to a} g(x) \neq 0$$

5. 
$$\lim_{x \to a} [f(x)]^n = \left[\lim_{x \to a} f(x)\right]^n$$

6. 
$$\lim_{x \to \infty} \left[ \sqrt[n]{f(x)} \right] = \sqrt[n]{\lim_{x \to \infty} f(x)}$$

## Basic Limit Evaluations at $\pm \infty$

1. 
$$\lim_{x \to \infty} \mathbf{e}^x = \infty$$
 &  $\lim_{x \to -\infty} \mathbf{e}^x = 0$ 

2. 
$$\lim_{x \to \infty} \ln(x) = \infty$$
 &  $\lim_{x \to 0^+} \ln(x) = -\infty$ 

3. If 
$$r > 0$$
 then  $\lim_{x \to \infty} \frac{b}{x^r} = 0$ 

4. If 
$$r>0$$
 and  $x^r$  is real for negative  $x$  then  $\lim_{x\to -\infty}\frac{b}{x^r}=0$ 

5. 
$$n \text{ even} : \lim_{x \to +\infty} x^n = \infty$$

6. 
$$n$$
 odd:  $\lim_{x \to \infty} x^n = \infty$  &  $\lim_{x \to -\infty} x^n = -\infty$ 

7. 
$$n \text{ even}: \lim_{x \to \pm \infty} a x^n + \dots + b x + c = \text{sgn}(a) \infty$$

8. 
$$n \text{ odd}$$
:  $\lim_{x \to \infty} a x^n + \dots + b x + c = \operatorname{sgn}(a) \infty$ 

9. 
$$n \text{ odd}$$
:  $\lim_{x \to -\infty} a x^n + \cdots + c x + d = -\operatorname{sgn}(a) \infty$ 

Note : 
$$sgn(a) = 1$$
 if  $a > 0$  and  $sgn(a) = -1$  if  $a < 0$ .

### **Evaluation Techniques**

#### Continuous Functions

If f(x) is continuous at a then  $\lim_{x\to a} f(x) = f(a)$ 

## **Continuous Functions and Composition**

$$f(x)$$
 is continuous at  $b$  and  $\lim_{x \to a} g(x) = b$  then 
$$\lim_{x \to a} f\left(g(x)\right) = f\left(\lim_{x \to a} g(x)\right) = f\left(b\right)$$

### **Factor and Cancel**

$$\lim_{x \to 2} \frac{x^2 + 4x - 12}{x^2 - 2x} = \lim_{x \to 2} \frac{(x - 2)(x + 6)}{x(x - 2)}$$
$$= \lim_{x \to 2} \frac{x + 6}{x} = \frac{8}{2} = 4$$

#### Rationalize Numerator/Denominator

$$\begin{split} &\lim_{x\to 9} \frac{3-\sqrt{x}}{x^2-81} = \lim_{x\to 9} \frac{3-\sqrt{x}}{x^2-81} \ \frac{3+\sqrt{x}}{3+\sqrt{x}} \\ &= \lim_{x\to 9} \frac{9-x}{(x^2-81)(3+\sqrt{x})} = \lim_{x\to 9} \frac{-1}{(x+9)(3+\sqrt{x})} \\ &= \frac{-1}{(18)(6)} = -\frac{1}{108} \end{split}$$

## Combine Rational Expressions

$$\begin{split} &\lim_{h\to 0}\frac{1}{h}\left(\frac{1}{x+h}-\frac{1}{x}\right)=\lim_{h\to 0}\frac{1}{h}\left(\frac{x-(x+h)}{x(x+h)}\right)\\ &=\lim_{h\to 0}\frac{1}{h}\left(\frac{-h}{x(x+h)}\right)=\lim_{h\to 0}\frac{-1}{x(x+h)}=-\frac{1}{x^2} \end{split}$$

## L'Hospital's/L'Hôpital's Rule

If 
$$\lim_{x \to a} \frac{f(x)}{g(x)} = \frac{0}{0}$$
 or  $\lim_{x \to a} \frac{f(x)}{g(x)} = \frac{\pm \infty}{\pm \infty}$  then,

$$\lim_{x\to a}\frac{f(x)}{g(x)}=\lim_{x\to a}\frac{f'(x)}{g'(x)}\text{, $a$ is a number, $\infty$ or $-\infty$}$$

## Polynomials at Infinity

p(x) and q(x) are polynomials. To compute  $\lim_{x\to\pm\infty}\frac{p(x)}{q(x)}$  factor largest power of x in q(x) out of

both 
$$p(\boldsymbol{x})$$
 and  $q(\boldsymbol{x})$  then compute limit.

$$\begin{split} \lim_{x \to -\infty} \frac{3x^2 - 4}{5x - 2x^2} &= \lim_{x \to -\infty} \frac{x^2 \left(3 - \frac{4}{x^2}\right)}{x^2 \left(\frac{5}{x} - 2\right)} \\ &= \lim_{x \to -\infty} \frac{3 - \frac{4}{x^2}}{\frac{5}{x} - 2} = -\frac{3}{2} \end{split}$$

$$\lim_{x\to -2}g(x) \text{ where } g(x) = \left\{ \begin{array}{ll} x^2+5 & \text{if } x<-2\\ 1-3x & \text{if } x\geq -2 \end{array} \right.$$

Compute two one sided limits.

$$\lim_{\substack{x\rightarrow -2^-\\\lim x\rightarrow -2^+}}g(x)=\lim_{\substack{x\rightarrow -2^-\\x\rightarrow -2^+}}x^2+5=9$$

One sided limits are different so  $\lim_{x\to a} g(x)$  doesn't exist. If the two one sided limits had been equal then  $\lim_{x \to 0} g(x)$  would have existed and had the same value.

## Some Continuous Functions

Partial list of continuous functions and the values of x for which they are continuous.

- 1. Polynomials for all x.
- 2. Rational function, except for x's that give division by zero.
- 3.  $\sqrt[n]{x}$  (n odd) for all x.
- 4.  $\sqrt[n]{x}$  (n even) for all  $x \ge 0$ .
- e<sup>x</sup> for all x.

- 6. ln(x) for x > 0.
- 7. cos(x) and sin(x) for all x.
- 8. tan(x) and sec(x) provided  $x \neq \cdots, -\frac{3\pi}{2}, -\frac{\pi}{2}, \frac{\pi}{2}, \frac{3\pi}{2}, \cdots$
- 9. cot(x) and csc(x) provided  $x \neq \cdots, -2\pi, -\pi, 0, \pi, 2\pi, \cdots$

#### Intermediate Value Theorem

Suppose that f(x) is continuous on [a,b] and let M be any number between f(a) and f(b). Then there exists a number c such that a < c < b and f(c) = M.