


Chapter 1

Chemistry: The Study of Change

Copyright © McGraw-Hill Education. All rights reserved. No reproduction or distribution without the prior written consent of McGraw-Hill Education.



©Science Source

Defining Chemistry

Chemistry is the study of matter and the changes it undergoes

Matter is anything that occupies space and has mass.

A **substance** is a form of matter that has a definite composition and distinct properties.



liquid nitrogen




gold ingots



silicon crystals

The Scientific Method

The ***scientific method*** is a systematic approach to research.



A ***hypothesis*** is a tentative explanation for a set of observations.

This hypothesis is then tested by developing and carrying out experiments. Based on the results of these experiments, the hypothesis is revised to account for experimental evidence. Further observations must be consistent with the hypothesis - otherwise the hypothesis must be revised.

The Scientific Method

A ***hypothesis*** is a tentative explanation for a set of observations.




A ***law*** is a concise statement of a relationship between phenomena that is always the same under the same conditions.



$$\text{Force} = \text{mass} \times \text{acceleration}$$

A ***theory*** is a unifying principle that explains a body of facts and/or those laws that are based on them.

Atomic Theory



International System of Units (SI)

Base Quantity	Name of Unit	Symbol
Length	meter	m
Mass	kilogram	kg
Time	second	s
Electrical current	ampere	A
Temperature	kelvin	K
Amount of substance	mole	mol
Luminous intensity	candela	cd

Prefixes Used with SI Units

Prefix	Symbol	Meaning	Example
peta-	P	1,000,000,000,000,000, or 10^{15}	1 petameter (Pm) = 1×10^{15} m
tera-	T	1,000,000,000,000, or 10^{12}	1 terameter (Tm) = 1×10^{12} m
giga-	G	1,000,000,000, or 10^9	1 gigameter (Gm) = 1×10^9 m
mega-	M	1,000,000, or 10^6	1 megameter (Mm) = 1×10^6 m
kilo-	K	1,000, or 10^3	1 kilometer (km) = 1×10^3 m
deci-	d	1/10, or 10^{-1}	1 decimeter (dm) = 0.1 m
centi-	c	1/100, or 10^{-2}	1 centimeter (cm) = 0.01 m
milli-	m	1/1,000, or 10^{-3}	1 millimeter (mm) = 0.001 m
micro-	μ	1/1,000,000, or 10^{-6}	1 micrometer (μ m) = 1×10^{-6} m
nano-	n	1/1,000,000,000, or 10^{-9}	1 nanometer (nm) = 1×10^{-9} m
pico-	p	1/1,000,000,000,000, or 10^{-12}	1 picometer (pm) = 1×10^{-12} m
femto-	f	1/1,000,000,000,000,000, or 10^{-15}	1 femtometer (fm) = 1×10^{-15} m
atto-	a	1/1,000,000,000,000,000,000, or 10^{-18}	1 attometer (am) = 1×10^{-18} m

Mixtures


A ***mixture*** is a combination of two or more substances in which the substances retain their distinct identities.

1. ***Homogenous mixture*** – composition of the mixture is the same throughout.

soft drink, milk, solder




2. ***Heterogeneous mixture*** – composition is not uniform throughout.



cement, iron filings in sand

Physical means can be used to separate a mixture into its pure components.



distillation



magnet

Elements

An ***element*** is a substance that cannot be separated into simpler substances by ***chemical means***.

- 118 elements have been identified
 - 82 elements occur naturally on Earth
gold, aluminum, lead, oxygen, carbon, sulfur



- 36 elements have been created by scientists
technetium, americium, seaborgium

TABLE 1.1 **Some Common Elements and Their Symbols**

Name	Symbol	Name	Symbol	Name	Symbol
Aluminum	Al	Fluorine	F	Oxygen	O
Arsenic	As	Gold	Au	Phosphorus	P
Barium	Ba	Hydrogen	H	Platinum	Pt
Bismuth	Bi	Iodine	I	Potassium	K
Bromine	Br	Iron	Fe	Silicon	Si
Calcium	Ca	Lead	Pb	Silver	Ag
Carbon	C	Magnesium	Mg	Sodium	Na
Chlorine	Cl	Manganese	Mn	Sulfur	S
Chromium	Cr	Mercury	Hg	Tin	Sn
Cobalt	Co	Nickel	Ni	Tungsten	W
Copper	Cu	Nitrogen	N	Zinc	Zn

Compounds

A ***compound*** is a substance composed of atoms of two or more elements chemically united in fixed proportions.

Compounds can only be separated into their pure components (elements) by ***chemical*** means.



lithium fluoride




quartz



dry ice
(carbon dioxide)



Classifications of Matter



Volume

Volume – SI derived unit for volume is cubic meter (m^3)

Volume: 1000 cm^3 ;
 1000 mL ;
 1 dm^3 ;
 1 L



$$1 \text{ cm}^3 = (1 \times 10^{-2} \text{ m})^3 = 1 \times 10^{-6} \text{ m}^3$$

$$1 \text{ dm}^3 = (1 \times 10^{-1} \text{ m})^3 = 1 \times 10^{-3} \text{ m}^3$$

$$1 \text{ L} = 1000 \text{ mL} = 1000 \text{ cm}^3 = 1 \text{ dm}^3$$

$$1 \text{ mL} = 1 \text{ cm}^3$$



Density

The **density** of a substance is its mass per unit volume (the volumetric mass).

SI derived unit for density is **kg/m³**

$$1 \text{ g/cm}^3 = 1 \text{ g/mL} = 1000 \text{ kg/m}^3$$

$$\text{density} = \frac{\text{mass}}{\text{volume}}$$

$$d = \frac{m}{V}$$

TABLE 1.4

Densities of Some Substances at 25°C

Substance	Density (g/cm³)
Air*	0.001
Ethanol	0.79
Water	1.00
Mercury	13.6
Table salt	2.2
Iron	7.9
Gold	19.3
Osmium [†]	22.6

*Measured at 1 atmosphere.

[†]Osmium (Os) is the densest element known.

A piece of platinum metal with a density of 21.5 g/cm³ has a volume of 4.49 cm³. What is its mass?

$$d = \frac{m}{V}$$

$$m = d \times V$$

$$= 21.5 \text{ g/cm}^3 \times 4.49 \text{ cm}^3 = \underline{\underline{96.5 \text{ g}}}$$

EXAMPLE 1.1

Gold is a precious metal that is chemically unreactive. It is used mainly in jewelry, dentistry, and electronic devices. A piece of gold ingot with a mass of 301 g has a volume of 15.6 cm³. Calculate the density of gold.

Solution We are given the mass and volume and asked to calculate the density. Therefore, from Equation (1.1), we write

$$\begin{aligned} d &= \frac{m}{V} \\ &= \frac{301 \text{ g}}{15.6 \text{ cm}^3} \\ &= 19.3 \text{ g/cm}^3 \end{aligned}$$

EXAMPLE 1.2


The density of mercury, the only metal that is a liquid at room temperature, is 13.6 g/mL. Calculate the mass of 5.50 mL of the liquid.

Solution We are given the density and volume of a liquid and asked to calculate the mass of the liquid. We rearrange Equation (1.1) to give

$$\begin{aligned} m &= d \times V \\ &= 13.6 \frac{\text{g}}{\text{mL}} \times 5.50 \text{ mL} \\ &= 74.8 \text{ g} \end{aligned}$$

Practice Exercise The density of sulfuric acid in a certain car battery is 1.41 g/mL. Calculate the mass of 242 mL of the liquid.

A Comparison of Temperature Scales



$$K = ^\circ C + 273.15$$

$$273 \text{ K} = 0 ^\circ C$$

$$373 \text{ K} = 100 ^\circ C$$

$$^\circ F = \frac{9}{5} \times ^\circ C + 32$$

$$32 ^\circ F = 0 ^\circ C$$

$$212 ^\circ F = 100 ^\circ C$$

Convert 172.9 °F to degrees Celsius.

$$^{\circ}\text{F} = \frac{9}{5} \times ^{\circ}\text{C} + 32$$

$$^{\circ}\text{F} - 32 = \frac{9}{5} \times ^{\circ}\text{C}$$

$$\frac{5}{9} \times (^{\circ}\text{F} - 32) = ^{\circ}\text{C}$$

$$^{\circ}\text{C} = \frac{5}{9} \times (^{\circ}\text{F} - 32)$$

$$^{\circ}\text{C} = \frac{5}{9} \times (172.9 - 32) = \underline{\underline{78.3}}$$

EXAMPLE 1.3

(a) Solder is an alloy made of tin and lead that is used in electronic circuits. A certain solder has a melting point of 224°C . What is its melting point in degrees Fahrenheit?
(b) Helium has the lowest boiling point of all the elements at -452°F . Convert this temperature to degrees Celsius.
(c) Mercury, the only metal that exists as a liquid at room temperature, melts at -38.9°C . Convert its melting point to kelvins.

Solution These three parts require that we carry out temperature conversions, so we need Equations (1.2), (1.3), and (1.4). Keep in mind that the lowest temperature on the Kelvin scale is zero (0 K); therefore, it can never be negative.

(a) This conversion is carried out by writing

$$\frac{9^{\circ}\text{F}}{5^{\circ}\text{C}} \times (224^{\circ}\text{C}) + 32^{\circ}\text{F} = 435^{\circ}\text{F}$$

Scientific Notation

- The number of atoms in 12 g of carbon:

602,200,000,000,000,000,000,000,000

$$6.022 \times 10^{23}$$

- The mass of a single carbon atom in grams:

[illegible]

$$1.99 \times 10^{-23}$$

$$N \times 10^n$$

N is a number
between 1 and 10

n is a positive or negative integer

Scientific Notation

568.762

← move decimal left

$$n > 0$$

$$568.762 = 5.68762 \times 10^2$$

0.00000772

→ move decimal right

$$n < 0$$

$$0.00000772 = 7.72 \times 10^{-6}$$

- Addition or Subtraction

1. Write each quantity with the same exponent ***n***
2. Combine N_1 and N_2
3. The exponent, ***n***, remains the same

$$\begin{aligned} 4.31 \times 10^4 + 3.9 \times 10^3 &= \\ 4.31 \times 10^4 + 0.39 \times 10^4 &= \\ 4.70 \times 10^4 & \end{aligned}$$

Scientific Notation

Multiplication

1. Multiply N_1 and N_2
2. Add exponents n_1 and n_2

$$\begin{aligned}(4.0 \times 10^{-5}) \times (7.0 \times 10^3) &= \\ (4.0 \times 7.0) \times (10^{-5+3}) &= \\ 28 \times 10^{-2} &= \\ 2.8 \times 10^{-1}\end{aligned}$$

Division

1. Divide N_1 and N_2
2. Subtract exponents n_1 and n_2

$$\begin{aligned}8.5 \times 10^4 \div 5.0 \times 10^9 &= \\ (8.5 \div 5.0) \times 10^{4-9} &= \\ 1.7 \times 10^{-5}\end{aligned}$$

Significant Figures

- Any digit that is not zero is significant
1.234 kg **4** significant figures
- Zeros between nonzero digits are significant
606 m **3** significant figures
- Zeros to the left of the first nonzero digit are **not** significant
0.08 L **1** significant figure
- If a number is greater than 1, then all zeros to the right of the decimal point are significant
2.0 mg **2** significant figures
- If a number is less than 1, then only the zeros that are at the end and in the middle of the number are significant
0.00420 g **3** significant figures

(1) Express 568.762 in scientific notation:

$$568.762 = 5.68762 \times 10^2$$

Note that the decimal point is moved to the left by two places and $n = 2$

(2) Express 0.00000772 in scientific notation:

$$0.00000772 = 7.72 \times 10^{-6}$$

Here the decimal point is moved to the right by six places and $n = -6$

EXAMPLE 1.4

Determine the number of significant figures in the following measurements: (a) 478 cm, (b) 6.01 g, (c) 0.825 m, (d) 0.043 kg, (e) 1.310×10^{22} atoms, (f) 7000 mL.

Solution (a) Three, because each digit is a nonzero digit. (b) Three, because zeros between nonzero digits are significant. (c) Three, because zeros to the left of the first nonzero digit do not count as significant figures. (d) Two. Same reason as in (c). (e) Four, because the number is greater than one so all the zeros written to the right of the decimal point count as significant figures. (f) This is an ambiguous case. The number of significant figures may be four (7.000×10^3), three (7.00×10^3), two (7.0×10^3), or one (7×10^3). This example illustrates why scientific notation must be used to show the proper number of significant figures.

Significant Figures ₂

- Addition or Subtraction
- The answer cannot have more digits to the right of the decimal point than any of the original numbers.

89.332

+1.1 ← one significant figure after decimal point

90.432 ← roundoff to 90.4

3.70

-2.9133 ← two significant figure after decimal point

0.7867 ← roundoff to 0.79

Significant Figures ₃

- Multiplication or Division
- The number of significant figures in the result is set by the original number that has the ***smallest*** number of significant figures.

$$\begin{array}{ccc} 4.51 & \times 3.6666 & = 16.536366 = 16.5 \\ \uparrow & & \uparrow \\ 3 \text{ sig figs} & & \text{round to} \\ & & 3 \text{ sig figs} \end{array}$$

$$\begin{array}{ccc} 6.8 & \div 112.04 & = 0.0606926 = 0.061 \\ \uparrow & & \uparrow \\ 2 \text{ sig figs} & & \text{round to} \\ & & 2 \text{ sig figs} \end{array}$$

Significant Figures ₄

Exact Numbers

Numbers from definitions or numbers of objects are considered to have an infinite number of significant figures.

The average of three measured lengths: 6.64, 6.68 and 6.70?

$$\frac{6.64 + 6.68 + 6.70}{3} = 6.67333 = 6.67$$

Because 3 is an **exact number**

EXAMPLE 1.5

Carry out the following arithmetic operations to the correct number of significant figures: (a) $11,254.1 \text{ g} + 0.1983 \text{ g}$, (b) $66.59 \text{ L} - 3.113 \text{ L}$, (c) $8.16 \text{ m} \times 5.1355$, (d) $0.0154 \text{ kg} \div 88.3 \text{ mL}$, (e) $2.64 \times 10^3 \text{ cm} + 3.27 \times 10^2 \text{ cm}$.

Solution In addition and subtraction, the number of decimal places in the answer is determined by the number having the lowest number of decimal places. In multiplication and division, the significant number of the answer is determined by the number having the smallest number of significant figures.

$$\begin{array}{r} \text{(a)} \quad 11,254.1 \text{ g} \\ + \quad 0.1983 \text{ g} \\ \hline 11,254.2983 \text{ g} \leftarrow \text{round off to } 11,254.3 \text{ g} \end{array}$$

$$\begin{array}{r} \text{(b)} \quad 66.59 \text{ L} \\ - \quad 3.113 \text{ L} \\ \hline 63.477 \text{ L} \leftarrow \text{round off to } 63.48 \text{ L} \end{array}$$

$$\text{(c)} \quad 8.16 \text{ m} \times 5.1355 = 41.90568 \text{ m} \leftarrow \text{round off to } 41.9 \text{ m}$$

$$\begin{array}{l} \text{(d)} \quad \frac{0.0154 \text{ kg}}{88.3 \text{ mL}} = 0.000174405436 \text{ kg/mL} \leftarrow \text{round off to } 0.000174 \text{ kg/mL} \\ \text{or } 1.74 \times 10^{-4} \text{ kg/mL} \end{array}$$

$$\begin{array}{l} \text{(e)} \quad \text{First we change } 3.27 \times 10^2 \text{ cm to } 0.327 \times 10^3 \text{ cm and then carry out the addition} \\ (2.64 \text{ cm} + 0.327 \text{ cm}) \times 10^3. \text{ Following the procedure in (a), we find the answer is} \\ 2.97 \times 10^3 \text{ cm.} \end{array}$$

Dimensional Analysis Method of Solving Problems

1. Determine which unit conversion factor(s) are needed
2. Carry units through calculation
3. If all units cancel except for the ***desired unit(s)***, then the problem was solved correctly.

given quantity $\times$ conversion factor = desired quantity

$$\cancel{\text{given unit}} \times \frac{\text{desired unit}}{\cancel{\text{given unit}}} = \text{desired unit}$$

EXAMPLE 1.6

A person's average daily intake of glucose (a form of sugar) is 0.0833 pound (lb). What is this mass in milligrams (mg)? (1 lb = 453.6 g.)

Strategy The problem can be stated as

$$? \text{ mg} = 0.0833 \text{ lb}$$

The relationship between pounds and grams is given in the problem. This relationship will enable conversion from pounds to grams. A metric conversion is then needed to convert grams to milligrams ($1 \text{ mg} = 1 \times 10^{-3} \text{ g}$). Arrange the appropriate conversion factors so that pounds and grams cancel and the unit milligrams is obtained in your answer.

Solution The sequence of conversions is

pounds $\longrightarrow$ grams $\longrightarrow$ milligrams

Using the following conversion factors

$$\frac{453.6 \text{ g}}{1 \text{ lb}} \quad \text{and} \quad \frac{1 \text{ mg}}{1 \times 10^{-3} \text{ g}}$$

we obtain the answer in one step:

$$? \text{ mg} = 0.0833 \text{ lb} \times \frac{453.6 \text{ g}}{1 \text{ lb}} \times \frac{1 \text{ mg}}{1 \times 10^{-3} \text{ g}} = 3.78 \times 10^4 \text{ mg}$$

EXAMPLE 1.7

An average adult has 5.2 L of blood. What is the volume of blood in m^3 ?

Strategy The problem can be stated as

$$? \text{ m}^3 = 5.2 \text{ L}$$

How many conversion factors are needed for this problem? Recall that $1 \text{ L} = 1000 \text{ cm}^3$ and $1 \text{ cm} = 1 \times 10^{-2} \text{ m}$.

Solution We need two conversion factors here: one to convert liters to cm^3 and one to convert centimeters to meters:

$$\frac{1000 \text{ cm}^3}{1 \text{ L}} \quad \text{and} \quad \frac{1 \times 10^{-2} \text{ m}}{1 \text{ cm}}$$

Because the second conversion factor deals with length (cm and m) and we want volume here, it must therefore be cubed to give

$$\frac{1 \times 10^{-2} \text{ m}}{1 \text{ cm}} \times \frac{1 \times 10^{-2} \text{ m}}{1 \text{ cm}} \times \frac{1 \times 10^{-2} \text{ m}}{1 \text{ cm}} = \left(\frac{1 \times 10^{-2} \text{ m}}{1 \text{ cm}} \right)^3$$

This means that $1 \text{ cm}^3 = 1 \times 10^{-6} \text{ m}^3$. Now we can write

$$? \text{ m}^3 = 5.2 \text{ L} \times \frac{1000 \text{ cm}^3}{1 \text{ L}} \times \left(\frac{1 \times 10^{-2} \text{ m}}{1 \text{ cm}} \right)^3 = 5.2 \times 10^{-3} \text{ m}^3$$