



**Answer the following questions:**

**Q1: [4+5]**

(a) For the Markov process  $\{X_t\}$ ,  $t=0,1,2,\dots,n$  with states  $i_0, i_1, i_2, \dots, i_{n-1}, i_n$

Prove that:  $\Pr\{X_0 = i_0, X_1 = i_1, X_2 = i_2, \dots, X_n = i_n\} = p_{i_0} P_{i_0 i_1} P_{i_1 i_2} \dots P_{i_{n-1} i_n}$  where  $p_{i_0} = \Pr\{X_0 = i_0\}$

(b) A Markov chain  $X_0, X_1, X_2, \dots$  has the transition probability matrix

$$\mathbf{P} = \begin{matrix} & \begin{matrix} 0 & 1 & 2 \end{matrix} \\ \begin{matrix} 0 \\ 1 \\ 2 \end{matrix} & \left\| \begin{array}{ccc} 0.2 & 0.3 & 0.5 \\ 0.4 & 0.5 & 0.1 \\ 0.3 & 0.2 & 0.5 \end{array} \right\| \end{matrix}$$

and initial distribution  $p_0 = 0.3$  and  $p_1 = 0.7$ . Determine the following probabilities

i)  $\Pr\{X_0 = 1, X_1 = 1, X_2 = 0\}$

ii)  $\Pr\{X_2 = 0\}$

**Q2: [5+4]**

(a) Consider a spare parts inventory model in which either 0, 1, or 2 repair parts are demanded in any period, with  $\Pr\{\xi_n = 0\} = 0.3$ ,  $\Pr\{\xi_n = 1\} = 0.2$ ,  $\Pr\{\xi_n = 2\} = 0.5$  and suppose  $s=0$  and  $S=3$ . Determine the transition probability matrix for the Markov chain  $\{X_n\}$ , where  $X_n$  is defined to be the quantity on hand at the end of period  $n$ .

(b) Let  $X_n$  denote the quality of the  $n$ th item that produced in a certain factory with  $X_n = 0$  meaning “good” and  $X_n = 1$  meaning “defective”. Suppose that  $\{X_n\}$  be a Markov chain whose transition matrix is

$$P = \begin{matrix} & \begin{matrix} 0 & 1 \end{matrix} \\ \begin{matrix} 0 \\ 1 \end{matrix} & \left\| \begin{matrix} 0.99 & 0.01 \\ 0.12 & 0.88 \end{matrix} \right\| \end{matrix}$$

- i) What is the probability that the fourth item is defective given that the first item is good?
- ii) In the long run, what is the probability that an item produced by this system is good?

**Q3: [7]**

Consider the Markov chain whose transition probability matrix is given by

$$P = \begin{matrix} & \begin{matrix} 0 & 1 & 2 & 3 \end{matrix} \\ \begin{matrix} 0 \\ 1 \\ 2 \\ 3 \end{matrix} & \left\| \begin{matrix} 1 & 0 & 0 & 0 \\ 0.2 & 0.4 & 0.3 & 0.1 \\ 0.1 & 0.5 & 0.3 & 0.1 \\ 0 & 0 & 0 & 1 \end{matrix} \right\| \end{matrix}$$

- (i) Starting in state 1, determine the probability that the Markov chain ends in state 0.
- (ii) Determine the mean time to absorption.
- (iii) Sketch, the Markov chain diagram, and determine whether it's an absorbing chain or not.



## The Model Answer

### Q1: [4+5]

(a)

$$\begin{aligned} & \because \Pr\{X_0 = i_0, X_1 = i_1, X_2 = i_2, \dots, X_n = i_n\} \\ &= \Pr\{X_0 = i_0, X_1 = i_1, X_2 = i_2, \dots, X_{n-1} = i_{n-1}\} \cdot \Pr\{X_n = i_n | X_0 = i_0, X_1 = i_1, X_2 = i_2, \dots, X_{n-1} = i_{n-1}\} \\ &= \Pr\{X_0 = i_0, X_1 = i_1, X_2 = i_2, \dots, X_{n-1} = i_{n-1}\} \cdot P_{i_{n-1}i_n} \quad \text{Definition of Markov} \end{aligned}$$

By repeating this argument  $n - 1$  times

$$\begin{aligned} & \therefore \Pr\{X_0 = i_0, X_1 = i_1, X_2 = i_2, \dots, X_n = i_n\} \\ &= p_{i_0} P_{i_0i_1} P_{i_1i_2} \dots P_{i_{n-2}i_{n-1}} P_{i_{n-1}i_n} \quad \text{where } p_{i_0} = \Pr\{X_0 = i_0\} \text{ is obtained from the initial distribution of the process.} \end{aligned}$$

(b)

$$\begin{aligned} \text{i) } pr\{X_0 = 1, X_1 = 1, X_2 = 0\} &= p_1 P_{11} P_{10} \quad , \quad p_1 = pr\{X_0 = 1\} \\ &= 0.7(0.5)(0.4) \\ &= 0.14 \end{aligned}$$

$$\begin{aligned} \text{ii) } \because pr\{X_2 = 0\} &= pr\{X_2 = 0 | X_0 = 0\} pr\{X_0 = 0\} \\ &+ pr\{X_2 = 0 | X_0 = 1\} pr\{X_0 = 1\} \\ &= P_{00}^2 P_0 + P_{10}^2 P_1 \quad , \quad P_0 = 0.3, \quad P_1 = 0.7 \end{aligned}$$

$$\begin{aligned} \text{and } P^2 &= \begin{bmatrix} 0.2 & 0.3 & 0.5 \\ 0.4 & 0.5 & 0.1 \\ 0.3 & 0.2 & 0.5 \end{bmatrix} \begin{bmatrix} 0.2 & 0.3 & 0.5 \\ 0.4 & 0.5 & 0.1 \\ 0.3 & 0.2 & 0.5 \end{bmatrix} \\ &= \begin{bmatrix} 0.31 & 0.31 & 0.38 \\ 0.31 & 0.39 & 0.30 \\ 0.29 & 0.29 & 0.42 \end{bmatrix} \end{aligned}$$

$$\begin{aligned} \therefore pr\{X_2 = 0\} &= 0.31(0.3) + 0.31(0.7) \\ &= 0.31 \end{aligned}$$

### Q2: [5+4]

(a)

$$\begin{array}{c}
 -1 \quad 0 \quad 1 \quad 2 \quad 3 \\
 -1 \left\| \begin{array}{ccccc} 0 & 0 & 0.5 & 0.2 & 0.3 \\ 0 & 0 & 0.5 & 0.2 & 0.3 \\ 1 & 0.5 & 0.2 & 0.3 & 0 & 0 \\ 2 & 0 & 0.5 & 0.2 & 0.3 & 0 \\ 3 & 0 & 0 & 0.5 & 0.2 & 0.3 \end{array} \right\|
 \end{array}$$

$P_{ij} = \Pr(\xi_{n+1} = S - j)$  ,  $i \leq s$  for replenishment

$P_{-1,-1} = \Pr(\xi_{n+1} = 4) = 0$  ,  $P_{01} = \Pr(\xi_{n+1} = 2) = 0.5$

$P_{ij} = \Pr(\xi_{n+1} = i - j)$  ,  $s < i \leq S$  for non-replenishment

$P_{1,-1} = \Pr(\xi_{n+1} = 2) = 0.5$  ,  $P_{11} = \Pr(\xi_{n+1} = 0) = 0.3$ ,  $P_{21} = \Pr(\xi_{n+1} = 1) = 0.2$

(b)

i)

$$P^3 = \begin{bmatrix} 0.9737 & 0.0263 \\ 0.3152 & 0.6848 \end{bmatrix}$$

$$pr\{X_3 = 1 | X_0 = 0\} = p_{01}^3 = 0.0263$$

ii)

In the long run, the probability that an item produced by this system is good is given by:

$$\begin{aligned}
 b / (a + b) &= \frac{0.12}{0.01 + 0.12} \\
 &= \frac{12}{13} = 92.31 \% ,
 \end{aligned}$$

$$\text{where } \lim_{n \rightarrow \infty} P^n = \begin{bmatrix} \frac{b}{a+b} & \frac{a}{a+b} \\ \frac{b}{a+b} & \frac{a}{a+b} \end{bmatrix}$$

Q3: [7]

$$\mathbf{P} = \begin{array}{c}
 \quad \quad \quad 0 \quad 1 \quad 2 \quad 3 \\
 \begin{array}{c} 0 \\ 1 \\ 2 \\ 3 \end{array} \left\| \begin{array}{cccc} 1 & 0 & 0 & 0 \\ 0.2 & 0.4 & 0.3 & 0.1 \\ 0.1 & 0.5 & 0.3 & 0.1 \\ 0 & 0 & 0 & 1 \end{array} \right\|
 \end{array}$$

$$u_i = pr\{X_T = 0 | X_0 = i\} \text{ for } i=1,2,$$

$$\text{and } v_i = E[T | X_0 = i] \text{ for } i=1,2.$$

(i)

$$u_1 = p_{10} + p_{11}u_1 + p_{12}u_2$$

$$u_2 = p_{20} + p_{21}u_1 + p_{22}u_2$$

$\Rightarrow$

$$u_1 = 0.2 + 0.4u_1 + 0.3u_2$$

$$u_2 = 0.1 + 0.5u_1 + 0.3u_2$$

$\Rightarrow$

$$6u_1 - 3u_2 = 2 \quad (1)$$

$$5u_1 - 7u_2 = -1 \quad (2)$$

Solving (1) and (2), we get

$$u_1 = \frac{17}{27} \text{ and } u_2 = \frac{16}{27}$$

Starting in state 1, the probability that the Markov chain ends in state 0 is

$$u_1 = u_{10} = \frac{17}{27} = 0.6296$$

(ii) Also, the mean time to absorption can be found as follows

$$v_1 = 1 + p_{11}v_1 + p_{12}v_2$$

$$v_2 = 1 + p_{21}v_1 + p_{22}v_2$$

$\Rightarrow$

$$v_1 = 1 + 0.4v_1 + 0.3v_2$$

$$v_2 = 1 + 0.5v_1 + 0.3v_2$$

$\Rightarrow$


$$6v_1 - 3v_2 = 10 \quad (1)$$

$$5v_1 - 7v_2 = -10 \quad (2)$$

Solving (1) and (2), we get

$$v_1 = v_{10} = \frac{100}{27} \approx 3.7$$

(iii) It's an absorbing Markov Chain.



Markov Chain Diagram