

Chapter 1

Systems of Linear Equations

1.1 Introduction to Systems of Linear Equations

1.2 Gaussian Elimination and Gauss-Jordan Elimination

1.1 Introduction to Systems of Linear Equations

- a linear equation in n variables:

$$a_1x_1 + a_2x_2 + a_3x_3 + \dots + a_nx_n = b$$

$a_1, a_2, a_3, \dots, a_n, b$: real number

a_1 : leading coefficient

x_1 : leading variable

- Notes:

- (1) Linear equations have no products or roots of variables and no variables involved in trigonometric, exponential, or logarithmic functions.
- (2) Variables appear only to the first power.

■ Ex 1: (Linear or Nonlinear)

Linear (a) $3x + 2y = 7$

(b) $\frac{1}{2}x + y - \pi z = \sqrt{2}$ Linear

Linear (c) $x_1 - 2x_2 + 10x_3 + x_4 = 0$ (d) $(\sin \frac{\pi}{2})x_1 - 4x_2 = e^2$ Linear

Nonlinear (e) $xy + z = 2$ (f) $e^x - 2y = 4$ Nonlinear

not the first power *Exponential*

Nonlinear (g) $\sin x_1 + 2x_2 - 3x_3 = 0$ (h) $\frac{1}{x} + \frac{1}{y} = 4$ Nonlinear

trigonometric functions *not the first power*

-
- a solution of a linear equation in n variables:

$$a_1x_1 + a_2x_2 + a_3x_3 + \cdots + a_nx_n = b$$

$$x_1 = s_1, x_2 = s_2, x_3 = s_3, \cdots, x_n = s_n$$

such that $a_1s_1 + a_2s_2 + a_3s_3 + \cdots + a_ns_n = b$

- **Solution set:**

the set of all solutions of a linear equation

- Ex 2 : (Parametric representation of a solution set)

$$x_1 + 2x_2 = 4$$

a solution: $(2, 1)$, i.e. $x_1 = 2, x_2 = 1$

If you solve for x_1 in terms of x_2 , you obtain

$$x_1 = 4 - 2x_2,$$

By letting $x_2 = t$ you can represent the solution set as

$$x_1 = 4 - 2t$$

And the solutions are $\{(4 - 2t, t) \mid t \in R\}$ or $\{(s, 2 - \frac{1}{2}s) \mid s \in R\}$

-
- a system of m linear equations in n variables:

$$a_{11}x_1 + a_{12}x_2 + a_{13}x_3 + \cdots + a_{1n}x_n = b_1$$

$$a_{21}x_1 + a_{22}x_2 + a_{23}x_3 + \cdots + a_{2n}x_n = b_2$$

$$a_{31}x_1 + a_{32}x_2 + a_{33}x_3 + \cdots + a_{3n}x_n = b_3$$

$$\vdots$$

$$a_{m1}x_1 + a_{m2}x_2 + a_{m3}x_3 + \cdots + a_{mn}x_n = b_m$$

- **Consistent:**

A system of linear equations has at least one solution.

- **Inconsistent:**

A system of linear equations has no solution.

- Notes:

Every system of linear equations has either


(1) exactly one solution,

(2) infinitely many solutions, or

(3) no solution.


■ Ex 4: (Solution of a system of linear equations)

(1) $x + y = 3$
 $x - y = -1$
two intersecting lines




exactly one solution

(2) $x + y = 3$
 $2x + 2y = 6$
two coincident lines



infinite number

(3) $x + y = 3$
 $x + y = 1$
two parallel lines



no solution

-
- **Ex 5:** (Using back substitution to solve a system in row echelon form)

$$x - 2y = 5 \quad (1)$$

$$y = -2 \quad (2)$$

Sol: By substituting $y = -2$ into (1), you obtain

$$x - 2(-2) = 5$$

$$x = 1$$

The system has exactly one solution: $x = 1, y = -2$

-
- Ex 6: (Using back substitution to solve a system in row echelon form)

$$x - 2y + 3z = 9 \quad (1)$$

$$y + 3z = 5 \quad (2)$$

$$z = 2 \quad (3)$$

Sol: Substitute $z = 2$ into (2)

$$y + 3(2) = 5$$

$$y = -1$$

and substitute $y = -1$ and $z = 2$ into (1)

$$x - 2(-1) + 3(2) = 9$$

$$x = 1$$

The system has exactly one solution:

$$x = 1, y = -1, z = 2$$

- **Equivalent:**

Two systems of linear equations are called **equivalent** if they have precisely the same solution set.

- **Notes:**

Each of the following operations on a system of linear equations produces an equivalent system.

(1) Interchange two equations.

(2) Multiply an equation by a nonzero constant.

(3) Add a multiple of an equation to another equation.

-
- Ex 7: Solve a system of linear equations (consistent system)

$$x - 2y + 3z = 9 \quad (1)$$

$$-x + 3y = -4 \quad (2)$$

$$2x - 5y + 5z = 17 \quad (3)$$

Sol: $(1) + (2) \rightarrow (2)$

$$\begin{array}{rcl} x - 2y + 3z & = & 9 \\ & y + 3z & = 5 \end{array} \quad (4)$$

$$2x - 5y + 5z = 17$$

$(1) \times (-2) + (3) \rightarrow (3)$

$$\begin{array}{rcl} x - 2y + 3z & = & 9 \\ & y + 3z & = 5 \\ & -y - z & = -1 \end{array} \quad (5)$$

$$(4) + (5) \rightarrow (5)$$

$$\begin{array}{rclcl} x & - & 2y & + & 3z & = & 9 \\ & & y & + & 3z & = & 5 \\ & & & & 2z & = & 4 \end{array} \quad (6)$$

$$(6) \times \frac{1}{2} \rightarrow (6)$$

$$\begin{array}{rclcl} x & - & 2y & + & 3z & = & 9 \\ & & y & + & 3z & = & 5 \\ & & & & z & = & 2 \end{array}$$

So the solution is $x = 1$, $y = -1$, $z = 2$ (only one solution)

-
- Ex 8: Solve a system of linear equations (inconsistent system)

$$x_1 - 3x_2 + x_3 = 1 \quad (1)$$

$$2x_1 - x_2 - 2x_3 = 2 \quad (2)$$

$$x_1 + 2x_2 - 3x_3 = -1 \quad (3)$$

Sol: $(1) \times (-2) + (2) \rightarrow (2)$

$(1) \times (-1) + (3) \rightarrow (3)$

$$\begin{array}{rclcl} x_1 & - & 3x_2 & + & x_3 & = & 1 \\ & & 5x_2 & - & 4x_3 & = & 0 \end{array} \quad (4)$$

$$\begin{array}{rclcl} & & 5x_2 & - & 4x_3 & = & -2 \end{array} \quad (5)$$

$$(4) \times (-1) + (5) \rightarrow (5)$$

$$x_1 - 3x_2 + x_3 = 1$$

$$5x_2 - 4x_3 = 0$$

$$\boxed{0 = -2} \text{ (a false statement)}$$

So the system has no solution (an inconsistent system).

-
- Ex 9: Solve a system of linear equations (infinitely many solutions)

$$x_2 - x_3 = 0 \quad (1)$$

$$x_1 - 3x_3 = -1 \quad (2)$$

$$-x_1 + 3x_2 = 1 \quad (3)$$

Sol: $(1) \leftrightarrow (2)$

$$x_1 - 3x_3 = -1 \quad (1)$$

$$x_2 - x_3 = 0 \quad (2)$$

$$-x_1 + 3x_2 = 1 \quad (3)$$

$(1) + (3) \rightarrow (3)$

$$x_1 - 3x_3 = -1$$

$$x_2 - x_3 = 0$$

$$3x_2 - 3x_3 = 0 \quad (4)$$

$$\begin{array}{rclcl} x_1 & & - & 3x_3 & = & -1 \\ & x_2 & - & x_3 & = & 0 \end{array}$$

$$\Rightarrow x_2 = x_3, \quad x_1 = -1 + 3x_3$$

$$\text{let } x_3 = t$$

$$\text{then } x_1 = 3t - 1,$$

$$x_2 = t, \quad t \in \mathbb{R}$$

$$x_3 = t,$$

So this system has infinitely many solutions.

1.2 Gaussian Elimination and Gauss-Jordan Elimination

- $m \times n$ matrix:

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \cdots & a_{2n} \\ a_{31} & a_{32} & a_{33} & \cdots & a_{3n} \\ \vdots & & & & \\ a_{m1} & a_{m2} & a_{m3} & \cdots & a_{mn} \end{bmatrix} \quad \begin{array}{l} m \text{ rows} \\ n \text{ columns} \end{array}$$

- Notes:

- (1) Every entry a_{ij} in a matrix is a number.
- (2) A matrix with m rows and n columns is said to be of size $m \times n$.
- (3) If $m = n$, then the matrix is called square of order n .
- (4) For a square matrix, the entries $a_{11}, a_{22}, \dots, a_{nn}$ are called the main diagonal entries.

■ Ex 1:	Matrix	Size
	$[2]$	1×1
	$\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$	2×2
	$\left[1 \quad -3 \quad 0 \quad \frac{1}{2} \right]$	1×4
	$\begin{bmatrix} e & \pi \\ 2 & \sqrt{2} \\ -7 & 4 \end{bmatrix}$	3×2

■ Note:

One very common use of matrices is to represent a system of linear equations.

-
- a system of m equations in n variables:

$$a_{11}x_1 + a_{12}x_2 + a_{13}x_3 + \dots + a_{1n}x_n = b_1$$

$$a_{21}x_1 + a_{22}x_2 + a_{23}x_3 + \dots + a_{2n}x_n = b_2$$

$$a_{31}x_1 + a_{32}x_2 + a_{33}x_3 + \dots + a_{3n}x_n = b_3$$

$$\vdots$$

$$a_{m1}x_1 + a_{m2}x_2 + a_{m3}x_3 + \dots + a_{mn}x_n = b_m$$

Matrix form:

$$Ax = b$$

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \cdots & a_{2n} \\ a_{31} & a_{32} & a_{33} & \cdots & a_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & a_{m3} & \cdots & a_{mn} \end{bmatrix} \quad x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \quad b = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix}$$

- Augmented matrix:

$$\left[\begin{array}{ccccc|c} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} & b_1 \\ a_{21} & a_{22} & a_{23} & \cdots & a_{2n} & b_2 \\ a_{31} & a_{32} & a_{33} & \cdots & a_{3n} & b_3 \\ & \vdots & & & & \\ a_{m1} & a_{m2} & a_{m3} & \cdots & a_{mn} & b_m \end{array} \right] = [A \mid b]$$

- Coefficient matrix:

$$\left[\begin{array}{ccccc} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \cdots & a_{2n} \\ a_{31} & a_{32} & a_{33} & \cdots & a_{3n} \\ & \vdots & & & \\ a_{m1} & a_{m2} & a_{m3} & \cdots & a_{mn} \end{array} \right] = A$$

- Elementary row operation:

(1) Interchange two rows.

$$r_{ij} : R_i \leftrightarrow R_j$$

(2) Multiply a row by a nonzero constant.

$$r_i^{(k)} : (k)R_i \rightarrow R_i$$

(3) Add a multiple of a row to another row.

$$r_{ij}^{(k)} : (k)R_i + R_j \rightarrow R_j$$

- Row equivalent:

Two matrices are said to be **row equivalent** if one can be obtained from the other by a finite sequence of elementary row operation.

■ Ex 2: (Elementary row operation)

$$\begin{bmatrix} 0 & 1 & 3 & 4 \\ -1 & 2 & 0 & 3 \\ 2 & -3 & 4 & 1 \end{bmatrix} \xrightarrow{r_{12}} \begin{bmatrix} -1 & 2 & 0 & 3 \\ 0 & 1 & 3 & 4 \\ 2 & -3 & 4 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 2 & -4 & 6 & -2 \\ 1 & 3 & -3 & 0 \\ 5 & -2 & 1 & 2 \end{bmatrix} \xrightarrow{r_1^{(\frac{1}{2})}} \begin{bmatrix} 1 & -2 & 3 & -1 \\ 1 & 3 & -3 & 0 \\ 5 & -2 & 1 & 2 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 2 & -4 & 3 \\ 0 & 3 & -2 & -1 \\ 2 & 1 & 5 & -2 \end{bmatrix} \xrightarrow{r_{13}^{(-2)}} \begin{bmatrix} 1 & 2 & -4 & 3 \\ 0 & 3 & -2 & -1 \\ 0 & -3 & 13 & -8 \end{bmatrix}$$

■ Ex 3: Using elementary row operations to solve a system

Linear System

Associated
Augmented Matrix

Elementary
Row Operation

$$\begin{array}{rrcr} x & - & 2y & + & 3z & = & 9 \\ -x & + & 3y & & & = & -4 \\ 2x & - & 5y & + & 5z & = & 17 \end{array} \quad \left[\begin{array}{cccc} 1 & -2 & 3 & 9 \\ -1 & 3 & 0 & -4 \\ 2 & -5 & 5 & 17 \end{array} \right]$$

$$\begin{array}{rrcr} x & - & 2y & + & 3z & = & 9 \\ & & y & + & 3z & = & 5 \\ 2x & - & 5y & + & 5z & = & 17 \end{array} \quad \left[\begin{array}{cccc} 1 & -2 & 3 & 9 \\ 0 & 1 & 3 & 5 \\ 2 & -5 & 5 & 17 \end{array} \right]$$

$$r_{12}^{(1)} : (1)R_1 + R_2 \rightarrow R_2$$

$$\begin{array}{rrcr} x & - & 2y & + & 3z & = & 9 \\ & & y & + & 3z & = & 5 \\ & - & y & - & z & = & -1 \end{array} \quad \left[\begin{array}{cccc} 1 & -2 & 3 & 9 \\ 0 & 1 & 3 & 5 \\ 0 & -1 & -1 & -1 \end{array} \right]$$

$$r_{13}^{(-2)} : (-2)R_1 + R_3 \rightarrow R_3$$

Linear System

Associated
Augmented Matrix

Elementary
Row Operation

$$\begin{array}{rcrcrcrcrcl} x & - & 2y & + & 3z & = & 9 \\ & & y & + & 3z & = & 5 \\ & & & & 2z & = & 4 \end{array}$$

$$\left[\begin{array}{cccc} 1 & -2 & 3 & 9 \\ 0 & 1 & 3 & 5 \\ 0 & 0 & 2 & 4 \end{array} \right]$$

$$r_{23}^{(1)} : (1)R_2 + R_3 \rightarrow R_3$$

$$\begin{array}{rcrcrcrcrcl} x & - & 2y & + & 3z & = & 9 \\ & & y & + & 3z & = & 5 \\ & & & & z & = & 2 \end{array}$$

$$\left[\begin{array}{cccc} 1 & -2 & 3 & 9 \\ 0 & 1 & 3 & 5 \\ 0 & 0 & 1 & 2 \end{array} \right]$$

$$r_3^{(\frac{1}{2})} : (\frac{1}{2})R_3 \rightarrow R_3$$

$$\longrightarrow \begin{array}{rcrcrcrcl} x & & & = & 1 \\ & y & & = & -1 \\ & & z & = & 2 \end{array}$$

-
- Row-echelon form: (1, 2, 3)
 - Reduced row-echelon form: (1, 2, 3, 4)

(1) All row consisting entirely of zeros occur at the bottom of the matrix.

(2) For each row that does not consist entirely of zeros, the first nonzero entry is 1 (called **a leading 1**).

(3) For two successive (nonzero) rows, the leading 1 in the higher row is farther to the left than the leading 1 in the lower row.

(4) Every column that has a leading 1 has zeros in every position above and below its leading 1.

■ Ex 4: (Row-echelon form or reduced row-echelon form)

$$\begin{bmatrix} 1 & 2 & -1 & 4 \\ 0 & 1 & 0 & 3 \\ 0 & 0 & 1 & -2 \end{bmatrix}$$

(row - echelon form)

$$\begin{bmatrix} 0 & 1 & 0 & 5 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

(reduced row - echelon form)

$$\begin{bmatrix} 1 & -5 & 2 & -1 & 3 \\ 0 & 0 & 1 & 3 & -2 \\ 0 & 0 & 0 & 1 & 4 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

(row - echelon form)

$$\begin{bmatrix} 1 & 0 & 0 & -1 \\ 0 & 1 & 0 & 2 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

(reduced row - echelon form)

$$\begin{bmatrix} 1 & 2 & -3 & 4 \\ 0 & 2 & 1 & -1 \\ 0 & 0 & 1 & -3 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 2 & -1 & 2 \\ 0 & 0 & 0 & 0 \\ 0 & 1 & 2 & -4 \end{bmatrix}$$

- **Gaussian elimination:**

The procedure for reducing a matrix to a row-echelon form.

- **Gauss-Jordan elimination:**

The procedure for reducing a matrix to a reduced row-echelon form.

- **Notes:**

(1) Every matrix has an unique reduced row echelon form.

(2) A row-echelon form of a given matrix is not unique.

(Different sequences of row operations can produce different row-echelon forms.)

■ Ex: (Procedure of Gaussian elimination and Gauss-Jordan elimination)

$$\begin{bmatrix} 0 & 0 & -2 & 0 & 8 & 12 \\ 2 & 4 & -10 & 6 & 12 & 28 \\ 2 & 4 & -5 & 6 & -5 & 4 \end{bmatrix} \xrightarrow{r_{12}} \begin{bmatrix} 2 & 4 & -10 & 6 & 12 & 28 \\ 0 & 0 & -2 & 0 & 8 & 12 \\ 2 & 4 & -5 & 6 & -5 & 4 \end{bmatrix}$$

↑ The first nonzero column ← Produce leading 1

$$\xrightarrow{r_1^{(\frac{1}{2})}} \begin{bmatrix} 1 & 2 & -5 & 3 & 6 & 14 \\ 0 & 0 & -2 & 0 & 8 & 12 \\ 2 & 4 & -5 & 6 & -5 & 4 \end{bmatrix} \xrightarrow{r_{13}^{(-2)}} \begin{bmatrix} 1 & 4 & -3 & 2 & 6 & 14 \\ 0 & 0 & -2 & 0 & 8 & 12 \\ 0 & 0 & 5 & 0 & -17 & -24 \end{bmatrix}$$

leading 1 Produce leading 1
 ↑ Zeros elements below leading 1 ↑ The first nonzero Submatrix column

$$\begin{array}{c}
 \xrightarrow{r_2^{(-\frac{1}{2})}} \begin{bmatrix} 1 & 4 & -3 & 2 & 6 & 14 \\ 0 & 0 & \textcircled{1} & 0 & -4 & -6 \\ 0 & 0 & \textcircled{5} & 0 & -17 & -24 \end{bmatrix} \xrightarrow{r_{23}^{(-5)}} \begin{bmatrix} 1 & 4 & -3 & 2 & 6 & 14 \\ 0 & 0 & 1 & 0 & -4 & -6 \\ 0 & 0 & 0 & 0 & \boxed{3} & \boxed{6} \end{bmatrix}
 \end{array}$$

leading 1
 Zeros elements below leading 1
 Submatrix
 Produce leading 1

$$\begin{array}{c}
 \xrightarrow{r_3^{(\frac{1}{3})}} \begin{bmatrix} 1 & 4 & \boxed{-3} & 2 & \boxed{6} & 14 \\ 0 & 0 & 1 & 0 & \boxed{-4} & -6 \\ 0 & 0 & 0 & 0 & \textcircled{1} & 2 \end{bmatrix} \xrightarrow{r_{31}^{(-6)}} \begin{bmatrix} 1 & 4 & -3 & 2 & 0 & 2 \\ 0 & 0 & 1 & 0 & -4 & -6 \\ 0 & 0 & 0 & 0 & 1 & 2 \end{bmatrix}
 \end{array}$$

Zeros elsewhere
 leading 1
 (row - echelon form)
 (row - echelon form)

$$\begin{array}{c}
 \xrightarrow{r_{32}^{(4)}} \begin{bmatrix} 1 & 4 & -3 & 2 & 0 & 2 \\ 0 & 0 & 1 & 0 & 0 & 2 \\ 0 & 0 & 0 & 0 & 1 & 2 \end{bmatrix} \xrightarrow{r_{21}^{(3)}} \begin{bmatrix} 1 & 4 & 0 & 2 & 0 & 8 \\ 0 & 0 & 1 & 0 & 0 & 2 \\ 0 & 0 & 0 & 0 & 1 & 2 \end{bmatrix}
 \end{array}$$

(row - echelon form)
 (reduced row - echelon form)

- Ex 7: Solve a system by Gauss-Jordan elimination method (only one solution)

$$\begin{array}{rclcl} x & - & 2y & + & 3z & = & 9 \\ -x & + & 3y & & & = & -4 \\ 2x & - & 5y & + & 5z & = & 17 \end{array}$$

Sol:

augmented matrix

$$\begin{bmatrix} 1 & -2 & 3 & 9 \\ -1 & 3 & 0 & -4 \\ 2 & -5 & 5 & 17 \end{bmatrix} \xrightarrow{r_{12}^{(1)}, r_{13}^{(-2)}} \begin{bmatrix} 1 & -2 & 3 & 9 \\ 0 & 1 & 3 & 5 \\ 0 & -1 & -1 & -1 \end{bmatrix} \xrightarrow{r_{23}^{(1)}} \begin{bmatrix} 1 & -2 & 3 & 9 \\ 0 & 1 & 3 & 5 \\ 0 & 0 & 2 & 4 \end{bmatrix}$$

$$\xrightarrow{r_3^{(\frac{1}{2})}} \begin{bmatrix} 1 & -2 & 3 & 9 \\ 0 & 1 & 3 & 5 \\ 0 & 0 & 1 & 2 \end{bmatrix} \xrightarrow{r_{21}^{(2)}, r_{32}^{(-3)}, r_{31}^{(-9)}} \begin{bmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & 2 \end{bmatrix} \longrightarrow \begin{array}{lcl} x & = & 1 \\ y & = & -1 \\ z & = & 2 \end{array}$$

(row - echelon form)

(reduced row - echelon form)

$$\begin{array}{rcl} x & = & 1 \\ y & = & -1 \\ z & = & 2 \end{array}$$

-
- Ex 8 : Solve a system by Gauss-Jordan elimination method
(infinitely many solutions)

$$\begin{array}{rclcl} 2x_1 + & 4x_2 - & 2x_3 & = & 0 \\ 3x_1 + & 5x_2 & & = & 1 \end{array}$$

Sol: augmented matrix

$$\left[\begin{array}{cccc} 2 & 4 & -2 & 0 \\ 3 & 5 & 0 & 1 \end{array} \right] \xrightarrow{r_1^{(\frac{1}{2})}, r_{12}^{(-3)}, r_2^{(-1)}, r_{21}^{(-2)}} \left[\begin{array}{cccc} 1 & 0 & 5 & 2 \\ 0 & 1 & -3 & -1 \end{array} \right] \text{ (reduced row - echelon form)}$$

the corresponding system of equations is

$$\begin{array}{rclcl} x_1 & + & 5x_3 & = & 2 \\ & x_2 & - 3x_3 & = & -1 \end{array}$$

leading variable : x_1, x_2

free variable : x_3

$$\begin{aligned}x_1 &= 2 - 5x_3 \\x_2 &= -1 + 3x_3\end{aligned}$$

Let $x_3 = t$

$$\begin{aligned}x_1 &= 2 - 5t, \\x_2 &= -1 + 3t, \quad t \in \mathbb{R} \\x_3 &= t,\end{aligned}$$

So this system has infinitely many solutions.

- Homogeneous systems of linear equations:

A system of linear equations is said to be **homogeneous** if all the constant terms are zero.

$$\begin{array}{ccccccccc} a_{11}x_1 + & a_{12}x_2 + & a_{13}x_3 + & \cdots + & a_{1n}x_n & = & 0 \\ a_{21}x_1 + & a_{22}x_2 + & a_{23}x_3 + & \cdots + & a_{2n}x_n & = & 0 \\ a_{31}x_1 + & a_{32}x_2 + & a_{33}x_3 + & \cdots + & a_{3n}x_n & = & 0 \\ & & \vdots & & & & \\ a_{m1}x_1 + & a_{m2}x_2 + & a_{m3}x_3 + & \cdots + & a_{mn}x_n & = & 0 \end{array}$$

- **Trivial solution:**

$$x_1 = x_2 = x_3 = \cdots = x_n = 0$$

- **Nontrivial solution:**

other solutions

- **Notes:**

- (1) Every homogeneous system of linear equations is consistent.
- (2) If the homogenous system has fewer equations than variables, then it must have an infinite number of solutions.
- (3) For a homogeneous system, exactly one of the following is true.
 - (a) The system has only the trivial solution.
 - (b) The system has infinitely many nontrivial solutions in addition to the trivial solution.

-
- Ex 9: Solve the following homogeneous system

$$x_1 - x_2 + 3x_3 = 0$$

$$2x_1 + x_2 + 3x_3 = 0$$

Sol: augmented matrix

$$\begin{bmatrix} 1 & -1 & 3 & 0 \\ 2 & 1 & 3 & 0 \end{bmatrix} \xrightarrow{r_{12}^{(-2)}, r_2^{(\frac{1}{3})}, r_{21}^{(1)}} \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 1 & -1 & 0 \end{bmatrix} \text{ (reduced row - echelon form)}$$

leading variable : x_1, x_2

free variable : x_3

Let $x_3 = t$

$$x_1 = -2t, x_2 = t, x_3 = t, t \in R$$

When $t = 0, x_1 = x_2 = x_3 = 0$ (trivial solution)

Keywords in Section 1.2:

- matrix: مصفوفة
- row: صف
- column: عمود
- entry: **عنصر**
- size: حجم
- square matrix: مصفوفة مربعة
- symmetric matrix: مصفوفة متماثلة
- trace of a matrix: أثر المصفوفة
- order: ترتيب
- main diagonal: قطر رئيسي
- augmented matrix: مصفوفة موسعة
- coefficient matrix: معامل المصفوفة