

Chapter 5

Dot, Inner and Cross Products

5.1 Length of a vector

5.2 Dot Product

5.3 Inner Product

5.4 Cross Product

5.1 Length and Dot Product in R^n

- **Length:**

The length of a vector $\mathbf{v} = (v_1, v_2, \dots, v_n)$ in R^n is given by

$$\|\mathbf{v}\| = \sqrt{v_1^2 + v_2^2 + \dots + v_n^2}$$

- **Notes:** The length of a vector is also called its **norm**.

- **Notes: Properties of length**

(1) $\|\mathbf{v}\| \geq 0$

(2) $\|\mathbf{v}\| = 1 \Rightarrow \mathbf{v}$ is called a **unit vector**.

(3) $\|\mathbf{v}\| = 0$ iff $\mathbf{v} = \mathbf{0}$

(4) $\|c\mathbf{v}\| = |c|\|\mathbf{v}\|$

■ Ex 1:

(a) In R^5 , the length of $\mathbf{v} = (0, -2, 1, 4, -2)$ is given by

$$\|\mathbf{v}\| = \sqrt{0^2 + (-2)^2 + 1^2 + 4^2 + (-2)^2} = \sqrt{25} = 5$$

(b) In R^3 the length of $\mathbf{v} = (\frac{2}{\sqrt{17}}, \frac{-2}{\sqrt{17}}, \frac{3}{\sqrt{17}})$ is given by

$$\|\mathbf{v}\| = \sqrt{\left(\frac{2}{\sqrt{17}}\right)^2 + \left(\frac{-2}{\sqrt{17}}\right)^2 + \left(\frac{3}{\sqrt{17}}\right)^2} = \sqrt{\frac{17}{17}} = 1$$

(\mathbf{v} is a unit vector)

-
- A standard unit vector in R^n :

$$\{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n\} = \{(1, 0, \dots, 0), (0, 1, \dots, 0), (0, 0, \dots, 1)\}$$

- Ex:

the standard unit vector in R^2 : $\{i, j\} = \{(1, 0), (0, 1)\}$

the standard unit vector in R^3 : $\{i, j, k\} = \{(1, 0, 0), (0, 1, 0), (0, 0, 1)\}$

- Notes: (Two nonzero vectors are parallel)

$$\mathbf{u} = c\mathbf{v}$$

(1) $c > 0 \Rightarrow \mathbf{u}$ and \mathbf{v} have the same direction

(2) $c < 0 \Rightarrow \mathbf{u}$ and \mathbf{v} have the opposite direction

- **Thm 5.1: (Length of a scalar multiple)**

Let \mathbf{v} be a vector in R^n and c be a scalar. Then

$$\|c\mathbf{v}\| = |c| \|\mathbf{v}\|$$

Pf:

$$\mathbf{v} = (v_1, v_2, \dots, v_n)$$

$$\Rightarrow c\mathbf{v} = (cv_1, cv_2, \dots, cv_n)$$

$$\|c\mathbf{v}\| = \|(cv_1, cv_2, \dots, cv_n)\|$$

$$= \sqrt{(cv_1)^2 + (cv_2)^2 + \dots + (cv_n)^2}$$

$$= \sqrt{c^2(v_1^2 + v_2^2 + \dots + v_n^2)}$$

$$= |c| \sqrt{v_1^2 + v_2^2 + \dots + v_n^2}$$

$$= |c| \|\mathbf{v}\|$$

- **Thm 5.2: (Unit vector in the direction of \mathbf{v})**

If \mathbf{v} is a nonzero vector in R^n , then the vector $\mathbf{u} = \frac{\mathbf{v}}{\|\mathbf{v}\|}$

has length 1 and has the same direction as \mathbf{v} . This vector \mathbf{u} is called the **unit vector in the direction of \mathbf{v}** .

Pf:

$$\mathbf{v} \text{ is nonzero} \Rightarrow \|\mathbf{v}\| \neq 0 \Rightarrow \frac{1}{\|\mathbf{v}\|} > 0$$

$$\Rightarrow \mathbf{u} = \frac{1}{\|\mathbf{v}\|} \mathbf{v} \quad (\mathbf{u} \text{ has the same direction as } \mathbf{v})$$

$$\|\mathbf{u}\| = \left\| \frac{\mathbf{v}}{\|\mathbf{v}\|} \right\| = \frac{1}{\|\mathbf{v}\|} \|\mathbf{v}\| = 1 \quad (\mathbf{u} \text{ has length } 1)$$

■ **Notes:**

(1) The vector $\frac{\mathbf{v}}{\|\mathbf{v}\|}$ is called the unit vector in the direction of \mathbf{v} .

(2) The process of finding the unit vector in the direction of \mathbf{v} is called **normalizing** the vector \mathbf{v} .

■ Ex 2: (Finding a unit vector)

Find the unit vector in the direction of $\mathbf{v} = (3, -1, 2)$,
and verify that this vector has length 1.

Sol:

$$\mathbf{v} = (3, -1, 2) \Rightarrow \|\mathbf{v}\| = \sqrt{3^2 + (-1)^2 + 2^2} = \sqrt{14}$$

$$\Rightarrow \frac{\mathbf{v}}{\|\mathbf{v}\|} = \frac{(3, -1, 2)}{\sqrt{3^2 + (-1)^2 + 2^2}} = \frac{1}{\sqrt{14}} (3, -1, 2) = \left(\frac{3}{\sqrt{14}}, \frac{-1}{\sqrt{14}}, \frac{2}{\sqrt{14}} \right)$$

$$\therefore \sqrt{\left(\frac{3}{\sqrt{14}} \right)^2 + \left(\frac{-1}{\sqrt{14}} \right)^2 + \left(\frac{2}{\sqrt{14}} \right)^2} = \sqrt{\frac{14}{14}} = 1$$

$$\therefore \frac{\mathbf{v}}{\|\mathbf{v}\|} \text{ is a unit vector.}$$

- Distance between two vectors:

The **distance** between two vectors \mathbf{u} and \mathbf{v} in R^n is

$$d(\mathbf{u}, \mathbf{v}) = \|\mathbf{u} - \mathbf{v}\|$$

- Notes: (Properties of distance)

(1) $d(\mathbf{u}, \mathbf{v}) \geq 0$

(2) $d(\mathbf{u}, \mathbf{v}) = 0$ if and only if $\mathbf{u} = \mathbf{v}$

(3) $d(\mathbf{u}, \mathbf{v}) = d(\mathbf{v}, \mathbf{u})$

- Ex 3: (Finding the distance between two vectors)

The distance between $\mathbf{u}=(0, 2, 2)$ and $\mathbf{v}=(2, 0, 1)$ is

$$\begin{aligned}d(\mathbf{u}, \mathbf{v}) &= \|\mathbf{u} - \mathbf{v}\| = \|(0 - 2, 2 - 0, 2 - 1)\| \\&= \sqrt{(-2)^2 + 2^2 + 1^2} = 3\end{aligned}$$

Keywords in Section 5.1:

- length: طول
- norm: معيار
- unit vector: متجه الوحدة
- standard unit vector : متجه الوحدة الأساسي
- normalizing: معايرة
- distance: المسافة
- angle: زاوية
- triangle inequality: متباينة المثلث
- Pythagorean theorem: نظرية فيثاغورس

5.2 Dot Product

- Dot product in R^n :

The **dot product** of $\mathbf{u} = (u_1, u_2, \dots, u_n)$ and $\mathbf{v} = (v_1, v_2, \dots, v_n)$ is the scalar quantity

$$\mathbf{u} \cdot \mathbf{v} = u_1 v_1 + u_2 v_2 + \dots + u_n v_n$$

- Ex 4: (Finding the dot product of two vectors)

The dot product of $\mathbf{u} = (1, 2, 0, -3)$ and $\mathbf{v} = (3, -2, 4, 2)$ is

$$\mathbf{u} \cdot \mathbf{v} = (1)(3) + (2)(-2) + (0)(4) + (-3)(2) = -7$$

- **Thm 5.3: (Properties of the dot product)**

If \mathbf{u} , \mathbf{v} , and \mathbf{w} are vectors in R^n and c is a scalar, then the following properties are true.

$$(1) \quad \mathbf{u} \cdot \mathbf{v} = \mathbf{v} \cdot \mathbf{u}$$

$$(2) \quad \mathbf{u} \cdot (\mathbf{v} + \mathbf{w}) = \mathbf{u} \cdot \mathbf{v} + \mathbf{u} \cdot \mathbf{w}$$

$$(3) \quad c(\mathbf{u} \cdot \mathbf{v}) = (c\mathbf{u}) \cdot \mathbf{v} = \mathbf{u} \cdot (c\mathbf{v})$$

$$(4) \quad \mathbf{v} \cdot \mathbf{v} = \|\mathbf{v}\|^2$$

$$(5) \quad \mathbf{v} \cdot \mathbf{v} \geq 0 \quad \text{and} \quad \mathbf{v} \cdot \mathbf{v} = 0 \quad \text{if and only if} \quad \mathbf{v} = \mathbf{0}$$

- Euclidean n -space:

R^n was defined to be the *set* of all order n -tuples of real numbers. When R^n is combined with the standard operations of vector addition, scalar multiplication, vector length, and the dot product, the resulting vector space is called **Euclidean n -space**.

■ Ex 5: (Finding dot products)

$$\mathbf{u} = (2, -2), \mathbf{v} = (5, 8), \mathbf{w} = (-4, 3)$$

$$(a) \mathbf{u} \cdot \mathbf{v} \quad (b) (\mathbf{u} \cdot \mathbf{v})\mathbf{w} \quad (c) \mathbf{u} \cdot (2\mathbf{v}) \quad (d) \|\mathbf{w}\|^2 \quad (e) \mathbf{u} \cdot (\mathbf{v} - 2\mathbf{w})$$

Sol:

$$(a) \mathbf{u} \cdot \mathbf{v} = (2)(5) + (-2)(8) = -6$$

$$(b) (\mathbf{u} \cdot \mathbf{v})\mathbf{w} = -6\mathbf{w} = -6(-4, 3) = (24, -18)$$

$$(c) \mathbf{u} \cdot (2\mathbf{v}) = 2(\mathbf{u} \cdot \mathbf{v}) = 2(-6) = -12$$

$$(d) \|\mathbf{w}\|^2 = \mathbf{w} \cdot \mathbf{w} = (-4)(-4) + (3)(3) = 25$$

$$(e) \mathbf{v} - 2\mathbf{w} = (5 - (-8), 8 - 6) = (13, 2)$$

$$\mathbf{u} \cdot (\mathbf{v} - 2\mathbf{w}) = (2)(13) + (-2)(2) = 26 - 4 = 22$$

■ Ex 6: (Using the properties of the dot product)

Given $\mathbf{u} \cdot \mathbf{u} = 39$ $\mathbf{u} \cdot \mathbf{v} = -3$ $\mathbf{v} \cdot \mathbf{v} = 79$

Find $(\mathbf{u} + 2\mathbf{v}) \cdot (3\mathbf{u} + \mathbf{v})$

Sol:

$$\begin{aligned}(\mathbf{u} + 2\mathbf{v}) \cdot (3\mathbf{u} + \mathbf{v}) &= \mathbf{u} \cdot (3\mathbf{u} + \mathbf{v}) + 2\mathbf{v} \cdot (3\mathbf{u} + \mathbf{v}) \\&= \mathbf{u} \cdot (3\mathbf{u}) + \mathbf{u} \cdot \mathbf{v} + (2\mathbf{v}) \cdot (3\mathbf{u}) + (2\mathbf{v}) \cdot \mathbf{v} \\&= 3(\mathbf{u} \cdot \mathbf{u}) + \mathbf{u} \cdot \mathbf{v} + 6(\mathbf{v} \cdot \mathbf{u}) + 2(\mathbf{v} \cdot \mathbf{v}) \\&= 3(\mathbf{u} \cdot \mathbf{u}) + 7(\mathbf{u} \cdot \mathbf{v}) + 2(\mathbf{v} \cdot \mathbf{v}) \\&= 3(39) + 7(-3) + 2(79) = 254\end{aligned}$$

- Thm 5.4: (The Cauchy - Schwarz inequality)

If \mathbf{u} and \mathbf{v} are vectors in R^n , then

$$|\mathbf{u} \cdot \mathbf{v}| \leq \|\mathbf{u}\| \|\mathbf{v}\| \quad (|\mathbf{u} \cdot \mathbf{v}| \text{ denotes the absolute value of } \mathbf{u} \cdot \mathbf{v})$$

- Ex 7: (An example of the Cauchy - Schwarz inequality)

Verify the Cauchy - Schwarz inequality for $\mathbf{u}=(1, -1, 3)$
and $\mathbf{v}=(2, 0, -1)$

Sol: $\mathbf{u} \cdot \mathbf{v} = -1, \quad \mathbf{u} \cdot \mathbf{u} = 11, \quad \mathbf{v} \cdot \mathbf{v} = 5$


$$\Rightarrow |\mathbf{u} \cdot \mathbf{v}| = |-1| = 1$$

$$\|\mathbf{u}\| \|\mathbf{v}\| = \sqrt{\mathbf{u} \cdot \mathbf{u}} \cdot \sqrt{\mathbf{v} \cdot \mathbf{v}} = \sqrt{11} \cdot \sqrt{5} = \sqrt{55}$$

$$\therefore |\mathbf{u} \cdot \mathbf{v}| \leq \|\mathbf{u}\| \|\mathbf{v}\|$$

- The angle between two vectors in R^n :

$$\cos \theta = \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{u}\| \|\mathbf{v}\|}, 0 \leq \theta \leq \pi$$



- Note:

The angle between the zero vector and another vector is not defined.

■ Ex 8: (Finding the angle between two vectors)

$$\mathbf{u} = (-4, 0, 2, -2) \quad \mathbf{v} = (2, 0, -1, 1)$$

Sol:

$$\|\mathbf{u}\| = \sqrt{\mathbf{u} \cdot \mathbf{u}} = \sqrt{(-4)^2 + 0^2 + 2^2 + (-2)^2} = \sqrt{24}$$

$$\|\mathbf{v}\| = \sqrt{\mathbf{v} \cdot \mathbf{v}} = \sqrt{2^2 + (0)^2 + (-1)^2 + 1^2} = \sqrt{6}$$

$$\mathbf{u} \cdot \mathbf{v} = (-4)(2) + (0)(0) + (2)(-1) + (-2)(1) = -12$$

$$\Rightarrow \cos \theta = \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{u}\| \|\mathbf{v}\|} = \frac{-12}{\sqrt{24}\sqrt{6}} = -\frac{12}{\sqrt{144}} = -1$$

$$\Rightarrow \theta = \pi \quad \therefore \mathbf{u} \text{ and } \mathbf{v} \text{ have opposite directions. } (\mathbf{u} = -2\mathbf{v})$$

- Orthogonal vectors:

Two vectors \mathbf{u} and \mathbf{v} in R^n are orthogonal if

$$\mathbf{u} \cdot \mathbf{v} = 0$$

- Note:

The vector $\mathbf{0}$ is said to be orthogonal to every vector.

- Ex 10: (Finding orthogonal vectors)

Determine all vectors in R^n that are orthogonal to $\mathbf{u}=(4, 2)$.

Sol:

$$\mathbf{u} = (4, 2) \quad \text{Let } \mathbf{v} = (v_1, v_2)$$

$$\Rightarrow \mathbf{u} \cdot \mathbf{v} = (4, 2) \cdot (v_1, v_2)$$

$$= 4v_1 + 2v_2$$

$$= 0$$

$$\begin{bmatrix} 4 & 2 & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & \frac{1}{2} & 0 \end{bmatrix}$$

$$\Rightarrow v_1 = \frac{-t}{2}, \quad v_2 = t$$

$$\therefore \mathbf{v} = \left(\frac{-t}{2}, t \right), \quad t \in R$$

- **Thm 5.5: (The triangle inequality)**

If \mathbf{u} and \mathbf{v} are vectors in R^n , then $\|\mathbf{u} + \mathbf{v}\| \leq \|\mathbf{u}\| + \|\mathbf{v}\|$

Pf:

$$\begin{aligned}\|\mathbf{u} + \mathbf{v}\|^2 &= (\mathbf{u} + \mathbf{v}) \cdot (\mathbf{u} + \mathbf{v}) \\ &= \mathbf{u} \cdot (\mathbf{u} + \mathbf{v}) + \mathbf{v} \cdot (\mathbf{u} + \mathbf{v}) = \mathbf{u} \cdot \mathbf{u} + 2(\mathbf{u} \cdot \mathbf{v}) + \mathbf{v} \cdot \mathbf{v} \\ &= \|\mathbf{u}\|^2 + 2(\mathbf{u} \cdot \mathbf{v}) + \|\mathbf{v}\|^2 \leq \|\mathbf{u}\|^2 + 2|\mathbf{u} \cdot \mathbf{v}| + \|\mathbf{v}\|^2 \\ &\leq \|\mathbf{u}\|^2 + 2\|\mathbf{u}\|\|\mathbf{v}\| + \|\mathbf{v}\|^2 \\ &= (\|\mathbf{u}\| + \|\mathbf{v}\|)^2\end{aligned}$$

$$\therefore \|\mathbf{u} + \mathbf{v}\| \leq \|\mathbf{u}\| + \|\mathbf{v}\|$$

- **Note:**

Equality occurs in the triangle inequality if and only if the vectors \mathbf{u} and \mathbf{v} have the same direction.

- **Thm 5.6: (The Pythagorean theorem)**

If **u** and **v** are vectors in R^n , then **u** and **v** are orthogonal if and only if

$$\|\mathbf{u} + \mathbf{v}\|^2 = \|\mathbf{u}\|^2 + \|\mathbf{v}\|^2$$

- Dot product and matrix multiplication:

$$\mathbf{u} = \begin{bmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{bmatrix} \quad \mathbf{v} = \begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix} \quad (\text{A vector } \mathbf{u} = (u_1, u_2, \dots, u_n) \text{ in } R^n \text{ is represented as an } n \times 1 \text{ column matrix})$$

$$\mathbf{u} \cdot \mathbf{v} = \mathbf{u}^T \mathbf{v} = [u_1 \quad u_2 \quad \cdots \quad u_n] \begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{bmatrix} = [u_1 v_1 + u_2 v_2 + \cdots + u_n v_n]$$

Keywords in Section 5.2:

- dot product: الضرب النقطي
- Euclidean n -space: فضاء نوني اقليدي
- Cauchy-Schwarz inequality: متباينة كوشي-شوارز
- triangle inequality: متباينة المثلث
- Pythagorean theorem: نظرية فيثاغورس

5.3 Inner Product

- Inner product:

Let \mathbf{u} , \mathbf{v} , and \mathbf{w} be vectors in a vector space V , and let c be any scalar. An inner product on V is a function that associates a real number $\langle \mathbf{u}, \mathbf{v} \rangle$ with each pair of vectors \mathbf{u} and \mathbf{v} and satisfies the following axioms.

$$(1) \quad \langle \mathbf{u}, \mathbf{v} \rangle = \langle \mathbf{v}, \mathbf{u} \rangle$$

$$(2) \quad \langle \mathbf{u}, \mathbf{v} + \mathbf{w} \rangle = \langle \mathbf{u}, \mathbf{v} \rangle + \langle \mathbf{u}, \mathbf{w} \rangle$$

$$(3) \quad c \langle \mathbf{u}, \mathbf{v} \rangle = \langle c\mathbf{u}, \mathbf{v} \rangle$$

$$(4) \quad \langle \mathbf{v}, \mathbf{v} \rangle \geq 0 \quad \langle \mathbf{v}, \mathbf{v} \rangle = 0 \text{ if and only if } \mathbf{v} = \mathbf{0}$$

- **Note:**

$\mathbf{u} \cdot \mathbf{v}$ = dot product (Euclidean inner product for R^n)

$\langle \mathbf{u}, \mathbf{v} \rangle$ = general inner product for vector space V

- **Note:**

A vector space V with an inner product is called an **inner product space**.

Vector space: $(V, +, \bullet)$

Inner product space: $(V, +, \bullet, \langle, \rangle)$

- Ex 1: (The Euclidean inner product for R^n)

Show that the dot product in R^n satisfies the four axioms of an inner product.

Sol:

$$\mathbf{u} = (u_1, u_2, \dots, u_n) \quad , \quad \mathbf{v} = (v_1, v_2, \dots, v_n)$$

$$\langle \mathbf{u}, \mathbf{v} \rangle = \mathbf{u} \cdot \mathbf{v} = u_1 v_1 + u_2 v_2 + \dots + u_n v_n$$

By Theorem 5.3, this dot product satisfies the required four axioms. Thus it is an inner product on R^n .

■ Ex 2: (A different inner product for R^n)

Show that the function defines an inner product on R^2 , where $\mathbf{u} = (u_1, u_2)$ and $\mathbf{v} = (v_1, v_2)$.

$$\langle \mathbf{u}, \mathbf{v} \rangle = u_1 v_1 + 2u_2 v_2$$

Sol:

$$(a) \quad \langle \mathbf{u}, \mathbf{v} \rangle = u_1 v_1 + 2u_2 v_2 = v_1 u_1 + 2v_2 u_2 = \langle \mathbf{v}, \mathbf{u} \rangle$$

$$(b) \quad \mathbf{w} = (w_1, w_2)$$

$$\begin{aligned} \Rightarrow \langle \mathbf{u}, \mathbf{v} + \mathbf{w} \rangle &= u_1(v_1 + w_1) + 2u_2(v_2 + w_2) \\ &= u_1 v_1 + u_1 w_1 + 2u_2 v_2 + 2u_2 w_2 \\ &= (u_1 v_1 + 2u_2 v_2) + (u_1 w_1 + 2u_2 w_2) \\ &= \langle \mathbf{u}, \mathbf{v} \rangle + \langle \mathbf{u}, \mathbf{w} \rangle \end{aligned}$$

$$(c) \quad c \langle \mathbf{u}, \mathbf{v} \rangle = c(u_1 v_1 + 2u_2 v_2) = (cu_1)v_1 + 2(cu_2)v_2 = \langle c\mathbf{u}, \mathbf{v} \rangle$$

$$(d) \quad \langle \mathbf{v}, \mathbf{v} \rangle = v_1^2 + 2v_2^2 \geq 0$$

$$\langle \mathbf{v}, \mathbf{v} \rangle = 0 \Rightarrow v_1^2 + 2v_2^2 = 0 \quad \Rightarrow \quad v_1 = v_2 = 0 \quad (\mathbf{v} = \mathbf{0})$$

■ **Note:** (An inner product on \mathbb{R}^n)

$$\langle \mathbf{u}, \mathbf{v} \rangle = c_1 u_1 v_1 + c_2 u_2 v_2 + \cdots + c_n u_n v_n, \quad c_i > 0$$

- Ex 3: (A function that is not an inner product)

Show that the following function is not an inner product on R^3 .

$$\langle \mathbf{u} \cdot \mathbf{v} \rangle = u_1 v_1 - 2u_2 v_2 + u_3 v_3$$

Sol:

Let $\mathbf{v} = (1, 2, 1)$

Then $\langle \mathbf{v}, \mathbf{v} \rangle = (1)(1) - 2(2)(2) + (1)(1) = -6 < 0$

Axiom 4 is not satisfied.

Thus this function is not an inner product on R^3 .

- **Thm 5.7: (Properties of inner products)**

Let \mathbf{u} , \mathbf{v} , and \mathbf{w} be vectors in an inner product space V , and let c be any real number.

$$(1) \quad \langle \mathbf{0}, \mathbf{v} \rangle = \langle \mathbf{v}, \mathbf{0} \rangle = 0$$

$$(2) \quad \langle \mathbf{u} + \mathbf{v}, \mathbf{w} \rangle = \langle \mathbf{u}, \mathbf{w} \rangle + \langle \mathbf{v}, \mathbf{w} \rangle$$

$$(3) \quad \langle \mathbf{u}, c\mathbf{v} \rangle = c \langle \mathbf{u}, \mathbf{v} \rangle$$

- **Norm (length) of \mathbf{u} :**

$$\|\mathbf{u}\| = \sqrt{\langle \mathbf{u}, \mathbf{u} \rangle}$$

- **Note:**

$$\|\mathbf{u}\|^2 = \langle \mathbf{u}, \mathbf{u} \rangle$$

-
- Distance between \mathbf{u} and \mathbf{v} :

$$d(\mathbf{u}, \mathbf{v}) = \|\mathbf{u} - \mathbf{v}\| = \sqrt{\langle \mathbf{u} - \mathbf{v}, \mathbf{u} - \mathbf{v} \rangle}$$

- Angle between two nonzero vectors \mathbf{u} and \mathbf{v} :

$$\cos \theta = \frac{\langle \mathbf{u}, \mathbf{v} \rangle}{\|\mathbf{u}\| \|\mathbf{v}\|}, \quad 0 \leq \theta \leq \pi$$

- Orthogonal: $(\mathbf{u} \perp \mathbf{v})$

\mathbf{u} and \mathbf{v} are **orthogonal** if $\langle \mathbf{u}, \mathbf{v} \rangle = 0$.

- Notes:

(1) If $\|\mathbf{v}\| = 1$, then \mathbf{v} is called a **unit vector**.

(2) $\|\mathbf{v}\| \neq 1$
 $\mathbf{v} \neq 0$ $\xrightarrow{\text{Normalizing}}$ $\frac{\mathbf{v}}{\|\mathbf{v}\|}$ (the unit vector in the direction of \mathbf{v})
not a unit vector

■ Ex 6: (Finding inner product)

$\langle p, q \rangle = a_0b_0 + a_1b_1 + \cdots + a_nb_n$ is an inner product

Let $p(x) = 1 - 2x^2$, $q(x) = 4 - 2x + x^2$ be polynomials in $P_2(x)$

(a) $\langle p, q \rangle = ?$ (b) $\|q\| = ?$ (c) $d(p, q) = ?$

Sol:

$$(a) \quad \langle p, q \rangle = (1)(4) + (0)(-2) + (-2)(1) = 2$$

$$(b) \quad \|q\| = \sqrt{\langle q, q \rangle} = \sqrt{4^2 + (-2)^2 + 1^2} = \sqrt{21}$$

$$(c) \quad \because p - q = -3 + 2x - 3x^2$$

$$\therefore d(p, q) = \|p - q\| = \sqrt{\langle p - q, p - q \rangle}$$

$$= \sqrt{(-3)^2 + 2^2 + (-3)^2} = \sqrt{22}$$

- Properties of norm:

(1) $\|\mathbf{u}\| \geq 0$

(2) $\|\mathbf{u}\| = 0$ if and only if $\mathbf{u} = \mathbf{0}$

(3) $\|c\mathbf{u}\| = |c| \|\mathbf{u}\|$

- Properties of distance:

(1) $d(\mathbf{u}, \mathbf{v}) \geq 0$

(2) $d(\mathbf{u}, \mathbf{v}) = 0$ if and only if $\mathbf{u} = \mathbf{v}$

(3) $d(\mathbf{u}, \mathbf{v}) = d(\mathbf{v}, \mathbf{u})$

■ **Thm 5.8 :**

Let \mathbf{u} and \mathbf{v} be vectors in an inner product space V .

(1) Cauchy-Schwarz inequality:

$$|\langle \mathbf{u}, \mathbf{v} \rangle| \leq \|\mathbf{u}\| \|\mathbf{v}\| \quad \text{Theorem 5.4}$$

(2) Triangle inequality:

$$\|\mathbf{u} + \mathbf{v}\| \leq \|\mathbf{u}\| + \|\mathbf{v}\| \quad \text{Theorem 5.5}$$

(3) Pythagorean theorem :

\mathbf{u} and \mathbf{v} are orthogonal if and only if

$$\|\mathbf{u} + \mathbf{v}\|^2 = \|\mathbf{u}\|^2 + \|\mathbf{v}\|^2 \quad \text{Theorem 5.6}$$

- **Orthogonal projections in inner product spaces:**

Let \mathbf{u} and \mathbf{v} be two vectors in an inner product space V , such that $\mathbf{v} \neq \mathbf{0}$. Then the **orthogonal projection of \mathbf{u} onto \mathbf{v}** is given by

$$\text{proj}_{\mathbf{v}} \mathbf{u} = \frac{\langle \mathbf{u}, \mathbf{v} \rangle}{\langle \mathbf{v}, \mathbf{v} \rangle} \mathbf{v}$$

- **Note:**

If \mathbf{v} is a unit vector, then $\langle \mathbf{v}, \mathbf{v} \rangle = \|\mathbf{v}\|^2 = 1$.

The formula for the orthogonal projection of \mathbf{u} onto \mathbf{v} takes the following simpler form.

$$\text{proj}_{\mathbf{v}} \mathbf{u} = \langle \mathbf{u}, \mathbf{v} \rangle \mathbf{v}$$

- Ex 10: (Finding an orthogonal projection in R^3)

Use the Euclidean inner product in R^3 to find the orthogonal projection of $\mathbf{u}=(6, 2, 4)$ onto $\mathbf{v}=(1, 2, 0)$.

Sol:

$$\because \langle \mathbf{u}, \mathbf{v} \rangle = (6)(1) + (2)(2) + (4)(0) = 10$$

$$\langle \mathbf{v}, \mathbf{v} \rangle = 1^2 + 2^2 + 0^2 = 5$$

$$\therefore \text{proj}_{\mathbf{v}} \mathbf{u} = \frac{\mathbf{u} \cdot \mathbf{v}}{\mathbf{v} \cdot \mathbf{v}} \mathbf{v} = \frac{10}{5} (1, 2, 0) = (2, 4, 0)$$

- Note:

$\mathbf{u} - \text{proj}_{\mathbf{v}} \mathbf{u} = (6, 2, 4) - (2, 4, 0) = (4, -2, 4)$ is orthogonal to $\mathbf{v} = (1, 2, 0)$.

- **Thm 5.9: (Orthogonal projection and distance)**

Let \mathbf{u} and \mathbf{v} be two vectors in an inner product space V , such that $\mathbf{v} \neq \mathbf{0}$. Then

$$d(\mathbf{u}, \text{proj}_{\mathbf{v}} \mathbf{u}) < d(\mathbf{u}, c\mathbf{v}), \quad c \neq \frac{\langle \mathbf{u}, \mathbf{v} \rangle}{\langle \mathbf{v}, \mathbf{v} \rangle}$$

Keywords in Section 5.2:

- inner product: ضرب داخلي
- inner product space: فضاء الضرب الداخلي
- norm: معيار
- distance: مسافة
- angle: زاوية
- orthogonal: متعامد
- unit vector: متجه وحدة
- normalizing: معايرة
- Cauchy – Schwarz inequality: متباينة كوشي - شوارز
- triangle inequality: متباينة المثلث
- Pythagorean theorem: نظرية فيثاغورس
- orthogonal projection: إسقاط عمودي

5.4 Cross Product

- **Cross product in R^3 :**

The **cross product** of $\mathbf{u} = (u_1, u_2, u_3)$ and $\mathbf{v} = (v_1, v_2, v_3)$ is the vector quantity

$$\mathbf{w} = \mathbf{u} \times \mathbf{v} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \end{vmatrix} = \left(\begin{vmatrix} u_2 & u_3 \\ v_2 & v_3 \end{vmatrix}, -\begin{vmatrix} u_1 & u_3 \\ v_1 & v_3 \end{vmatrix}, \begin{vmatrix} u_1 & u_2 \\ v_1 & v_2 \end{vmatrix} \right)$$

$$\mathbf{w} = \mathbf{u} \times \mathbf{v} = (u_2 v_3 - u_3 v_2, u_3 v_1 - u_1 v_3, u_1 v_2 - u_2 v_1)$$

- **Ex 11: (Finding the cross product of two vectors)**

The cross product of $\mathbf{u}=(1, 2, 0)$ and $\mathbf{v}=(3, -2, 4)$ is

$$\mathbf{w} = \mathbf{u} \times \mathbf{v} = (8, -4, -8)$$

- **Thm 5.10: Relationships involving cross product and dot product**

Let \mathbf{u} , \mathbf{v} and \mathbf{w} be 3 vectors in \mathbb{R}^3 , then:

$$\begin{aligned} (a) \quad & \mathbf{u} \cdot (\mathbf{u} \times \mathbf{v}) = 0 && (\mathbf{u} \times \mathbf{v} \text{ is orthogonal to } \mathbf{u}) \\ (b) \quad & \mathbf{v} \cdot (\mathbf{u} \times \mathbf{v}) = 0 && (\mathbf{u} \times \mathbf{v} \text{ is orthogonal to } \mathbf{v}) \\ (c) \quad & \|\mathbf{u} \times \mathbf{v}\|^2 = \|\mathbf{u}\|^2 \|\mathbf{v}\|^2 - (\mathbf{u} \cdot \mathbf{v})^2 && (\text{Lagrange's identity}) \\ (d) \quad & \mathbf{u} \times (\mathbf{v} \times \mathbf{w}) = (\mathbf{u} \cdot \mathbf{w})\mathbf{v} - (\mathbf{u} \cdot \mathbf{v})\mathbf{w} && (\text{relationship between cross and dot products}) \\ (e) \quad & (\mathbf{u} \times \mathbf{v}) \times \mathbf{w} = (\mathbf{u} \cdot \mathbf{w})\mathbf{v} - (\mathbf{v} \cdot \mathbf{w})\mathbf{u} && (\text{relationship between cross and dot products}) \end{aligned}$$

- **Thm 5.11: Properties of involving cross product**


Let \mathbf{u} , \mathbf{v} and \mathbf{w} be 3 vectors in \mathbb{R}^3 and k a scalar, then:

$$\begin{aligned} (a) \quad & \mathbf{u} \times \mathbf{v} = -(\mathbf{v} \times \mathbf{u}) \\ (b) \quad & \mathbf{u} \times (\mathbf{v} + \mathbf{w}) = (\mathbf{u} \times \mathbf{v}) + (\mathbf{u} \times \mathbf{w}) \\ (c) \quad & (\mathbf{u} + \mathbf{v}) \times \mathbf{w} = (\mathbf{u} \times \mathbf{w}) + (\mathbf{v} \times \mathbf{w}) \\ (d) \quad & k(\mathbf{u} \times \mathbf{v}) = (k\mathbf{u}) \times \mathbf{v} = \mathbf{u} \times (k\mathbf{v}) \\ (e) \quad & \mathbf{u} \times \mathbf{0} = \mathbf{0} \times \mathbf{u} = \mathbf{0} \\ (f) \quad & \mathbf{u} \times \mathbf{u} = \mathbf{0} \end{aligned}$$

- **Thm 5.12: Scalar triple product**

Let \mathbf{u} , \mathbf{v} and \mathbf{w} be 3 vectors in \mathbb{R}^3 , then:

$$V = \mathbf{w} \bullet (\mathbf{u} \times \mathbf{v}) = \begin{vmatrix} u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \\ w_1 & w_2 & w_3 \end{vmatrix} = \text{Volume of the parallelepiped determined by the 3 vectors}$$



Keywords in Section 5.4:

- Cross product: ضرب خارجي
- Scalar triple product: الضرب الثلاثي العددي
- angle: زاوية
- orthogonal: متعامد