

STAT 109
Biostatistics

CHAPTER 4: Probabilistic Features of Certain Data Distribution (Probability Distributions)

September 2025

NOTE: This presentation is based on the presentation prepared thankfully by Professor Abdullah al-Shiha.

Last Edited by: Kholoud Basalim

4.1 Introduction

4.2 Probability Distributions of Discrete Random Variables

4.3 Binomial Distribution

4.4 Poisson Distribution

4.5 Continuous Probability Distributions

4.6 Normal Distribution

4.7 Normal Distribution Application

4.8 t-Distribution

Introduction

The concept of random variables is very important in Statistics. Some events can be defined using random variables.

There are two types of random variables:

1. Discrete Random variables
2. Continuous Random Variables

Probability Distributions of Discrete Random Variables (r.v.):

Definition :

The probability distribution of a discrete random variable is a table, graph, formula or other device used to specify all possible values of the random variable along with their respective probabilities.

Examples of discrete r .v's :

- The no. of patients visiting KKUH in a week.
- The no. of times a person had a cold in last year.

Example:

Consider the following discrete random variable :

X = The number of times a Saudi person had a cold in January 2010

Suppose we are able to count the no. of Saudis which $X = x$:

x (no. of colds a Saudi person had in January 2010)	Frequency of x (no. of Saudi people who had a cold x times in January 2010)
0	10,000,000
1	3,000,000
2	2,000,000
3	1,000,000
Total	$N = 16,000,000$

X = The number of times a Saudi person had a cold in January 2010

Note that the possible values of the random variable X are:

$$x = 0, 1, 2, 3$$

Definition random variable

Experiment:

Selecting a person at random

Define the event:

$(X = 0)$ = The event that the selected person had no cold.

$(X = 1)$ = The event that the selected person had 1 cold.

$(X = 2)$ = The event that the selected person had 2 colds.

$(X = 3)$ = The event that the selected person had 3 colds.

In general

$(X = x)$ = The event that the selected person had x colds.

Value of the random variable

For this experiment, there are $n(\Omega) = 16,000,000$ equally likely outcomes .

The number of elements of the event $(X = x)$ is :

$n(X = x)$ = no. of Saudi people who had a cold x times in January 2010 .
= frequency of x .

The probability of event $(X = x)$ is :

$$P(X = x) = \frac{n(X=x)}{n(\Omega)} = \frac{n(X=x)}{16,000,000} , \quad \text{for } x = 0, 1, 2, 3$$

x	frequency of x $n(X = x)$	$P(X = x) = \frac{n(X = x)}{16,000,000}$ (Relative frequency)
0	10000000	0.6250
1	3000000	0.1875
2	2000000	0.1250
3	1000000	0.0625
Total	16000000	1

Note :

$$P(X = x) = \frac{n(X=x)}{16,000,000} = \text{Relative frequency} = \frac{\text{frequency}}{16,000,000}$$

The probability distribution of the discrete random variable X is given by the following table:

x	$P(X = x) = f(x)$
0	0.6250 $\leftarrow P(X = 0)$
1	0.1874 $\leftarrow P(X = 1)$
2	0.1250 $\leftarrow P(X = 2)$
3	0.0625 $\leftarrow P(X = 3)$
Total	1

Notes:

- The probability distribution of any discrete random variable X must satisfy the following two properties:

$$(1) \quad 0 \leq P(X = x) \leq 1 \qquad (2) \quad \sum_x P(X = x) = 1$$

- Using the probability distribution of a discrete r.v. we can find the probability of r.v. X .

Example:

Consider the discrete r.v. X in the previous example.

$$(1) P(X \geq 2) = P(X = 2) + P(X = 3) = 0.1250 + 0.0625 = 0.1875$$

$$(2) P(X > 2) = P(X = 3) = 0.0625 \quad [\text{note: } P(X > 2) \neq P(X \geq 2)]$$

$$(3) P(1 \leq X < 3) = P(X = 1) + P(X = 2) = 0.1875 + 0.1250 = 0.3125$$

$$(4) P(X \leq 2) = P(X = 0) + P(X = 1) + P(X = 2) \\ = 0.6250 + 0.1875 + 0.1250 = 0.9375$$

Another solution :

$$P(X \leq 2) = 1 - P(X > 2) \\ = 1 - P(X = 3) \\ = 1 - 0.0625 \\ = 0.9375$$

x	P(X = x)
0	0.6250
1	0.1875
2	0.1250
3	0.0625
Total	1

$$(5) P(-1 < X < 2) = P(X = 0) + P(X = 1) = 0.6250 + 0.1875 = 0.8125$$

$$(6) P(-1.5 < X < 1.3) = P(X = 0) + P(X = 1) = 0.6250 + 0.1875 = 0.8125$$

$$(7) P(X = 3.5) = P(3.5) = 0$$

$$(8) P(X < 10) = P(X = 0) + P(X = 1) + P(X = 2) + P(X = 3) \\ = 0.6250 + 0.1875 + 0.1250 + 0.0625 = 1$$

(9) The probability that the selected person had at least 2 colds :

$$P(X \geq 2) = P(X = 2) + P(X = 3) = 0.1875$$

(10) The probability that the selected person had at most 2 colds :

$$P(X \leq 2) = P(X = 2) + P(X = 1) + P(X = 0) = 0.9375$$

(11) The probability that the selected person had more than 2 colds :

$$P(X > 2) = P(X = 3) = 0.0625$$

(12) The probability that the selected person had less than 2 colds :

$$P(X < 2) = P(X = 0) + P(X = 1) = 0.8125$$

x	P(X = x)
0	0.6250
1	0.1875
2	0.1250
3	0.0625
Total	1


Graphical Presentation:

The probability distribution of a discrete r. v. X can be graphically represented .

Example:

The probability distribution of the random variable in the previous example is:

The graphical presentation of this probability distribution is given by the following figure:



x	P(X = x)
0	0.6250
1	0.1875
2	0.1250
3	0.0625
Total	1

Find the Probability that the selected person had no cold in January 2010 ?

$$P(X = 0) = 0.6$$

Mean and Variance of a Discrete Random Variable :

Mean :

The mean (or expected value) of a discrete random variable X is denoted by μ_X or μ .

It is defined by:

$$\mu = \sum_x x P(X = x)$$

Variance:

The variance of a discrete random variable X is denoted by σ_X^2 or σ^2

It is defined by:

$$\sigma^2 = \sum_x (x - \mu)^2 P(X = x)$$

Standard Deviation:

$$\sigma = \sqrt{\text{variance}}$$

Example:

We wish to calculate the **mean** and the **variance** of the discrete r. v. X whose probability distribution is given by the following table:

x	P(X=x)
0	0.05
1	0.25
2	0.45
3	0.25
Total	1

Solution:

$$\begin{aligned}\mu &= \sum_x x P(X = x) \\ &= (0 \times 0.05) + (1 \times 0.25) + (2 \times 0.45) + (3 \times 0.25) = \mathbf{1.9 \text{ Mean}}\end{aligned}$$

$$\begin{aligned}\sigma^2 &= \sum_x (x - \mu)^2 P(X = x) \\ &= \sum_x (x - 1.9)^2 P(X = x) \\ &= \mathbf{0.69 \text{ Variance}}\end{aligned}$$

$$\sigma = \sqrt{\text{variance}} = \sqrt{0.69} = \mathbf{0.83066 \text{ Standard Deviation}}$$

x	P(X=x)	xP(X=x)
0	0.05	0
1	0.25	0.25
2	0.45	0.9
3	0.25	0.75
Total	1	1.9

Mean : $\mu = \sum_x x P(X = x)$

Cumulative Distributions:

The cumulative distribution function of a discrete r. v. X is defined by :

$$P(X \leq x) = \sum_{a \leq x} P(X = a)$$

Example:

Calculate the cumulative distribution of the discrete r. v. X whose probability distribution is given by the following table:

x	$P(X=x)$	$P(X \leq x)$
0	0.05	0.05
1	0.25	$0.25+0.05= 0.30$
2	0.45	$0.45+0.25+0.05= 0.75$
3	0.25	$0.25+0.45+0.25+0.05= 1$
Total	1	--

Use the cumulative distribution to find:

$P(X \leq 2), P(X < 2),$

$P(X \leq 1.5), P(X < 1.5),$

$P(X > 1), P(X \geq 1) .$

cumulative distribution

$$F(x) = P(X \leq x)$$

Solution :

The cumulative distribution of X is :

Using the cumulative distribution :

$$P(X \leq 2) = 0.75$$

$$P(X < 2) = P(X \leq 1) = 0.30$$

$$P(X < 1.5) = P(X \leq 1) = 0.30$$

$$P(X \leq 1.5) = P(X \leq 1) = 0.30$$

$$\begin{aligned} P(X > 1) &= 1 - P(X \leq 1) \\ &= 1 - 0.3 \\ &= 0.7 \end{aligned}$$

$$\begin{aligned} P(X \geq 1) &= 1 - P(X < 1) \\ &= 1 - P(X \leq 0) \\ &= 1 - 0.05 \\ &= 0.95 \end{aligned}$$

x	P(X=x)	P(X ≤ x)
0	0.05	0.05 ← P(X ≤ 0)
1	0.25	0.25+0.05= 0.30 ← P(X ≤ 1)
2	0.45	0.45+0.25+0.05= 0.75 ← P(X ≤ 2)
3	0.25	0.25+0.45+0.25+0.05= 1 ← P(X ≤ 3)
Total	1	--

Note:

$$P(X \leq 0) = P(X = 0)$$

$$P(X \leq 1) = P(X = 0) + P(X = 1)$$

$$P(X \leq 2) = P(X = 0) + P(X = 1) + P(X = 2)$$

$$P(X \leq 3) = P(X = 0) + P(X = 1) + P(X = 2) + P(X = 3)$$

Probability Distribution

x	$P(X=x)$
0	0.05
1	0.25
2	0.45
3	0.25
Total	1



Cumulative Distribution

x	$P(X \leq x)$
0	0.05
1	$0.25+0.05=$ 0.30
2	$0.45+0.25+0.05=$ 0.75
3	$0.25+0.45+0.25+0.05=$ 1
Total	--


Cumulative Distribution

x	$P(X \leq x)$
0	0.05
1	0.30
2	0.75
3	1
Total	--



Probability Distribution

x	$P(X=x)$
0	0.05
1	$0.30 - 0.05 =$ 0.25
2	$0.75 - 0.30 =$ 0.45
3	$1 - 0.75 =$ 0.25
Total	1



Complement of probability:

- $P(X \leq a) = 1 - P(X > a)$
- $P(X < a) = 1 - P(X \geq a)$
- $P(X \geq a) = 1 - P(X < a)$
- $P(X > a) = 1 - P(X \leq a)$

At least **a** :

$$P(X \geq a)$$

At most **a** :

$$P(X \leq a)$$

Less than **a** :

$$P(X < a)$$

More than **a**:

$$P(X > a)$$

Exactly **a**:

$$P(X = a)$$

Between **a** and **b** :

$$P(a < X < b)$$

Example: (Reading Assignment)

Given the following probability distribution of a discrete random variable X representing the number of defective teeth of the patient visiting a certain dental clinic :

A- Find the value of K.

B- Find the flowing probabilities:

1- $P(X < 3)$

2- $P(X \leq 3)$

3- $P(X < 6)$

4- $P(X < 1)$

5- $P(X = 3.5)$

C- Find the probability that the patient has at least 4 defective teeth.

D- Find the probability that the patient has at most 2 defective teeth.

E- Find the expected number of defective teeth (mean of X).

F- Find the variance of X.

x	P(X=x)
1	0.25
2	0.35
3	0.20
4	0.15
5	K

Solution:

A- Find the value of K. **The probability distribution of any discrete random variable X must satisfy**

$$\begin{aligned}\sum_x P(X = x) &= 1 \\ 0.25 + 0.35 + 0.20 + 0.15 + K &= 1 \\ 0.95 + K &= 1 \\ K &= 0.05\end{aligned}$$

B- Find the following probabilities:

- 1- $P(X < 3) = P(X=1) + P(X=2) = 0.25 + 0.35 = 0.60$
- 2- $P(X \leq 3) = P(X=1) + P(X=2) + P(X=3) = 0.25 + 0.35 + 0.20 = 0.8$
- 3- $P(X < 6) = P(X=1) + P(X=2) + P(X=3) + P(X=4) + P(X=5) = 1$
- 4- $P(X < 1) = 0$
- 5- $P(X = 3.5) = 0$

x	P(X=x)
1	0.25
2	0.35
3	0.20
4	0.15
5	K = 0.05
Total	1

C- Find the probability that the patient has **at least 4** defective teeth.

$$P(X \geq 4) = P(X = 4) + P(X = 5) = 0.15 + 0.05 = 0.20$$

D- Find the probability that the patient has **at most 2** defective teeth.

$$P(X \leq 2) = P(X = 1) + P(X = 2) = 0.25 + 0.35 = 0.6$$

x	P(X=x)	x P(X=x)	(x - μ)	(x - μ) ²	(x - μ) ² P(X = x)
1	0.25	0.25	-1.4	1.96	0.49
2	0.35	0.70	-0.4	0.16	0.056
3	0.20	0.60	0.6	0.36	0.072
4	0.15	0.60	1.6	2.56	0.384
5	0.05	0.25	2.6	6.76	0.338
Total	1	2.4	-	-	1.34

Mean : $\mu = \sum_x x P(X = x)$



Variance: $\sigma^2 = \sum_x (x - \mu)^2 P(X = x)$



E- Find the expected number of defective teeth (mean of X).

$$\mu = \sum_x x P(X = x)$$

$$= (1 \times 0.25) + (2 \times 0.35) + (3 \times 0.2) + (4 \times 0.15) + (5 \times 0.05) = 2.4$$

F- Find the variance of X.

$$\sigma^2 = \sum_x (x - \mu)^2 P(X = x) = 1.34$$

$$\sigma = \sqrt{\text{variance}} = \sqrt{1.34} = 1.1576 \text{ Standard Deviation}$$

Notation (n!) : **n!** is read (**n factorial**) .

It defined by :

$$n! = n(n - 1)(n - 2) \dots (2)(1) \quad \text{for } n \geq 1$$

Examples :

$$5! = 5 \times 4 \times 3 \times 2 \times 1 = 120$$

Combinations:

The number of different ways for selecting r objects from n distinct objects is denoted by C_r^n or $\binom{n}{r}$ and is given by :

$$C(\text{Number of items in set}, \text{Number of items selected from the set}) = C_r^n = \frac{n!}{(n-r)!r!}$$

Notes:

1. C_r^n is read as “ n “ choose “ r ”.
2. $C_n^n = 1$, $C_0^n = 1$
3. $C_r^n = C_{n-r}^n$ (for example : $C_3^{10} = C_7^{10}$)
4. C_r^n = number of unordered subsets of a set of (n) objects such that each subset contains (r) objects.

Example: For $n = 4$ and $r = 2$:

$$C_2^4 = \frac{4!}{2!(4-2)!} = \frac{4!}{2! \times 2!} = \frac{4 \times 3 \times 2 \times 1}{(2 \times 1)(2 \times 1)} = 6$$

$C_2^4 = 6$ = The number of different ways for selecting 2 objects from 4 distinct objects.

Example:

Suppose that we have the set **{a, b, c, d}** of (n=4) objects.

We wish to choose a subset of two objects. The possible subsets of this set with 2 elements in each subset are:

{a , b}, {a , c}, {a , d}, {b , d}, {b , c}, {c , d}

The number of these subset is : $C_2^4 = 6$

4.3 Binomial Distribution:

Bernoulli Trial:

Is an experiment with only two possible outcomes: **S = success** and **F= failure**

(Boy or girl , Saudi or non-Saudi , sick or well , dead or alive).

- Binomial distribution is a discrete distribution.

Binomial Distribution:

- Binomial distribution is used to model an experiment for which:

1. The experiment has a sequence of **n** Bernoulli trials.
2. The probability of success is **$P(S) = p$** , and the probability of failure is **$P(F) = 1 - p = q$** .
3. The probability of success **$P(S) = p$** is constant for each trial .
4. The trials are independent; that is the outcome of one trial hasno effect on the outcome of any other trial.

In this type of experiment, we are interested in the discrete r. v. representing the number of successes in the n trials.

X = The number of successes in the n trials

The possible values of X (number of success in n trials) are:

$$x = 0, 1, 2, \dots, n$$

The r.v. X has a **binomial distribution with parameters n and p** , and we write:

$$X \sim \text{Binomial}(n, p)$$

n =number of trials
 P = probability of success

The probability distribution of X is given by:

$$P(X=x) = \begin{cases} C_x^n p^x q^{n-x} & ; \text{ for } x=0,1,2,\dots,n \\ 0 & ; \text{ otherwise} \end{cases} ; q=1-p$$

Where : $C_x^n = \frac{n!}{x!(n-x)!}$

We can write the probability distribution of **X** as a table *as follows* :

x	$P(X=x) = C_x^n p^x q^{n-x}$
0	$C_0^n p^0 q^{n-0} = q^n$
1	$C_1^n p^1 q^{n-1}$
2	$C_2^n p^2 q^{n-2}$
⋮	⋮
⋮	⋮
⋮	⋮
n-1	$C_{n-1}^n p^{n-1} q^1$
n	$C_n^n p^n q^0 = p^n$
Total	1

Result: (Mean and Variance for Binomial distribution)

If $X \sim \text{Binomial}(n, p)$, then

1-The mean (expected value):

$$\mu = np$$

2-The variance:

$$\sigma^2 = npq$$

3- The standard deviation:

$$\sigma = \sqrt{npq}$$

Example:

Suppose that the probability that a Saudi man has **high blood pressure is 0.15** . Suppose that we randomly **select a sample of 6 Saudi men**.

1. Find the probability distribution of the random variable (X) representing the number of men with high blood pressure in the sample.
2. Find the expected number of men with high blood pressure in the sample (mean of X).
3. Find the variance X.
4. What is the probability that there will be exactly 2 men with high blood pressure?
5. What is the probability that there will be at most 2 men with high blood pressure?
6. What is the probability that there will be at least 4 men with high blood pressure?

Solution:

We are interested in the following random variable:

X = The number of men with high blood pressure in the sample of 6 men.

Notes:

- Bernoulli trial: diagnosing whether a man has a high bloodpressure or not.
There are two outcomes for each trial:

S = Success: The man has high blood pressure

F = failure: The man does not have high blood pressure.

Number of trials = 6 (we need to check 6 men)

-Probability of success: $P(S) = p = 0.15$

- Probability of failure: $P(F) = q = 1 - p = 0.85$

- Number of trials: $n = 6$

- The trials are independent because the result of each man does not affect the result of any other man since the selection was made at random.

- The random variable **X** has a binomial distribution with parameters: **$n = 6$ and $p = 0.15$** , that is:

$$\mathbf{X \sim \text{Binomial}(n, p)}$$

$$\mathbf{X \sim \text{Binomial}(6, 0.15)}$$

- The possible values of **X** are:

$$\mathbf{x = 0, 1, 2, 3, 4, 5, 6}$$

1. The probability distribution of X is:

$$P(X=x)=\begin{cases} C_x^n p^x q^{n-x} & ; \quad \text{for } x=0,1,2,\dots,n \\ 0 & ; \quad \text{otherwise} \end{cases}$$

$$P(X=x)=\begin{cases} C_x^6 (0.15)^x (0.85)^{6-x} & ; \quad \text{for } x=0,1,2,3,4,5,6 \\ 0 & ; \quad \text{otherwise} \end{cases}$$

The probabilities of all values of X are:

$$P(X=0)= C_0^6 (0.15)^0 (0.85)^{6-0} = 0.37715$$

$$P(X=1)= C_1^6 (0.15)^1 (0.85)^{6-1} = 0.39933$$

$$P(X=2)= C_2^6 (0.15)^2 (0.85)^{6-2} = 0.17618$$

$$P(X=3)= C_3^6 (0.15)^3 (0.85)^{6-3} = 0.04145$$

$$P(X=4)= C_4^6 (0.15)^4 (0.85)^{6-4} = 0.00549$$


$$P(X=5)= C_5^6 (0.15)^5 (0.85)^{6-5} = 0.00039$$

$$P(X=6)= C_6^6 (0.15)^6 (0.85)^{6-6} = 0.00001$$

The probability distribution of X can be presented by the following table:

x	$P(X = x)$
0	0.37715
1	0.39933
2	0.17618
3	0.04145
4	0.00549
5	0.00039
6	0.00001
Total	1

The probability distribution of X can be presented by the following graph:



2. The **mean** of the distribution (the **expected number** of men out of 6 with high blood pressure) is:

$$\mu = np = 6 \times 0.15 = 0.9$$

3. The **variance** is:

$$\sigma^2 = npq = 6 \times 0.15 \times 0.85 = 0.765$$

The **standard deviation**:

$$\sigma = \sqrt{npq} = \sqrt{0.765}$$

4. The probability that there will be **exactly 2 men** with high blood pressure is:

$$P(X = 2) = 0.17618$$

$$P(X = 2) = C_2^6 (0.15)^2 (0.85)^{6-2}$$

x	P(X=x) $C_x^6 (0.15)^x (0.85)^{6-x}$
0	0.37715
1	0.39933
2	0.17618
3	0.04145
4	0.00549
5	0.00039
6	0.00001
Total	1

5. The probability that there will be **at most 2 men** with highblood pressure is:

$$\begin{aligned}
 P(X \leq 2) &= P(X = 0) + P(X = 1) + P(X = 2) \\
 &= 0.37715 + 0.39933 + 0.17618 \\
 &= 0.95266
 \end{aligned}$$

By calculator :

$$P(X \leq 2) = \sum_{x=0}^{x=2} (C_x^6) (0.15)^x (0.85)^{6-x}$$

6. The probability that there will be **at least 4 men** with highblood pressure is:

$$\begin{aligned}
 P(X \geq 4) &= P(X = 4) + P(X = 5) + P(X = 6) \\
 &= 0.00549 + 0.00039 + 0.00001 \\
 &= 0.00589
 \end{aligned}$$

By calculator :

$$P(X \geq 4) = \sum_{x=4}^{x=6} (C_x^6) (0.15)^x (0.85)^{6-x}$$

x	P(X=x)
	$C_x^6 (0.15)^x (0.85)^{6-x}$
0	0.37715
1	0.39933
2	0.17618
3	0.04145
4	0.00549
5	0.00039
6	0.00001
Total	1

Example: (Reading Assignment)

Suppose that **25% of the people in a certain population have low hemoglobin levels**. The experiment is to **choose 5 people at random from this population**.

Let the discrete random variable X be the number of people out of 5 with low hemoglobin levels .

- Find the probability distribution of X .
- Find the probability that **at least 2** people have low hemoglobin levels.
- Find the probability that **at most 3** people have low hemoglobin levels.
- Find the **expected number** of people with low hemoglobin levels out of the 5 people.
- Find the **variance** of the number of people with low hemoglobin levels out of the 5 people.

Solution:

X = the number of people out of 5 with low hemoglobin levels
The Bernoulli trial is the process of diagnosing the person

Success = the person has low hemoglobin

Failure = the person does not have low hemoglobin

$$\begin{aligned}
 n &= 5 && \text{(no. of trials)} \\
 p &= 0.25 && \text{(probability of success)} \\
 q &= 1 - p = 0.75 && \text{(probability of failure)}
 \end{aligned}$$

a) X has a binomial distribution with parameter $n = 5$ and $p = 0.25$

$$X \sim \text{Binomial}(5, 0.25)$$

The possible values of X are :

$$x = 0, 1, 2, 3, 4, 5$$

The probability distribution is:

$$P(X=x) = \begin{cases} C_x^5 (0.25)^x (0.75)^{5-x} & ; \text{ for } x=0,1,2,3,4,5 \\ 0 & ; \text{ otherwise} \end{cases}$$

x	P(X = x)
0	$C_0^5 (0.25)^0 (0.75)^{5-0} = 0.23730$
1	$C_1^5 (0.25)^1 (0.75)^{5-1} = 0.39551$
2	$C_2^5 (0.25)^2 (0.75)^{5-2} = 0.26367$
3	$C_3^5 (0.25)^3 (0.75)^{5-3} = 0.08789$
4	$C_4^5 (0.25)^4 (0.75)^{5-4} = 0.01465$
5	$C_5^5 (0.25)^5 (0.75)^{5-5} = 0.00098$
Total	1

b) The probability that at least 2 people have low hemoglobin levels:

$$\begin{aligned}
 P(X \geq 2) &= P(X = 2) + P(X = 3) + P(X = 4) + P(X = 5) \\
 &= 0.26367 + 0.08789 + 0.01465 + 0.00098 \\
 &= 0.036719
 \end{aligned}$$

By calculator :

$$P(X \geq 2) = \sum_{x=2}^5 (C_x^5) (0.25)^x (0.75)^{5-x}$$

c) The probability that at most 3 people have low hemoglobin levels:

$$\begin{aligned}
 P(X \leq 3) &= P(X = 0) + P(X = 1) + P(X = 2) + P(X = 3) \\
 &= 0.23730 + 0.39551 + 0.26367 + 0.08789 \\
 &= 0.98437
 \end{aligned}$$

By calculator :

$$P(X \leq 3) = \sum_{x=0}^3 (C_x^5) (0.25)^x (0.75)^{5-x}$$

d) The **expected number** of people with low hemoglobin levels out of the 5 people (**the mean of X**):

$$\mu = np = 5 \times 0.25 = 1.25$$

e) The **variance of the number** of people with low hemoglobin levels out of the 5 people (the variance of X) is:

$$\sigma^2 = npq = 5 \times 0.25 \times 0.75 = 0.9375$$

The standard deviation:

$$\sigma = \sqrt{npq} = \sqrt{0.9375}$$

4.4 The Poisson Distribution:

- It is a discrete distribution.
- The Poisson distribution is used to model a discrete r.v representing the number of occurrences of some random event in an interval of time or space (or some volume of matter).
- The possible values of X are:

$$x = 0, 1, 2, 3, \dots$$

- The discrete r. v. X is said to have a **Poisson distribution with parameter (average or mean) λ** if the probability distribution of X is given by :

$$P(X = x) = \begin{cases} \frac{e^{-\lambda} \lambda^x}{x!} & ; \quad \text{for } x = 0, 1, 2, \dots \\ 0 & ; \quad \text{otherwise} \end{cases}$$

Where $e = 2.71828$

We write :

$$X \sim \text{Poisson}(\lambda)$$

Result: (Mean and Variance of Poisson distribution)

If $X \sim \text{Poisson}(\lambda)$, then:

1-The mean (expected value/ average):

$$\mu = \lambda$$

2-The variance:

$$\sigma^2 = \lambda$$

3- The standard deviation:

$$\sigma = \sqrt{\lambda}$$

Example :

Some random quantities that can be modeled by Poisson distribution:

- No. of patients in a waiting room in an hours.
- No. of surgeries performed in a month.
- No. of rats in each house in a particular city.

Note:

- λ is the average (mean) of the distribution.
- If X = The number of patients seen in the emergency unit **in a day**, and if $X \sim \text{Poisson}(\lambda)$ then :

- 1) The average (mean) of patients seen every **day** in the emergency unit $= \lambda$.
- 2) The average (mean) of patients seen every **month** in the emergency unit $= 30\lambda$.
- 3) The average (mean) of patients seen every **year** in the emergency unit $= 360\lambda$.
- 4) The average (mean) of patients seen every **hour** in the emergency unit $= \lambda/24$.

Also, notice that:

2) Y = The number of patients seen **every month**, then:

$$Y \sim \text{Poisson}(\lambda^*),$$

$$\text{where } \lambda^* = 30 \lambda$$

3) W = The number of patients seen **every year**, then:

$$W \sim \text{Poisson}(\lambda^*),$$

$$\text{where } \lambda^* = 360 \lambda$$

4) V = The number of patients seen **every hour**, then:

$$V \sim \text{Poisson}(\lambda^*),$$

$$\text{where } \lambda^* = \lambda/24$$

Example:

Suppose that the number of snake bites cases seen at KKUH in **year** has a **Poisson distribution** with **average 6 bite cases**.

- (1) What is the probability that in a year:
 - (i) The no. of snake bite cases will be 7 ?
 - (ii) The no. of snake bite cases will be less than 2 ?
- (2) What is the probability that there will be 10 snake bite cases in 2 years?
- (3) What is the probability that there will be no snake bite cases in a month?
- (4) Find the probability that there will be more than or equal one snake bite cases in a month?
- (5) The mean of snake bite cases in a year.
- (6) The variance of snake bite cases in a month.

Solution:

X = no. of snake bite cases in a year.

$X \sim \text{Poisson } (\lambda = 6)$

year

$x = 0, 1, 2, \dots$

Note :The Poisson Distribution

$$P(X = x) = \frac{e^{-\lambda} \lambda^x}{x!}$$

(1) What is the probability that in a year:

$$(\lambda = 6)$$

- (i) The no. of snake bite cases will be 7 ?
- (ii) The no. of snake bite cases will be less than 2

$$i. \quad P(X = 7) = \frac{e^{-\lambda} \lambda^x}{x!} = \frac{e^{-6} 6^7}{7!} = 0.13678$$

$$ii. \quad P(X < 2) = \frac{e^{-6} 6^0}{0!} + \frac{e^{-6} 6^1}{1!} = 0.00248 + 0.01487 = 0.01735$$

(2) What is the probability that there will be 10 snake bite cases in 2 years?

$$(\lambda^* = 2\lambda = 12)$$

Y = no of snake bite cases in 2 years

$$Y \sim \text{Poisson} (\lambda^* = 12)$$

$$(\lambda^* = 2\lambda = 6 \times 2 = 12)$$

$$P(Y = y) = \frac{e^{-12} 12^y}{y!} \quad y = 0, 1, 2, \dots$$

$$P(Y = 10) = \frac{e^{-12} 12^{10}}{10!} = 0.1048$$

Note :The Poisson Distribution

$$P(X = x) = \frac{e^{-\lambda} \lambda^x}{x!}$$

(3) What is the probability that there will be no snake bite cases in a month

W= no of snake bite cases in a month

$$W \sim \text{Poisson} (\lambda^* = 0.5) \quad (\lambda^* = \frac{\lambda}{12} = \frac{6}{12} = 0.5)$$

$$P(W = w) = \frac{e^{-0.5} 0.5^w}{w!} \quad w = 0, 1, 2, \dots$$

$$P(W = 0) = \frac{e^{-0.5} 0.5^0}{0!} = 0.6065$$

$$\left(\lambda^* = \frac{\lambda}{12} = 0.5 \right)$$

(4) Find the probability that there will be more than or equal one snake bite cases in a month

$$P(W \geq 1) = P(W = 1) + P(W = 2) + P(W = 3) + \dots \quad (\text{The value of } x = 0, 1, 2, 3, 4, 5, \dots)$$

$$= 1 - P(W < 1)$$

$$= 1 - P(W = 0)$$

$$= 1 - \frac{e^{-0.5} 0.5^0}{0!}$$

$$= 1 - 0.6065 = 0.3935$$

Note :The Poisson Distribution

$$P(X = x) = \frac{e^{-\lambda} \lambda^x}{x!}$$

(5) The mean of snake bite cases in a year $\longrightarrow$ $(\lambda = 6)$

$$\mu = \lambda = 6$$

(6) The variance of snake bite cases in a month $\longrightarrow$ $\left(\lambda^* = \frac{\lambda}{12} = 0.5 \right)$

$$\sigma^2 = \lambda^* = \frac{\lambda}{12} = \frac{6}{12} = 0.5$$

(7) The standard deviation of snake bite cases in 2 years $\longrightarrow$ $(\lambda^* = 2\lambda = 12)$

$$\sigma = \sqrt{\lambda^*} = \sqrt{12} = 3.4641$$

Note :The Poisson Distribution

$$\begin{aligned}\mu &= \lambda \\ \sigma^2 &= \lambda \\ \sigma &= \sqrt{\lambda}\end{aligned}$$

(8) Find the probability that there will be more than 3 snake bite cases in 2 years :

$$(\lambda^* = 2\lambda = 12)$$

$$\begin{aligned} P(X > 3) &= 1 - P(X \leq 3) \\ &= 1 - [P(X = 0) + P(X = 1) + P(X = 2) + P(X = 3)] \\ &= 1 - 0.0023 \\ &= 0.9977 \end{aligned}$$

By calculator :

$$\begin{aligned} P(X > 3) &= 1 - P(X \leq 3) \\ &= 1 - \sum_{x=0}^{x=3} \left(\frac{e^{-12} \times 12^x}{x!} \right) \\ &= 0.9977 \end{aligned}$$

Note :The Poisson Distribution


$$P(X = x) = \frac{e^{-\lambda} \lambda^x}{x!}$$

	Discrete Distribution	
	Binomial	Poisson
parameters	n = trials P = probability of success	λ = average or mean
r.v	X = number of successes in n trials	X = number of occurrence of some event in an interval of time or space
Value of r.v	x = 0, 1, ..., n	x = 0, 1, 2,
write	$X \sim \text{Binomial}(n, p)$	$X \sim \text{Poisson}(\lambda)$
Probability distribution P(X=x)	$\binom{n}{x} p^x q^{n-x}$	$\frac{\lambda^x e^{-\lambda}}{x!}$
Mean Expected μ	np	λ
Variance σ^2	npq	λ
standard deviation σ	$\sqrt{npq}$	$\sqrt{\lambda}$

4.5 Continuous Probability Distributions:


For any continuous r.v. X , there exists a function $f(x)$, called the probability density function (pdf) of X , for which:

1. The total area under the curve of $f(x)$ equals to 1.




$$\text{Total area} = \int_{-\infty}^{\infty} f(x) dx = 1$$

2. The probability that X is between the points (a) and (b) equals to the area under the curve of $f(x)$ which is bounded by the point a and b.




$$P(a \leq X \leq b) = \int_a^b f(x) dx = \text{area}$$


3. In general, the probability of an interval event is given by the area under the curve of $f(x)$ and above that interval.



$$P(a \leq X \leq b) = \int_a^b f(x) dx = \text{area}$$



$$P(X \leq a) = \int_{-\infty}^a f(x) dx = \text{area}$$



$$P(X \geq b) = \int_b^{\infty} f(x) dx = \text{area}$$

Note:

If X is continuous r.v. then:


1. $P(X = a) = 0$ for any (a)
2. $P(X \leq a) = P(X < a)$
3. $P(X \geq b) = P(X > b)$

$$4. \quad P(a \leq X \leq b) = P(a \leq X < b) = P(a < X \leq b) = P(a < X < b)$$

$$5. \quad P(X \leq x) = \text{cumulative probability}$$


$$6. \quad P(X \geq a) = 1 - P(X < a) = 1 - P(X \leq a)$$

$$7. \quad P(a \leq X \leq b) = P(X \leq b) - P(X \leq a)$$



$$P(X \geq a) = 1 - P(X \leq a)$$

$$A = 1 - B, \quad \text{Total area} = 1$$



$$P(a \leq X \leq b) = P(X \leq b) - P(X \leq a)$$

$$\int_a^b f(x) dx = \int_{-\infty}^b f(x) dx - \int_{-\infty}^a f(x) dx$$

4.6 The Normal Distribution:

- One of the most important continuous distributions.
- Many measurable characteristics are normally or approximately normally distributed. (Examples: height, weight, ...)
- The probability density function of the normal distribution is given by:

$$f(x) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2} \quad ; \quad -\infty < x < \infty$$

where ($e = 2.71828$) and ($\pi = 3.14159$)



Mean = μ
Standard deviation = σ
Variance = σ^2

- The parameters of the distribution are the mean (μ) and the standard deviation (σ).

The continuous r.v. X which has a normal distribution has several important characteristics:

1. $-\infty < X < \infty$
2. The density function of X , $f(x)$, has a bell-Shaped curve:
3. The highest point of the curve of $f(x)$ at the mean (μ). (Mode= μ)
4. The curve of $f(x)$ is symmetric about the mean (μ).

Mean = Median = Mode = μ



5. The normal distribution **depends on two parameters** :

Mean = μ (determines the location)

Standard deviation = σ (variance σ^2) (determines the shape)

6. If the r.v. **X** is normally distributed with mean μ and standard deviation σ (variance σ^2), we write:

$X \sim \text{Normal}(\text{mean} = \mu, \text{varaince} = \sigma^2)$ **or** $X \sim N(\text{mean} = \mu, \text{varaince} = \sigma^2)$


7. The location of the normal distribution depends on μ and

The shape of the normal distribution depends on σ .


Note : The location of the normal distribution depends on μ and its shape depends on σ

Suppose we have two normal distributions:


———— $N(\mu_1, \sigma_1)$
 - - - - - $N(\mu_2, \sigma_2)$



$$\mu_1 < \mu_2, \sigma_1 = \sigma_2$$



$$\mu_1 = \mu_2, \sigma_1 < \sigma_2$$



$$\mu_1 < \mu_2, \sigma_1 < \sigma_2$$

The Standard Normal Distribution (Z):

The normal distribution with **mean** $\mu = 0$ and **variance** $\sigma^2 = 1$ is called **standard normal distribution (Z)** and is denoted by :


Normal(0 , 1) **or** N(0 , 1) .

The standard normal random variable is denoted by (Z) , and we write :




$$Z \sim N(0,1)$$

The probability density function (pdf) of $Z \sim N(0,1)$, is given by :

$$f(z) = \frac{1}{\sqrt{2\pi}} e^{-\frac{z^2}{2}} \quad , \quad -\infty < \mathbf{Z} < \infty$$




The standard normal distribution, Normal (0,1), is very important because probabilities of any normal distribution can be calculated from the probabilities of the standard normal distribution.



Result : (convert a normal distribution to a standard normal distribution)

$$\text{If } X \sim N(\mu, \sigma^2) , \text{ then } Z = \frac{x - \mu}{\sigma} \sim N(0,1)$$

$$Z = \frac{x - \text{mean}}{\text{Standard deviation}}$$



Calculating Probabilities of Normal (0,1): (Table Z)

For the standard normal distribution $Z \sim N(0,1)$, there is **a special table used to calculate probabilities** of the form:


$$P(Z \leq a)$$

$$1. P(Z \leq a) = \text{From the table}$$


$$2. P(Z \geq b) = 1 - P(Z \leq b)$$

$$3. P(a \leq Z \leq b) = P(Z \leq b) - P(Z \leq a)$$


$$4. P(Z = a) = 0 \text{ for every } (a)$$



$$P(Z \leq a) = \text{From the table}$$




$$P(Z \geq b) = 1 - P(Z \leq b)$$



$$P(a \leq Z \leq b) = P(Z \leq b) - P(Z \leq a)$$

Standard Normal Table

Areas Under the Standard Normal Curve



z	-0.09	-0.08	-0.07	-0.06	-0.05	-0.04	-0.03	-0.02	-0.01	-0.00	z
-3.50	0.00017	0.00017	0.00016	0.00015	0.00015	0.00014	0.00013	0.00012	0.00011	0.00010	-3.50
-3.40	0.00024	0.00025	0.00026	0.00027	0.00028	0.00029	0.00030	0.00031	0.00032	0.00034	-3.40
-3.30	0.00035	0.00036	0.00038	0.00039	0.00040	0.00042	0.00043	0.00045	0.00047	0.00048	-3.30
-3.20	0.00050	0.00052	0.00054	0.00056	0.00058	0.00060	0.00062	0.00064	0.00066	0.00069	-3.20
-3.10	0.00071	0.00074	0.00076	0.00079	0.00082	0.00084	0.00087	0.00090	0.00094	0.00097	-3.10
-3.00	0.00100	0.00104	0.00107	0.00111	0.00114	0.00118	0.00122	0.00126	0.00131	0.00135	-3.00
-2.90	0.00139	0.00144	0.00149	0.00154	0.00159	0.00164	0.00169	0.00175	0.00181	0.00187	-2.90
-2.80	0.00193	0.00199	0.00205	0.00212	0.00219	0.00226	0.00233	0.00240	0.00248	0.00256	-2.80
-2.70	0.00264	0.00272	0.00280	0.00289	0.00298	0.00307	0.00317	0.00326	0.00336	0.00347	-2.70
-2.60	0.00357	0.00368	0.00379	0.00391	0.00402	0.00415	0.00427	0.00440	0.00453	0.00466	-2.60
-2.50	0.00480	0.00494	0.00508	0.00523	0.00539	0.00554	0.00570	0.00587	0.00604	0.00621	-2.50
-2.40	0.00639	0.00657	0.00676	0.00695	0.00714	0.00734	0.00755	0.00776	0.00798	0.00820	-2.40
-2.30	0.00842	0.00866	0.00889	0.00914	0.00939	0.00964	0.00990	0.01017	0.01044	0.01072	-2.30
-2.20	0.01101	0.01130	0.01160	0.01191	0.01222	0.01255	0.01287	0.01321	0.01355	0.01390	-2.20
-2.10	0.01426	0.01463	0.01500	0.01539	0.01578	0.01618	0.01659	0.01700	0.01743	0.01786	-2.10
-2.00	0.01831	0.01876	0.01923	0.01970	0.02018	0.02068	0.02118	0.02169	0.02222	0.02275	-2.00
-1.90	0.02330	0.02385	0.02442	0.02500	0.02559	0.02619	0.02680	0.02743	0.02807	0.02872	-1.90
-1.80	0.02938	0.03005	0.03074	0.03144	0.03216	0.03288	0.03362	0.03438	0.03515	0.03593	-1.80
-1.70	0.03673	0.03754	0.03836	0.03920	0.04006	0.04093	0.04182	0.04272	0.04363	0.04457	-1.70
-1.60	0.04551	0.04648	0.04746	0.04846	0.04947	0.05050	0.05155	0.05262	0.05370	0.05480	-1.60
-1.50	0.05592	0.05705	0.05821	0.05938	0.06057	0.06178	0.06301	0.06426	0.06552	0.06681	-1.50
-1.40	0.06811	0.06944	0.07078	0.07213	0.07353	0.07493	0.07636	0.07780	0.07927	0.08076	-1.40
-1.30	0.08226	0.08379	0.08534	0.08691	0.08851	0.09012	0.09176	0.09342	0.09510	0.09680	-1.30
-1.20	0.09853	0.10027	0.10204	0.10383	0.10565	0.10749	0.10935	0.11123	0.11314	0.11507	-1.20
-1.10	0.11702	0.11900	0.12100	0.12302	0.12507	0.12714	0.12924	0.13136	0.13350	0.13567	-1.10
-1.00	0.13786	0.14007	0.14231	0.14457	0.14686	0.14917	0.15151	0.15386	0.15625	0.15866	-1.00
-0.90	0.16109	0.16354	0.16602	0.16853	0.17106	0.17361	0.17619	0.17879	0.18141	0.18406	-0.90
-0.80	0.18673	0.18943	0.19215	0.19489	0.19766	0.20045	0.20327	0.20611	0.20897	0.21186	-0.80
-0.70	0.21476	0.21770	0.22065	0.22363	0.22663	0.22965	0.23270	0.23576	0.23885	0.24196	-0.70
-0.60	0.24510	0.24825	0.25143	0.25463	0.25785	0.26109	0.26435	0.26763	0.27093	0.27425	-0.60
-0.50	0.27760	0.28096	0.28434	0.28774	0.29116	0.29460	0.29806	0.30153	0.30503	0.30854	-0.50
-0.40	0.31207	0.31561	0.31918	0.32276	0.32636	0.32997	0.33360	0.33724	0.3409	0.34458	-0.40
-0.30	0.34827	0.35197	0.35569	0.35942	0.36317	0.36693	0.37070	0.37448	0.37828	0.38209	-0.30
-0.20	0.38591	0.38974	0.39358	0.39743	0.40129	0.40517	0.40905	0.41294	0.41683	0.42074	-0.20
-0.10	0.42465	0.42858	0.43251	0.43644	0.44038	0.44433	0.44828	0.45224	0.45620	0.46017	-0.10
-0.00	0.46414	0.46812	0.47210	0.47608	0.48006	0.48405	0.48803	0.49202	0.49601	0.50000	-0.00

Value of

$P(Z \leq z)$

Value of z

$$P(Z < \boxed{\text{أطراف الجدول}}) = \boxed{\text{داخل الجدول}}$$

Example: Suppose that $Z \sim N(0,1)$

(1) $P(Z \leq 1.50) = 0.93319$

z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09	z
0.00	0.50000	0.50399	0.50798	0.51197	0.51595	0.51994	0.52392	0.52790	0.53188	0.53586	0.00
0.10	0.53983	0.54380	0.54776	0.55172	0.55567	0.55962	0.56356	0.56749	0.57142	0.57535	0.10
0.20	0.57926	0.58317	0.58706	0.59095	0.59483	0.59871	0.60257	0.60642	0.61026	0.61409	0.20
0.30	0.61791	0.62172	0.62552	0.62930	0.63307	0.63683	0.64058	0.64431	0.64803	0.65173	0.30
0.40	0.65542	0.65910	0.66276	0.66640	0.67003	0.67364	0.67724	0.68082	0.68439	0.68793	0.40
0.50	0.69146	0.69497	0.69847	0.70194	0.70540	0.70884	0.71226	0.71566	0.71904	0.72240	0.50
0.60	0.72575	0.72907	0.73237	0.73565	0.73891	0.74215	0.74537	0.74857	0.75175	0.75490	0.60
0.70	0.75804	0.76115	0.76424	0.76730	0.77035	0.77337	0.77637	0.77935	0.78230	0.78524	0.70
0.80	0.78814	0.79103	0.79389	0.79673	0.79955	0.80234	0.80511	0.80785	0.81057	0.81327	0.80
0.90	0.81594	0.81859	0.82121	0.82381	0.82639	0.82894	0.83147	0.83398	0.83646	0.83891	0.90
1.00	0.84134	0.84375	0.84614	0.84849	0.85083	0.85314	0.85543	0.85769	0.85993	0.86214	1.00
1.10	0.86433	0.86650	0.86864	0.87076	0.87286	0.87493	0.87698	0.87900	0.88100	0.88298	1.10
1.20	0.88493	0.88686	0.88877	0.89065	0.89251	0.89435	0.89617	0.89796	0.89973	0.90147	1.20
1.30	0.90320	0.90490	0.90658	0.90824	0.90988	0.91149	0.91309	0.91466	0.91621	0.91774	1.30
1.40	0.91924	0.92073	0.92220	0.92364	0.92507	0.92647	0.92785	0.92922	0.93056	0.93189	1.40
1.50	0.93319	0.93448	0.93574	0.93699	0.93822	0.93943	0.94062	0.94179	0.94295	0.94408	1.50
1.60	0.94520	0.94630	0.94738	0.94845	0.94950	0.95053	0.95154	0.95254	0.95352	0.95449	1.60

Example: Suppose that $Z \sim N(0,1)$

(2) $P(Z \geq 0.98) = 1 - P(Z \leq 0.98)$
 $= 1 - 0.83646$
 $= 0.1635$

z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09	z
0.00	0.50000	0.50399	0.50798	0.51197	0.51595	0.51994	0.52392	0.52790	0.53188	0.53586	0.00
0.10	0.53983	0.54380	0.54776	0.55172	0.55567	0.55962	0.56356	0.56749	0.57142	0.57535	0.10
0.20	0.57926	0.58317	0.58706	0.59095	0.59483	0.59871	0.60257	0.60642	0.61026	0.61409	0.20
0.30	0.61791	0.62172	0.62552	0.62930	0.63307	0.63683	0.64058	0.64431	0.64803	0.65173	0.30
0.40	0.65542	0.65910	0.66276	0.66640	0.67003	0.67364	0.67724	0.68082	0.68439	0.68793	0.40
0.50	0.69146	0.69497	0.69847	0.70194	0.70540	0.70884	0.71226	0.71566	0.71904	0.72240	0.50
0.60	0.72575	0.72907	0.73237	0.73565	0.73891	0.74215	0.74537	0.74857	0.75175	0.75490	0.60
0.70	0.75804	0.76115	0.76424	0.76730	0.77035	0.77337	0.77637	0.77935	0.78230	0.78524	0.70
0.80	0.78814	0.79103	0.79389	0.79673	0.79955	0.80234	0.80511	0.80785	0.81057	0.81327	0.80
0.90	0.81594	0.81859	0.82121	0.82381	0.82639	0.82894	0.83147	0.83398	0.83646	0.83891	0.90
1.00	0.84134	0.84375	0.84614	0.84849	0.85083	0.85314	0.85543	0.85769	0.85993	0.86214	1.00
1.10	0.86433	0.86650	0.86864	0.87076	0.87286	0.87493	0.87698	0.87900	0.88100	0.88298	1.10
1.20	0.88493	0.88686	0.88877	0.89065	0.89251	0.89435	0.89617	0.89796	0.89973	0.90147	1.20
1.30	0.90320	0.90490	0.90658	0.90824	0.90988	0.91149	0.91309	0.91466	0.91621	0.91774	1.30
1.40	0.91924	0.92073	0.92220	0.92364	0.92507	0.92647	0.92785	0.92922	0.93056	0.93189	1.40
1.50	0.93319	0.93448	0.93574	0.93699	0.93822	0.93943	0.94062	0.94179	0.94295	0.94408	1.50
1.60	0.94520	0.94630	0.94738	0.94845	0.94950	0.95053	0.95154	0.95254	0.95352	0.95449	1.60

Example: Suppose that $Z \sim N(0,1)$

(3) $P(-1.33 \leq Z \leq 2.42)$

$= P(Z \leq 2.42) - P(Z \leq -1.33)$

$= 0.99224 - 0.09176$


$= 0.9004$

z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09	z
0.00	0.50000	0.50399	0.50798	0.51197	0.51595	0.51994	0.52392	0.52790	0.53188	0.53586	0.00
0.10	0.53983	0.54380	0.54776	0.55172	0.55567	0.55962	0.56356	0.56749	0.57142	0.57535	0.10
0.20	0.57926	0.58317	0.58706	0.59095	0.59483	0.59871	0.60257	0.60642	0.61026	0.61409	0.20
0.30	0.61791	0.62172	0.62552	0.62930	0.63307	0.63683	0.64058	0.64431	0.64803	0.65173	0.30
0.40	0.65542	0.65910	0.66276	0.66640	0.67003	0.67364	0.67724	0.68082	0.68439	0.68793	0.40
0.50	0.69146	0.69497	0.69847	0.70194	0.70540	0.70884	0.71226	0.71566	0.71904	0.72240	0.50
0.60	0.72575	0.72907	0.73237	0.73565	0.73891	0.74215	0.74537	0.74857	0.75175	0.75490	0.60
0.70	0.75804	0.76115	0.76424	0.76730	0.77035	0.77337	0.77637	0.77935	0.78230	0.78524	0.70
0.80	0.78814	0.79103	0.79389	0.79673	0.79955	0.80234	0.80511	0.80785	0.81057	0.81327	0.80
0.90	0.81594	0.81859	0.82121	0.82381	0.82639	0.82894	0.83147	0.83398	0.83646	0.83891	0.90
1.00	0.84134	0.84375	0.84614	0.84849	0.85083	0.85314	0.85543	0.85769	0.85993	0.86214	1.00
1.10	0.86433	0.86650	0.86864	0.87076	0.87286	0.87493	0.87698	0.87900	0.88100	0.88298	1.10
1.20	0.88493	0.88686	0.88877	0.89065	0.89251	0.89435	0.89617	0.89796	0.89973	0.90147	1.20
1.30	0.90320	0.90490	0.90658	0.90824	0.90988	0.91149	0.91309	0.91466	0.91621	0.91774	1.30
1.40	0.91924	0.92073	0.92220	0.92364	0.92507	0.92647	0.92785	0.92922	0.93056	0.93189	1.40
1.50	0.93319	0.93448	0.93574	0.93699	0.93822	0.93943	0.94062	0.94179	0.94295	0.94408	1.50
1.60	0.94520	0.94630	0.94738	0.94845	0.94950	0.95053	0.95154	0.95254	0.95352	0.95449	1.60
1.70	0.95543	0.95637	0.95728	0.95818	0.95907	0.95994	0.96080	0.96164	0.96246	0.96327	1.70
1.80	0.96407	0.96485	0.96562	0.96638	0.96712	0.96784	0.96856	0.96926	0.96995	0.97062	1.80
1.90	0.97128	0.97193	0.97257	0.97320	0.97381	0.97441	0.97500	0.97558	0.97615	0.97670	1.90
2.00	0.97725	0.97778	0.97831	0.97882	0.97932	0.97982	0.98030	0.98077	0.98124	0.98169	2.00
2.10	0.98214	0.98257	0.98300	0.98341	0.98382	0.98422	0.98461	0.98500	0.98537	0.98574	2.10
2.20	0.98610	0.98645	0.98679	0.98713	0.98745	0.98778	0.98809	0.98840	0.98870	0.98899	2.20
2.30	0.98928	0.98956	0.98983	0.99010	0.99036	0.99061	0.99086	0.99111	0.99134	0.99158	2.30
2.40	0.99180	0.99207	0.99224	0.99245	0.99266	0.99286	0.99305	0.99324	0.99343	0.99361	2.40
2.50	0.99379	0.99396	0.99413	0.99430	0.99446	0.99461	0.99477	0.99492	0.99506	0.99520	2.50

z	-0.09	-0.08	-0.07	-0.06	-0.05	-0.04	-0.03	-0.02	-0.01	-0.00	z
-3.50	0.00017	0.00017	0.00018	0.00019	0.00019	0.00020	0.00021	0.00022	0.00022	0.00023	-3.50
-3.40	0.00024	0.00025	0.00026	0.00027	0.00028	0.00029	0.00030	0.00031	0.00032	0.00034	-3.40
-3.30	0.00035	0.00036	0.00038	0.00039	0.00040	0.00042	0.00043	0.00045	0.00047	0.00048	-3.30
-3.20	0.00050	0.00052	0.00054	0.00056	0.00058	0.00060	0.00062	0.00064	0.00066	0.00069	-3.20
-3.10	0.00071	0.00074	0.00076	0.00079	0.00082	0.00084	0.00087	0.00090	0.00094	0.00097	-3.10
-3.00	0.00100	0.00104	0.00107	0.00111	0.00114	0.00118	0.00122	0.00126	0.00131	0.00135	-3.00
-2.90	0.00139	0.00144	0.00149	0.00154	0.00159	0.00164	0.00169	0.00175	0.00181	0.00187	-2.90
-2.80	0.00193	0.00199	0.00205	0.00212	0.00219	0.00226	0.00233	0.00240	0.00248	0.00256	-2.80
-2.70	0.00264	0.00272	0.00280	0.00289	0.00298	0.00307	0.00317	0.00326	0.00336	0.00347	-2.70
-2.60	0.00357	0.00368	0.00379	0.00391	0.00402	0.00415	0.00427	0.00440	0.00453	0.00466	-2.60
-2.50	0.00480	0.00494	0.00508	0.00523	0.00539	0.00554	0.00570	0.00587	0.00604	0.00621	-2.50
-2.40	0.00639	0.00657	0.00676	0.00695	0.00714	0.00734	0.00755	0.00776	0.00798	0.00820	-2.40
-2.30	0.00842	0.00866	0.00889	0.00914	0.00939	0.00964	0.00990	0.01017	0.01044	0.01072	-2.30
-2.20	0.01101	0.01130	0.01160	0.01191	0.01222	0.01255	0.01287	0.01321	0.01355	0.01390	-2.20
-2.10	0.01426	0.01463	0.01500	0.01539	0.01578	0.01618	0.01659	0.01700	0.01743	0.01786	-2.10
-2.00	0.01831	0.01876	0.01923	0.01970	0.02018	0.02068	0.02118	0.02169	0.02222	0.02275	-2.00
-1.90	0.02330	0.02385	0.02442	0.02500	0.02559	0.02619	0.02680	0.02743	0.02807	0.02872	-1.90
-1.80	0.02938	0.03005	0.03074	0.03144	0.03216	0.03288	0.03362	0.03438	0.03515	0.03593	-1.80
-1.70	0.03673	0.03754	0.03836	0.03920	0.04006	0.04093	0.04182	0.04272	0.04363	0.04457	-1.70
-1.60	0.04551	0.04648	0.04746	0.04846	0.04947	0.05050	0.05155	0.05262	0.05370	0.05480	-1.60
-1.50	0.05592	0.05705	0.05821	0.05938	0.06057	0.06178	0.06301	0.06426	0.06552	0.06681	-1.50
-1.40	0.06811	0.06944	0.07078	0.07215	0.07353	0.07493	0.07636	0.07780	0.07927	0.08076	-1.40
-1.30	0.08226	0.08379	0.08534	0.08691	0.08851	0.08912	0.09176	0.09342	0.09510	0.09680	-1.30
-1.20	0.09853	0.10027	0.10204	0.10383	0.10565	0.10749	0.10935	0.11123	0.11314	0.11507	-1.20
-1.10	0.11702	0.11900	0.12100	0.12302	0.12507	0.12714	0.12924	0.13136	0.13350	0.13567	-1.10

Example: Suppose that $Z \sim N(0,1)$

(4) $P(Z \leq 0) = 0.5$ and $P(Z \geq 0) = 1 - P(Z < 0) = 1 - 0.5 = 0.5$



z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09	z
0.00	0.50000	0.50399	0.50798	0.51197	0.51595	0.51994	0.52392	0.52790	0.53188	0.53586	0.00
0.10	0.53983	0.54380	0.54776	0.55172	0.55567	0.55962	0.56356	0.56749	0.57142	0.57535	0.10
0.20	0.57926	0.58317	0.58706	0.59095	0.59483	0.59871	0.60257	0.60642	0.61026	0.61409	0.20
0.30	0.61791	0.62172	0.62552	0.62930	0.63307	0.63683	0.64058	0.64431	0.64803	0.65173	0.30
0.40	0.65542	0.65910	0.66276	0.66640	0.67003	0.67364	0.67724	0.68082	0.68439	0.68793	0.40
0.50	0.69146	0.69497	0.69847	0.70194	0.70540	0.70884	0.71226	0.71566	0.71904	0.72240	0.50
0.60	0.72575	0.72907	0.73237	0.73565	0.73891	0.74215	0.74537	0.74857	0.75175	0.75490	0.60
0.70	0.75804	0.76115	0.76424	0.76730	0.77035	0.77337	0.77637	0.77935	0.78230	0.78524	0.70
0.80	0.78814	0.79103	0.79389	0.79673	0.79955	0.80234	0.80511	0.80785	0.81057	0.81327	0.80
0.90	0.81594	0.81859	0.82121	0.82381	0.82639	0.82894	0.83147	0.83398	0.83646	0.83891	0.90


(5) $P(Z = 1.25) = 0$





Notation:

$$Z_A \gg P(Z \leq z) = A$$

• Z_A = The Z-value that leaves an area of A to the left.



For example:



Result:

Since the pdf of $Z \sim N(0,1)$ is **symmetric about 0**, we have:

$$Z_A = -Z_{1-A}$$


For example:

$$Z_{0.35} = -Z_{1-0.35} = -Z_{0.65}$$

$$Z_{0.86} = -Z_{1-0.86} = -Z_{0.14}$$

$$Z_{0.025} = -Z_{1-0.025} = -Z_{0.975} = -1.96$$

$$Z_{0.975} = 1.96$$



Example :

$Z_{0.975} = ??$ $\gggg P(Z \leq z) = 0.975$, Find the value of z ?

0.975 = probability > 0.5 (Use positive table)

z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09	z
0.00	0.50000	0.50399	0.50798	0.51197	0.51595	0.51994	0.52392	0.52790	0.53188	0.53586	0.00
0.10	0.53983	0.54380	0.54776	0.55172	0.55567	0.55962	0.56356	0.56749	0.57142	0.57535	0.10
0.20	0.57926	0.58317	0.58706	0.59095	0.59483	0.59871	0.60257	0.60642	0.61026	0.61409	0.20
0.30	0.61791	0.62172	0.62552	0.62930	0.63307	0.63683	0.64058	0.64431	0.64803	0.65173	0.30
0.40	0.65542	0.65910	0.66276	0.66640	0.67003	0.67364	0.67724	0.68082	0.68439	0.68793	0.40
0.50	0.69146	0.69497	0.69847	0.70194	0.70540	0.70884	0.71226	0.71566	0.71904	0.72240	0.50
0.60	0.72575	0.72907	0.73237	0.73565	0.73891	0.74215	0.74537	0.74857	0.75175	0.75490	0.60
0.70	0.75804	0.76115	0.76424	0.76730	0.77035	0.77337	0.77637	0.77935	0.78230	0.78524	0.70
0.80	0.78814	0.79103	0.79389	0.79673	0.79955	0.80234	0.80511	0.80785	0.81057	0.81327	0.80
0.90	0.81594	0.81859	0.82121	0.82381	0.82639	0.82894	0.83147	0.83398	0.83646	0.83891	0.90
1.00	0.84134	0.84375	0.84614	0.84849	0.85083	0.85314	0.85543	0.85769	0.85993	0.86214	1.00
1.10	0.86433	0.86650	0.86864	0.87076	0.87286	0.87493	0.87698	0.87900	0.88100	0.88298	1.10
1.20	0.88493	0.88686	0.88877	0.89065	0.89251	0.89435	0.89617	0.89796	0.89973	0.90147	1.20
1.30	0.90320	0.90490	0.90658	0.90824	0.90988	0.91149	0.91309	0.91466	0.91621	0.91774	1.30
1.40	0.91924	0.92073	0.92220	0.92364	0.92507	0.92647	0.92785	0.92922	0.93056	0.93189	1.40
1.50	0.93319	0.93448	0.93574	0.93699	0.93822	0.93943	0.94062	0.94179	0.94295	0.94408	1.50
1.60	0.94520	0.94630	0.94738	0.94845	0.94950	0.95053	0.95154	0.95254	0.95352	0.95449	1.60
1.70	0.95543	0.95637	0.95728	0.95818	0.95907	0.95994	0.96080	0.96164	0.96246	0.96327	1.70
1.80	0.96407	0.96485	0.96562	0.96638	0.96712	0.96784	0.96856	0.96926	0.96995	0.97062	1.80
1.90	0.97128	0.97193	0.97257	0.97320	0.97381	0.97441	0.97500	0.97558	0.97615	0.97670	1.90
2.00	0.97725	0.97778	0.97831	0.97882	0.97932	0.97982	0.98030	0.98077	0.98124	0.98169	2.00
2.10	0.98214	0.98257	0.98300	0.98341	0.98382	0.98422	0.98461	0.98500	0.98537	0.98574	2.10

$$Z = 1.90 + 0.06 = 1.96$$

$$Z_{0.975} = 1.96$$

Example :

$Z_{0.025} = ??$ $\gggg P(Z \leq z) = 0.025$, Find the value of z ?

0.025=probability < 0.5 (Use Negative table)

z	-0.09	-0.08	-0.07	-0.06	-0.05	-0.04	-0.03	-0.02	-0.01	-0.00	z
-3.50	0.00017	0.00017	0.00018	0.00019	0.00019	0.00020	0.00021	0.00022	0.00022	0.00023	-3.50
-3.40	0.00024	0.00025	0.00026	0.00027	0.00028	0.00029	0.00030	0.00031	0.00032	0.00034	-3.40
-3.30	0.00035	0.00036	0.00038	0.00039	0.00040	0.00042	0.00043	0.00045	0.00047	0.00048	-3.30
-3.20	0.00050	0.00052	0.00054	0.00056	0.00058	0.00060	0.00062	0.00064	0.00066	0.00069	-3.20
-3.10	0.00071	0.00074	0.00076	0.00079	0.00082	0.00084	0.00087	0.00090	0.00094	0.00097	-3.10
-3.00	0.00100	0.00104	0.00107	0.00111	0.00114	0.00118	0.00122	0.00126	0.00131	0.00135	-3.00
-2.90	0.00139	0.00144	0.00149	0.00154	0.00159	0.00164	0.00169	0.00175	0.00181	0.00187	-2.90
-2.80	0.00193	0.00199	0.00205	0.00212	0.00219	0.00226	0.00233	0.00240	0.00248	0.00256	-2.80
-2.70	0.00264	0.00272	0.00280	0.00289	0.00298	0.00307	0.00317	0.00326	0.00336	0.00347	-2.70
-2.60	0.00357	0.00368	0.00379	0.00391	0.00402	0.00415	0.00427	0.00440	0.00453	0.00466	-2.60
-2.50	0.00480	0.00494	0.00508	0.00523	0.00539	0.00554	0.00570	0.00587	0.00604	0.00621	-2.50
-2.40	0.00639	0.00657	0.00676	0.00695	0.00714	0.00734	0.00755	0.00776	0.00798	0.00820	-2.40
-2.30	0.00842	0.00866	0.00889	0.00914	0.00939	0.00964	0.00990	0.01017	0.01044	0.01072	-2.30
-2.20	0.01101	0.01130	0.01160	0.01191	0.01222	0.01255	0.01287	0.01321	0.01355	0.01390	-2.20
-2.10	0.01426	0.01463	0.01500	0.01539	0.01578	0.01618	0.01659	0.01700	0.01743	0.01786	-2.10
-2.00	0.01831	0.01876	0.01923	0.01970	0.02018	0.02068	0.02118	0.02169	0.02222	0.02275	-2.00
-1.90	0.02330	0.02385	0.02442	0.02500	0.02559	0.02619	0.02680	0.02743	0.02807	0.02872	-1.90
-1.80	0.02938	0.03005	0.03074	0.03144	0.03216	0.03288	0.03362	0.03438	0.03515	0.03593	-1.80
-1.70	0.03673	0.03754	0.03836	0.03920	0.04006	0.04093	0.04182	0.04272	0.04363	0.04457	-1.70
-1.60	0.04551	0.04648	0.04746	0.04846	0.04947	0.05050	0.05155	0.05262	0.05370	0.05480	-1.60

$$Z = (-1.90) + (-0.06) = -1.96$$

$$Z_{0.025} = -1.96$$

Example:

Suppose that $Z \sim N(0,1)$.

If $P(Z < a) = 0.95053 \gggg Z_{0.95053}$

Find the value of a ?? $a = 1.65$

0.95053 =probability > 0.5 (Use positive table)

z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09	z
0.00	0.50000	0.50399	0.50798	0.51197	0.51595	0.51994	0.52392	0.52790	0.53188	0.53586	0.00
0.10	0.53983	0.54380	0.54776	0.55172	0.55567	0.55962	0.56356	0.56749	0.57142	0.57535	0.10
0.20	0.57926	0.58317	0.58706	0.59095	0.59483	0.59871	0.60257	0.60642	0.61026	0.61409	0.20
0.30	0.61791	0.62172	0.62552	0.62930	0.63307	0.63683	0.64058	0.64431	0.64803	0.65173	0.30
0.40	0.65542	0.65910	0.66276	0.66640	0.67003	0.67364	0.67724	0.68082	0.68439	0.68793	0.40
0.50	0.69146	0.69497	0.69847	0.70194	0.70540	0.70884	0.71226	0.71566	0.71904	0.72240	0.50
0.60	0.72575	0.72907	0.73237	0.73565	0.73891	0.74215	0.74537	0.74857	0.75175	0.75490	0.60
0.70	0.75804	0.76115	0.76424	0.76730	0.77035	0.77337	0.77637	0.77935	0.78230	0.78524	0.70
0.80	0.78814	0.79103	0.79389	0.79673	0.79955	0.80234	0.80511	0.80785	0.81057	0.81327	0.80
0.90	0.81594	0.81859	0.82121	0.82381	0.82639	0.82894	0.83147	0.83398	0.83646	0.83891	0.90
1.00	0.84134	0.84375	0.84614	0.84849	0.85083	0.85314	0.85543	0.85769	0.85993	0.86214	1.00
1.10	0.86433	0.86650	0.86864	0.87076	0.87286	0.87493	0.87698	0.87900	0.88100	0.88298	1.10
1.20	0.88493	0.88686	0.88877	0.89065	0.89251	0.89435	0.89617	0.89796	0.89973	0.90147	1.20
1.30	0.90320	0.90490	0.90658	0.90824	0.90988	0.91149	0.91309	0.91466	0.91621	0.91774	1.30
1.40	0.91924	0.92073	0.92220	0.92364	0.92507	0.92647	0.92785	0.92922	0.93056	0.93189	1.40
1.50	0.93319	0.93448	0.93574	0.93699	0.93822	0.93943	0.94062	0.94179	0.94295	0.94408	1.50
1.60	0.94520	0.94630	0.94738	0.94845	0.94950	0.95053	0.95154	0.95254	0.95352	0.95449	1.60
1.70	0.95543	0.95637	0.95728	0.95818	0.95907	0.95994	0.96080	0.96164	0.96246	0.96327	1.70
1.80	0.96407	0.96485	0.96562	0.96638	0.96712	0.96784	0.96856	0.96926	0.96995	0.97062	1.80

$Z = 1.60 + 0.05 = 1.65$

$Z_{0.95053} = 1.65$

Example: Suppose that $Z \sim N(0,1)$.

Find the value of k such that

$$P(Z \leq k) = 0.02068 \gggg Z_{0.02068}$$

Solution:

0.02068 = probability < 0.5 (Use Negative table)

z	-0.09	-0.08	-0.07	-0.06	-0.05	-0.04	-0.03	-0.02	-0.01	-0.00	z
-3.50	0.00017	0.00017	0.00018	0.00019	0.00019	0.00020	0.00021	0.00022	0.00022	0.00023	-3.50
-3.40	0.00024	0.00025	0.00026	0.00027	0.00028	0.00029	0.00030	0.00031	0.00032	0.00034	-3.40
-3.30	0.00035	0.00036	0.00038	0.00039	0.00040	0.00042	0.00043	0.00045	0.00047	0.00048	-3.30
-3.20	0.00050	0.00052	0.00054	0.00056	0.00058	0.00060	0.00062	0.00064	0.00066	0.00069	-3.20
-3.10	0.00071	0.00074	0.00076	0.00079	0.00082	0.00084	0.00087	0.00090	0.00094	0.00097	-3.10
-3.00	0.00100	0.00104	0.00107	0.00111	0.00114	0.00118	0.00122	0.00126	0.00131	0.00135	-3.00
-2.90	0.00139	0.00144	0.00149	0.00154	0.00159	0.00164	0.00169	0.00175	0.00181	0.00187	-2.90
-2.80	0.00193	0.00199	0.00205	0.00212	0.00219	0.00226	0.00233	0.00240	0.00248	0.00256	-2.80
-2.70	0.00264	0.00272	0.00280	0.00289	0.00298	0.00307	0.00317	0.00326	0.00336	0.00347	-2.70
-2.60	0.00357	0.00368	0.00379	0.00391	0.00402	0.00415	0.00427	0.00440	0.00453	0.00466	-2.60
-2.50	0.00480	0.00494	0.00508	0.00523	0.00539	0.00554	0.00570	0.00587	0.00604	0.00621	-2.50
-2.40	0.00639	0.00657	0.00676	0.00695	0.00714	0.00734	0.00755	0.00776	0.00798	0.00820	-2.40
-2.30	0.00842	0.00866	0.00889	0.00914	0.00939	0.00964	0.00990	0.01017	0.01044	0.01072	-2.30
-2.20	0.01101	0.01130	0.01160	0.01191	0.01222	0.01255	0.01287	0.01321	0.01355	0.01390	-2.20
-2.10	0.01426	0.01463	0.01500	0.01539	0.01578	0.01618	0.01659	0.01700	0.01743	0.01786	-2.10
-2.00	0.01831	0.01876	0.01923	0.01970	0.02018	0.02068	0.02118	0.02169	0.02222	0.02275	-2.00
-1.90	0.02330	0.02385	0.02442	0.02500	0.02559	0.02619	0.02680	0.02743	0.02807	0.02872	-1.90
-1.80	0.02938	0.03005	0.03074	0.03144	0.03216	0.03288	0.03362	0.03438	0.03515	0.03593	-1.80

$$Z = (-2.00) + (-0.04) = -2.04$$

$$K = -2.04$$

$$Z_{0.02068} = -2.04$$

Example:

If $Z \sim N(0,1)$, then:

$$Z_{0.90} = ?? \gg P(Z \leq z) = 0.90$$

0.90 = probability > 0.5 (Use positive table)

z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09	z
0.00	0.50000	0.50399	0.50798	0.51197	0.51595	0.51994	0.52392	0.52790	0.53188	0.53586	0.00
0.10	0.53983	0.54380	0.54776	0.55172	0.55567	0.55962	0.56356	0.56749	0.57142	0.57535	0.10
0.20	0.57926	0.58317	0.58706	0.59095	0.59483	0.59871	0.60257	0.60642	0.61026	0.61409	0.20
0.30	0.61791	0.62172	0.62552	0.62930	0.63307	0.63683	0.64058	0.64431	0.64803	0.65173	0.30
0.40	0.65542	0.65910	0.66276	0.66640	0.67003	0.67364	0.67724	0.68082	0.68439	0.68793	0.40
0.50	0.69146	0.69497	0.69847	0.70194	0.70540	0.70884	0.71226	0.71566	0.71904	0.72240	0.50
0.60	0.72575	0.72907	0.73237	0.73565	0.73891	0.74215	0.74537	0.74857	0.75175	0.75490	0.60
0.70	0.75804	0.76115	0.76424	0.76730	0.77035	0.77337	0.77637	0.77935	0.78230	0.78524	0.70
0.80	0.78814	0.79103	0.79389	0.79673	0.79955	0.80234	0.80511	0.80785	0.81057	0.81327	0.80
0.90	0.81594	0.81859	0.82121	0.82381	0.82639	0.82894	0.83147	0.83398	0.83646	0.83891	0.90
1.00	0.84134	0.84375	0.84614	0.84849	0.85083	0.85314	0.85543	0.85769	0.85993	0.86214	1.00
1.10	0.86433	0.86650	0.86864	0.87076	0.87286	0.87493	0.87698	0.87900	0.88100	0.88298	1.10
1.20	0.88493	0.88686	0.88877	0.89065	0.89251	0.89435	0.89617	0.89796	0.89973	0.90147	1.20
1.30	0.90320	0.90490	0.90658	0.90824	0.90988	0.91149	0.91309	0.91466	0.91621	0.91774	1.30
1.40	0.91924	0.92073	0.92220	0.92364	0.92507	0.92647	0.92785	0.92922	0.93056	0.93189	1.40
1.50	0.93319	0.93448	0.93574	0.93699	0.93822	0.93943	0.94062	0.94179	0.94295	0.94408	1.50
1.60	0.94520	0.94630	0.94738	0.94845	0.94950	0.95053	0.95154	0.95254	0.95352	0.95449	1.60
1.70	0.95543	0.95637	0.95728	0.95818	0.95907	0.95994	0.96080	0.96164	0.96246	0.96327	1.70
1.80	0.96407	0.96485	0.96562	0.96638	0.96712	0.96784	0.96856	0.96926	0.96995	0.97062	1.80
1.90	0.97128	0.97193	0.97257	0.97320	0.97381	0.97441	0.97500	0.97558	0.97615	0.97670	1.90
2.00	0.97725	0.97778	0.97831	0.97882	0.97932	0.97982	0.98030	0.98077	0.98124	0.98169	2.00
2.10	0.98214	0.98257	0.98300	0.98341	0.98382	0.98422	0.98461	0.98500	0.98537	0.98574	2.10
2.20	0.98610	0.98645	0.98679	0.98713	0.98745	0.98778	0.98809	0.98840	0.98870	0.98899	2.20

$$Z_{0.90} = 1.285$$

Value of
Probability =
→ 0.90000

first number = 1.28
second number = 1.29
The value of $Z = (1.28 + 1.29) / 2$
= 1.285

Example:

If $Z \sim N(0,1)$, then:

$Z_{0.95} = ?? \gg P(Z \leq z) = 0.95$

0.95 =probability > 0.5 (Use positive table)

z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09	z
0.00	0.50000	0.50399	0.50798	0.51197	0.51595	0.51994	0.52392	0.52790	0.53188	0.53586	0.00
0.10	0.53983	0.54380	0.54776	0.55172	0.55567	0.55962	0.56356	0.56749	0.57142	0.57535	0.10
0.20	0.57926	0.58317	0.58706	0.59095	0.59483	0.59871	0.60257	0.60642	0.61026	0.61409	0.20
0.30	0.61791	0.62172	0.62552	0.62930	0.63307	0.63683	0.64058	0.64431	0.64803	0.65173	0.30
0.40	0.65542	0.65910	0.66276	0.66640	0.67003	0.67364	0.67724	0.68082	0.68439	0.68793	0.40
0.50	0.69146	0.69497	0.69847	0.70194	0.70540	0.70884	0.71226	0.71566	0.71904	0.72240	0.50
0.60	0.72575	0.72907	0.73237	0.73565	0.73891	0.74215	0.74537	0.74857	0.75175	0.75490	0.60
0.70	0.75804	0.76115	0.76424	0.76730	0.77035	0.77337	0.77637	0.77935	0.78230	0.78524	0.70
0.80	0.78814	0.79103	0.79389	0.79673	0.79955	0.80234	0.80511	0.80785	0.81057	0.81327	0.80
0.90	0.81594	0.81859	0.82121	0.82381	0.82639	0.82894	0.83147	0.83398	0.83646	0.83891	0.90
1.00	0.84134	0.84375	0.84614	0.84849	0.85083	0.85314	0.85543	0.85769	0.85993	0.86214	1.00
1.10	0.86433	0.86650	0.86864	0.87076	0.87286	0.87493	0.87698	0.87900	0.88100	0.88298	1.10
1.20	0.88493	0.88686	0.88877	0.89065	0.89251	0.89435	0.89617	0.89796	0.89973	0.90147	1.20
1.30	0.90320	0.90490	0.90658	0.90824	0.90988	0.91149	0.91309	0.91466	0.91621	0.91774	1.30
1.40	0.91924	0.92073	0.92220	0.92364	0.92507	0.92647	0.92785	0.92922	0.93056	0.93189	1.40
1.50	0.93319	0.93448	0.93574	0.93699	0.93822	0.93943	0.94062	0.94179	0.94295	0.94408	1.50
1.60	0.94520	0.94630	0.94738	0.94845	0.94950	0.95053	0.95154	0.95254	0.95352	0.95449	1.60
1.70	0.95543	0.95637	0.95728	0.95818	0.95907	0.95994	0.96080	0.96164	0.96246	0.96327	1.70
1.80	0.96407	0.96485	0.96562	0.96638	0.96712	0.96784	0.96856	0.96926	0.96995	0.97062	1.80
1.90	0.97128	0.97193	0.97257	0.97320	0.97381	0.97441	0.97500	0.97558	0.97615	0.97670	1.90
2.00	0.97725	0.97778	0.97831	0.97882	0.97932	0.97982	0.98030	0.98077	0.98124	0.98169	2.00

first number =1.64
second number =1.65

The value of Z = (1.64+1.65)/2
= 1.645

$Z_{0.95} = 1.645$

→ Probability = 0.95000

Example:

If $Z \sim N(0,1)$:

If $P(Z > z) = 0.63307$, find the value of z ?

$P(Z > z) = 0.63307$

$\gg P(Z < z) = 1 - 0.63307$

$P(Z < z) = 0.36693$

$Z = -0.34$

z	-0.09	-0.08	-0.07	-0.06	-0.05	-0.04	-0.03	-0.02	-0.01	-0.00	z
-3.50	0.00017	0.00017	0.00018	0.00019	0.00019	0.00020	0.00021	0.00022	0.00022	0.00023	-3.50
-3.40	0.00024	0.00025	0.00026	0.00027	0.00028	0.00029	0.00030	0.00031	0.00032	0.00034	-3.40
-3.30	0.00035	0.00036	0.00038	0.00039	0.00040	0.00042	0.00043	0.00045	0.00047	0.00048	-3.30
-3.20	0.00050	0.00052	0.00054	0.00056	0.00058	0.00060	0.00062	0.00064	0.00066	0.00069	-3.20
-3.10	0.00071	0.00074	0.00076	0.00079	0.00082	0.00084	0.00087	0.00090	0.00094	0.00097	-3.10
-3.00	0.00100	0.00104	0.00107	0.00111	0.00114	0.00118	0.00122	0.00126	0.00131	0.00135	-3.00
-2.90	0.00139	0.00144	0.00149	0.00154	0.00159	0.00164	0.00169	0.00175	0.00181	0.00187	-2.90
-2.80	0.00193	0.00199	0.00205	0.00212	0.00219	0.00226	0.00233	0.00240	0.00248	0.00256	-2.80
-2.70	0.00264	0.00272	0.00280	0.00289	0.00298	0.00307	0.00317	0.00326	0.00336	0.00347	-2.70
-2.60	0.00357	0.00368	0.00379	0.00391	0.00402	0.00415	0.00427	0.00440	0.00453	0.00466	-2.60
-2.50	0.00480	0.00494	0.00508	0.00523	0.00539	0.00554	0.00570	0.00587	0.00604	0.00621	-2.50
-2.40	0.00639	0.00657	0.00676	0.00695	0.00714	0.00734	0.00755	0.00776	0.00798	0.00820	-2.40
-2.30	0.00842	0.00866	0.00889	0.00914	0.00939	0.00964	0.00990	0.01017	0.01044	0.01072	-2.30
-2.20	0.01101	0.01130	0.01160	0.01191	0.01222	0.01255	0.01287	0.01321	0.01355	0.01390	-2.20
-2.10	0.01426	0.01463	0.01500	0.01539	0.01578	0.01618	0.01659	0.01700	0.01743	0.01786	-2.10
-2.00	0.01831	0.01876	0.01923	0.01970	0.02018	0.02068	0.02118	0.02169	0.02222	0.02275	-2.00
-1.90	0.02330	0.02385	0.02442	0.02500	0.02559	0.02619	0.02680	0.02743	0.02807	0.02872	-1.90
-1.80	0.02938	0.03005	0.03074	0.03144	0.03216	0.03288	0.03362	0.03438	0.03515	0.03593	-1.80
-1.70	0.03673	0.03754	0.03836	0.03920	0.04006	0.04093	0.04182	0.04272	0.04363	0.04457	-1.70
-1.60	0.04551	0.04648	0.04746	0.04846	0.04947	0.05050	0.05155	0.05262	0.05370	0.05480	-1.60
-1.50	0.05592	0.05705	0.05821	0.05938	0.06057	0.06178	0.06301	0.06426	0.06552	0.06681	-1.50
-1.40	0.06811	0.06944	0.07078	0.07215	0.07353	0.07493	0.07636	0.07780	0.07927	0.08076	-1.40
-1.30	0.08226	0.08379	0.08534	0.08691	0.08851	0.09012	0.09176	0.09342	0.09510	0.09680	-1.30
-1.20	0.09853	0.10027	0.10204	0.10383	0.10565	0.10749	0.10935	0.11123	0.11314	0.11507	-1.20
-1.10	0.11702	0.11900	0.12100	0.12302	0.12507	0.12714	0.12924	0.13136	0.13350	0.13567	-1.10
-1.00	0.13786	0.14007	0.14231	0.14457	0.14686	0.14917	0.15151	0.15386	0.15625	0.15866	-1.00
-0.90	0.16109	0.16354	0.16602	0.16853	0.17106	0.17361	0.17619	0.17879	0.18141	0.18406	-0.90
-0.80	0.18673	0.18943	0.19215	0.19489	0.19766	0.20045	0.20327	0.20611	0.20897	0.21186	-0.80
-0.70	0.21476	0.21770	0.22065	0.22363	0.22663	0.22965	0.23270	0.23576	0.23885	0.24196	-0.70
-0.60	0.24510	0.24825	0.25143	0.25463	0.25785	0.26109	0.26435	0.26763	0.27093	0.27425	-0.60
-0.50	0.27760	0.28096	0.28434	0.28774	0.29116	0.29460	0.29806	0.30153	0.30503	0.30854	-0.50
-0.40	0.31207	0.31561	0.31918	0.32276	0.32636	0.32997	0.33360	0.33724	0.3409	0.34458	-0.40
-0.30	0.34827	0.35197	0.35569	0.35942	0.36317	0.36693	0.37070	0.37448	0.37828	0.38209	-0.30

Example:

If $Z \sim N(0,1)$, then:

$$Z_{0.90} = 1.285$$

$$Z_{0.95} = 1.645$$





$$Z_{0.975} = 1.96$$

$$Z_{0.99} = 2.325$$

$$Z_{0.90} = (Z_{0.89973} + Z_{0.90147})/2 = (1.28 + 1.29)/2 = 1.285$$

$$Z_{0.95} = (Z_{0.94950} + Z_{0.95053})/2 = (1.64 + 1.65)/2 = 1.645$$

$$Z_{0.99} = (Z_{0.98983} + Z_{0.99010})/2 = (2.32 + 2.33)/2 = 2.325$$



Using the result:

$$Z_A = -Z_{1-A}$$

$$Z_{0.10} = -Z_{0.90} = -1.285$$

$$Z_{0.05} = -Z_{0.95} = -1.645$$

$$Z_{0.025} = -Z_{0.975} = -1.96$$

$$Z_{0.01} = -Z_{0.99} = -2.325$$

From previous examples:

$$Z_{0.90} = 1.285$$

$$Z_{0.95} = 1.645$$

$$Z_{0.975} = 1.96$$

$$Z_{0.99} = 2.325$$

Calculating Probabilities of Normal (μ, σ^2) :

Recall the result : **If** $X \sim N(\mu, \sigma^2)$, **then** $Z = \frac{x - \mu}{\sigma} \sim N(0, 1)$

$$\blacksquare \quad X \leq a \Leftrightarrow \frac{X - \mu}{\sigma} \leq \frac{a - \mu}{\sigma} \Leftrightarrow Z \leq \frac{a - \mu}{\sigma}$$

$$1. \quad P(X \leq a) = P\left(Z \leq \frac{a - \mu}{\sigma}\right) = \text{From the table.}$$

$$2. \quad P(X \geq a) = 1 - P(X \leq a) = 1 - P\left(Z \leq \frac{a - \mu}{\sigma}\right)$$

$$\begin{aligned} 3. \quad P(a \leq X \leq b) &= P(X \leq b) - P(X \leq a) \\ &= P\left(Z \leq \frac{b - \mu}{\sigma}\right) - P\left(Z \leq \frac{a - \mu}{\sigma}\right) \end{aligned}$$

$$4. \quad P(X = a) = 0, \text{ for every } a.$$

4.7 Normal Distribution Application:

Example :

Suppose that the hemoglobin levels of healthy adult males are approximately normally distributed with a mean of 16 and a variance of 0.81.

- 1- Find that probability that a randomly chosen healthy adult male has a hemoglobin level less than 14.
- 2- What is the percentage of healthy adult males who have hemoglobin level less than 14 ?
- 3- In a population of 10,000 healthy adult males, how many would you expect to have hemoglobin level less than 14 ?

Solution :

X = hemoglobin level for healthy adult males

Mean: $\mu = 16$


Variance: $\sigma^2 = 0.81$

Standard deviation : $\sigma = 0.9$

$X \sim \text{Normal} (\text{mean} = 16, \text{varaince} = 0.81)$

1- The probability that a randomly chosen healthy adult male has hemoglobin level less than 14 is :

$$\begin{aligned}
 P(X < 14) &= P(Z < \frac{14-\mu}{\sigma}) \\
 &= P(Z < \frac{14-16}{0.9}) \\
 &= P(Z < -2.22) \\
 &= 0.01321
 \end{aligned}$$



z	-0.09	-0.08	-0.07	-0.06	-0.05	-0.04	-0.03	-0.02	-0.01	-0.00	z
-3.50	0.00017	0.00017	0.00018	0.00019	0.00019	0.00020	0.00021	0.00022	0.00022	0.00023	-3.50
-3.40	0.00024	0.00025	0.00026	0.00027	0.00028	0.00029	0.00030	0.00031	0.00032	0.00034	-3.40
-3.30	0.00035	0.00036	0.00038	0.00039	0.00040	0.00042	0.00043	0.00045	0.00047	0.00048	-3.30
-3.20	0.00050	0.00052	0.00054	0.00056	0.00058	0.00060	0.00062	0.00064	0.00066	0.00069	-3.20
-3.10	0.00071	0.00074	0.00076	0.00079	0.00082	0.00084	0.00087	0.00090	0.00094	0.00097	-3.10
-3.00	0.00100	0.00104	0.00107	0.00111	0.00114	0.00118	0.00122	0.00126	0.00131	0.00135	-3.00
-2.90	0.00139	0.00144	0.00149	0.00154	0.00159	0.00164	0.00169	0.00175	0.00181	0.00187	-2.90
-2.80	0.00193	0.00199	0.00205	0.00212	0.00219	0.00226	0.00233	0.00240	0.00248	0.00256	-2.80
-2.70	0.00264	0.00272	0.00280	0.00289	0.00298	0.00307	0.00317	0.00326	0.00336	0.00347	-2.70
-2.60	0.00357	0.00368	0.00379	0.00391	0.00402	0.00415	0.00427	0.00440	0.00453	0.00466	-2.60
-2.50	0.00480	0.00494	0.00508	0.00523	0.00539	0.00554	0.00570	0.00587	0.00604	0.00621	-2.50
-2.40	0.00639	0.00657	0.00676	0.00695	0.00714	0.00734	0.00755	0.00776	0.00798	0.00820	-2.40
-2.30	0.00842	0.00866	0.00889	0.00914	0.00939	0.00964	0.00990	0.01017	0.01044	0.01072	-2.30
-2.20	0.01101	0.01128	0.01156	0.01184	0.01212	0.01241	0.01270	0.01321	0.01355	0.01390	-2.20
-2.10	0.01426	0.01463	0.01500	0.01539	0.01578	0.01618	0.01659	0.01700	0.01743	0.01786	-2.10
-2.00	0.01831	0.01876	0.01923	0.01970	0.02018	0.02068	0.02118	0.02169	0.02222	0.02275	-2.00
-1.90	0.02330	0.02385	0.02442	0.02500	0.02559	0.02619	0.02680	0.02743	0.02807	0.02872	-1.90
-1.80	0.02938	0.03005	0.03074	0.03144	0.03216	0.03288	0.03362	0.03438	0.03515	0.03593	-1.80

2- The percentage of healthy adult males who have hemoglobin level less than 14 is :

Percentage = Probability x 100%

$$P(X < 14) \times 100 \% = 0.01321 \times 100 \% = 1.321\%$$

3- In a population of 10000 healthy adult males, we would expect that the number of males with hemoglobin level less than 14 to be: (**N = population size = 10000**)

Expect number = Probability x N

$$P(X < 14) \times 10000 = 0.01321 \times 10000 = 132.1 \approx 133 \text{ males.}$$

Example:

Suppose that the birth weight of Saudi babies has a normal distribution with mean $\mu = 3.4$ and standard deviation $\sigma = 0.35$.

- 1- Find the probability that a randomly chosen Saudi baby has a birth weight between 3.0 and 4.0 kg.
- 2- What is the percentage of Saudi babies who have a birth weight between 3.0 and 4.0 kg?
- 3- In a population of 100000 Saudi babies, how many would you expect to have birth weight between 3.0 and 4.0 kg?

Solution :

X = weight of Saudi babies

Mean: $\mu = 3.4$

Variance: $\sigma^2 = (0.35)^2 = 0.1225$


Standard deviation : $\sigma = 0.35$

$X \sim \text{Normal}(\text{mean} = 3.4, \text{variance} = 0.1225)$

Solution:

1-The **probability** that a randomly chosen Saudi baby has a birth weight between 3.0 and 4.0 kg is :

$$\begin{aligned}
 P(3 < X < 4) &= P\left(\frac{3-\mu}{\sigma} < Z < \frac{4-\mu}{\sigma}\right) \\
 &= P\left(\frac{3-3.4}{0.35} < Z < \frac{4-3.4}{0.35}\right) \\
 &= P(-1.14 < Z < 1.71) \\
 &= P(Z < 1.71) - P(Z < -1.14) \\
 &= 0.9564 - 0.1271 \\
 &= 0.8293
 \end{aligned}$$



$$P(Z < 1.71) = 0.95637$$

z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09	z
0.00	0.50000	0.50399	0.50798	0.51197	0.51595	0.51994	0.52392	0.52790	0.53188	0.53586	0.00
0.10	0.53983	0.54380	0.54776	0.55172	0.55567	0.55962	0.56356	0.56749	0.57142	0.57535	0.10
0.20	0.57926	0.58317	0.58706	0.59095	0.59483	0.59871	0.60257	0.60642	0.61026	0.61409	0.20
0.30	0.61791	0.62172	0.62552	0.62930	0.63307	0.63683	0.64058	0.64431	0.64803	0.65173	0.30
0.40	0.65542	0.65910	0.66276	0.66640	0.67003	0.67364	0.67724	0.68082	0.68439	0.68793	0.40
0.50	0.69146	0.69497	0.69847	0.70194	0.70540	0.70884	0.71226	0.71566	0.71904	0.72240	0.50
0.60	0.72575	0.72910	0.73237	0.73565	0.73891	0.74215	0.74537	0.74857	0.75175	0.75490	0.60
0.70	0.75804	0.76115	0.76424	0.76730	0.77035	0.77337	0.77637	0.77935	0.78230	0.78524	0.70
0.80	0.78814	0.79103	0.79389	0.79673	0.79955	0.80234	0.80511	0.80785	0.81057	0.81327	0.80
0.90	0.81594	0.81859	0.82121	0.82381	0.82639	0.82894	0.83147	0.83398	0.83646	0.83891	0.90
1.00	0.84134	0.84375	0.84614	0.84849	0.85083	0.85314	0.85543	0.85769	0.85993	0.86214	1.00
1.10	0.86433	0.86650	0.86864	0.87076	0.87286	0.87493	0.87698	0.87900	0.88100	0.88298	1.10
1.20	0.88493	0.88686	0.88877	0.89065	0.89251	0.89435	0.89617	0.89796	0.89973	0.90147	1.20
1.30	0.90320	0.90490	0.90658	0.90824	0.90988	0.91149	0.91309	0.91466	0.91621	0.91774	1.30
1.40	0.91924	0.92073	0.92220	0.92364	0.92507	0.92647	0.92785	0.92922	0.93056	0.93189	1.40
1.50	0.93319	0.93458	0.93574	0.93699	0.93822	0.93943	0.94062	0.94179	0.94295	0.94408	1.50
1.60	0.94520	0.94620	0.94718	0.94815	0.94950	0.95053	0.95154	0.95254	0.95352	0.95449	1.60
1.70	0.95544	0.95637	0.95728	0.95818	0.95907	0.95994	0.96080	0.96164	0.96246	0.96327	1.70
1.80	0.96407	0.96485	0.96562	0.96638	0.96712	0.96784	0.96856	0.96926	0.96995	0.97062	1.80

$$P(Z < -1.14) = 0.12714$$

z	-0.09	-0.08	-0.07	-0.06	-0.05	-0.04	-0.03	-0.02	-0.01	-0.00	z
-3.50	0.00017	0.00017	0.00018	0.00019	0.00019	0.00020	0.00021	0.00022	0.00022	0.00023	-3.50
-3.40	0.00024	0.00025	0.00026	0.00027	0.00028	0.00029	0.00030	0.00031	0.00032	0.00034	-3.40
-3.30	0.00035	0.00036	0.00038	0.00039	0.00040	0.00042	0.00043	0.00045	0.00047	0.00048	-3.30
-3.20	0.00050	0.00052	0.00054	0.00056	0.00058	0.00060	0.00062	0.00064	0.00066	0.00069	-3.20
-3.10	0.00071	0.00074	0.00076	0.00079	0.00082	0.00084	0.00087	0.00090	0.00094	0.00097	-3.10
-3.00	0.00100	0.00104	0.00107	0.00111	0.00114	0.00118	0.00122	0.00126	0.00131	0.00135	-3.00
-2.90	0.00139	0.00144	0.00149	0.00154	0.00159	0.00164	0.00169	0.00175	0.00181	0.00187	-2.90
-2.80	0.00193	0.00199	0.00205	0.00212	0.00219	0.00226	0.00233	0.00240	0.00248	0.00256	-2.80
-2.70	0.00264	0.00272	0.00280	0.00289	0.00298	0.00307	0.00317	0.00326	0.00336	0.00347	-2.70
-2.60	0.00357	0.00368	0.00379	0.00391	0.00402	0.00415	0.00427	0.00440	0.00453	0.00466	-2.60
-2.50	0.00480	0.00494	0.00508	0.00523	0.00539	0.00554	0.00570	0.00587	0.00604	0.00621	-2.50
-2.40	0.00639	0.00657	0.00676	0.00695	0.00714	0.00734	0.00755	0.00776	0.00798	0.00820	-2.40
-2.30	0.00842	0.00866	0.00889	0.00914	0.00939	0.00964	0.00990	0.01017	0.01044	0.01072	-2.30
-2.20	0.01101	0.01130	0.01160	0.01191	0.01222	0.01255	0.01287	0.01321	0.01355	0.01390	-2.20
-2.10	0.01426	0.01463	0.01500	0.01539	0.01578	0.01618	0.01659	0.01700	0.01743	0.01786	-2.10
-2.00	0.01831	0.01876	0.01923	0.01970	0.02018	0.02068	0.02118	0.02169	0.02222	0.02275	-2.00
-1.90	0.02330	0.02385	0.02442	0.02500	0.02559	0.02619	0.02680	0.02743	0.02807	0.02872	-1.90
-1.80	0.02938	0.03005	0.03074	0.03144	0.03216	0.03288	0.03362	0.03438	0.03515	0.03593	-1.80
-1.70	0.03673	0.03754	0.03836	0.03920	0.04006	0.04093	0.04182	0.04272	0.04363	0.04457	-1.70
-1.60	0.04551	0.04648	0.04746	0.04846	0.04947	0.05050	0.05155	0.05262	0.05370	0.05480	-1.60
-1.50	0.05592	0.05705	0.05821	0.05938	0.06057	0.06178	0.06301	0.06426	0.06552	0.06681	-1.50
-1.40	0.06811	0.06944	0.07078	0.07215	0.07353	0.07493	0.07636	0.07780	0.07927	0.08076	-1.40
-1.30	0.08226	0.08379	0.08534	0.08691	0.08851	0.09012	0.09176	0.09342	0.09510	0.09680	-1.30
-1.20	0.09853	0.10027	0.10204	0.10383	0.10565	0.10749	0.10935	0.11123	0.11314	0.11507	-1.20
-1.10	0.11693	0.11881	0.12071	0.12263	0.12457	0.12714	0.12924	0.13136	0.13350	0.13567	-1.10
-1.00	0.13786	0.14007	0.14231	0.14457	0.14686	0.14917	0.15151	0.15386	0.15625	0.15866	-1.00

2-The **percentage** of Saudi babies who have a birth weight between 3.0 and 4.0 kg is

Percentage = Probability x 100%

$$P(3.0 < X < 4.0) \times 100\% = 0.8293 \times 100\% = 82.93\%$$

3- In a population of 100,000 Saudi babies, we would expect that the number of babies with birth weight between 3.0 and 4.0 kg to be:

Expect number = Probability x N

$$P(3.0 < X < 4.0) \times 100000 = 0.8293 \times 100000 = 82930 \text{ babies}$$

Example:

Given the normally distributed random variable X with mean 491 and standard deviation 119,

- If $P(X < k) = 0.9082$, the value of k is equal to

$$P(X < k) = 0.9082 \Rightarrow P\left(Z < \frac{k-491}{119}\right) = 0.9082 > 0.5$$

$$\frac{k-491}{119} = \frac{1.32+1.33}{2}$$

$$\frac{k-491}{119} = 1.325$$

$$k = (1.325 \times 119) + 491$$

$$\Rightarrow k = 648.675$$

Probability =
0.90820

z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09	z
0.00	0.50000	0.50399	0.50798	0.51197	0.51595	0.51994	0.52392	0.52790	0.53188	0.53586	0.00
0.10	0.53983	0.54380	0.54776	0.55172	0.55567	0.55962	0.56356	0.56749	0.57142	0.57535	0.10
0.20	0.57926	0.58317	0.58706	0.59095	0.59483	0.59871	0.60257	0.60642	0.61026	0.61409	0.20
0.30	0.61791	0.62172	0.62552	0.62930	0.63307	0.63683	0.64058	0.64431	0.64803	0.65173	0.30
0.40	0.65542	0.65910	0.66276	0.66640	0.67003	0.67364	0.67724	0.68082	0.68439	0.68793	0.40
0.50	0.69146	0.69497	0.69847	0.70194	0.70540	0.70884	0.71226	0.71566	0.71904	0.72240	0.50
0.60	0.72575	0.72907	0.73237	0.73565	0.73891	0.74215	0.74537	0.74857	0.75175	0.75490	0.60
0.70	0.75804	0.76115	0.76424	0.76730	0.77035	0.77337	0.77637	0.77935	0.78230	0.78524	0.70
0.80	0.78814	0.79103	0.79389	0.79673	0.79955	0.80234	0.80511	0.80785	0.81057	0.81327	0.80
0.90	0.81594	0.81859	0.82121	0.82381	0.82639	0.82894	0.83147	0.83398	0.83646	0.83891	0.90
1.00	0.84134	0.84375	0.84614	0.84849	0.85083	0.85314	0.85543	0.85769	0.85993	0.86214	1.00
1.10	0.86433	0.86650	0.86864	0.87076	0.87286	0.87493	0.87698	0.87900	0.88100	0.88298	1.10
1.20	0.88493	0.88686	0.88877	0.89065	0.89251	0.89435	0.89617	0.89796	0.89973	0.90147	1.20
1.30	0.90320	0.90490	0.90658	0.90824	0.90988	0.91149	0.91309	0.91466	0.91621	0.91774	1.30
1.40	0.91924	0.92073	0.92220	0.92364	0.92507	0.92647	0.92785	0.92922	0.93056	0.93189	1.40
1.50	0.93319	0.93448	0.93574	0.93699	0.93822	0.93943	0.94062	0.94179	0.94295	0.94408	1.50
1.60	0.94520	0.94630	0.94738	0.94845	0.94950	0.95053	0.95154	0.95254	0.95352	0.95449	1.60
1.70	0.95543	0.95637	0.95728	0.95818	0.95907	0.95994	0.96080	0.96164	0.96246	0.96327	1.70
1.80	0.96407	0.96485	0.96562	0.96638	0.96712	0.96784	0.96856	0.96926	0.96995	0.97062	1.80
1.90	0.97128	0.97193	0.97257	0.97320	0.97381	0.97441	0.97500	0.97558	0.97615	0.97670	1.90

If the random variable X has a normal distribution with the mean μ and the variance 1, and if then $P(X < 3) = 0.877$ then μ equal to

Given that $\sigma = 1$

$$P(X < 3) = 0.877 \Rightarrow P\left(Z < \frac{3-\mu}{1}\right) = 0.877$$

$$3 - \mu = \frac{1.16+1.17}{2}$$

$$3 - \mu = 1.165$$


$$\mu = 1.835$$

z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09	z
0.00	0.50000	0.50399	0.50798	0.51197	0.51595	0.51994	0.52392	0.52790	0.53188	0.53586	0.00
0.10	0.53983	0.54380	0.54776	0.55172	0.55567	0.55962	0.56356	0.56749	0.57142	0.57535	0.10
0.20	0.57926	0.58317	0.58706	0.59095	0.59483	0.59871	0.60257	0.60642	0.61026	0.61409	0.20
0.30	0.61791	0.62172	0.62552	0.62930	0.63307	0.63683	0.64058	0.64431	0.64803	0.65173	0.30
0.40	0.65542	0.65910	0.66276	0.66640	0.67003	0.67364	0.67724	0.68082	0.68439	0.68793	0.40
0.50	0.69146	0.69497	0.69847	0.70194	0.70540	0.70884	0.71226	0.71566	0.71904	0.72240	0.50
0.60	0.72575	0.72907	0.73237	0.73565	0.73891	0.74215	0.74537	0.74857	0.75175	0.75490	0.60
0.70	0.75804	0.76115	0.76424	0.76730	0.77035	0.77337	0.77637	0.77935	0.78230	0.78524	0.70
0.80	0.78814	0.79103	0.79389	0.79673	0.79955	0.80234	0.80511	0.80785	0.81057	0.81327	0.80
0.90	0.81594	0.81859	0.82121	0.82381	0.82639	0.82894	0.83147	0.83398	0.83646	0.83891	0.90
1.00	0.84134	0.84375	0.84614	0.84849	0.85083	0.85314	0.85543	0.85769	0.85993	0.86214	1.00
1.10	0.86433	0.86650	0.86864	0.87076	0.87286	0.87493	0.87698	0.87900	0.88100	0.88298	1.10
1.20	0.88493	0.88686	0.88877	0.89065	0.89251	0.89435	0.89617	0.89796	0.89973	0.90147	1.20
1.30	0.90320	0.90490	0.90658	0.90824	0.90988	0.91149	0.91309	0.91466	0.91621	0.91774	1.30
1.40	0.91924	0.92073	0.92220	0.92364	0.92507	0.92647	0.92785	0.92922	0.93056	0.93189	1.40
1.50	0.93319	0.93448	0.93574	0.93699	0.93822	0.93943	0.94062	0.94179	0.94295	0.94408	1.50

$$Z = \frac{1.16+1.17}{2} = 1.165$$


Probability = 0.87700

Standard Normal Table
Areas Under the Standard Normal Curve



z	-0.09	-0.08	-0.07	-0.06	-0.05	-0.04	-0.03	-0.02	-0.01	-0.00	z
-3.50	0.00017	0.00017	0.00018	0.00019	0.00019	0.00020	0.00021	0.00022	0.00022	0.00023	-3.50
-3.40	0.00024	0.00025	0.00026	0.00027	0.00028	0.00029	0.00030	0.00031	0.00032	0.00034	-3.40
-3.30	0.00035	0.00036	0.00038	0.00039	0.00040	0.00042	0.00043	0.00045	0.00047	0.00048	-3.30
-3.20	0.00050	0.00052	0.00054	0.00056	0.00058	0.00060	0.00062	0.00064	0.00066	0.00069	-3.20
-3.10	0.00071	0.00074	0.00076	0.00079	0.00082	0.00084	0.00087	0.00090	0.00094	0.00097	-3.10
-3.00	0.00100	0.00104	0.00107	0.00111	0.00114	0.00118	0.00122	0.00126	0.00131	0.00135	-3.00
-2.90	0.00139	0.00144	0.00149	0.00154	0.00159	0.00164	0.00169	0.00175	0.00181	0.00187	-2.90
-2.80	0.00193	0.00199	0.00205	0.00212	0.00219	0.00226	0.00233	0.00240	0.00248	0.00256	-2.80
-2.70	0.00264	0.00272	0.00280	0.00289	0.00298	0.00307	0.00317	0.00326	0.00336	0.00347	-2.70
-2.60	0.00357	0.00368	0.00379	0.00391	0.00402	0.00415	0.00427	0.00440	0.00453	0.00466	-2.60
-2.50	0.00480	0.00494	0.00508	0.00523	0.00539	0.00554	0.00570	0.00587	0.00604	0.00621	-2.50
-2.40	0.00639	0.00657	0.00676	0.00695	0.00714	0.00734	0.00755	0.00776	0.00798	0.00820	-2.40
-2.30	0.00842	0.00866	0.00889	0.00914	0.00939	0.00964	0.00990	0.01017	0.01044	0.01072	-2.30
-2.20	0.01101	0.01130	0.01160	0.01191	0.01222	0.01255	0.01287	0.01321	0.01355	0.01390	-2.20
-2.10	0.01426	0.01463	0.01500	0.01539	0.01578	0.01618	0.01659	0.01700	0.01743	0.01786	-2.10
-2.00	0.01831	0.01876	0.01923	0.01970	0.02018	0.02068	0.02118	0.02169	0.02222	0.02275	-2.00
-1.90	0.02330	0.02385	0.02442	0.02500	0.02559	0.02619	0.02680	0.02743	0.02807	0.02872	-1.90
-1.80	0.02938	0.03005	0.03074	0.03144	0.03216	0.03288	0.03362	0.03438	0.03515	0.03593	-1.80
-1.70	0.03673	0.03754	0.03836	0.03920	0.04006	0.04093	0.04182	0.04272	0.04363	0.04457	-1.70
-1.60	0.04551	0.04648	0.04746	0.04846	0.04947	0.05050	0.05155	0.05262	0.05370	0.05480	-1.60
-1.50	0.05592	0.05705	0.05821	0.05938	0.06057	0.06178	0.06301	0.06426	0.06552	0.06681	-1.50
-1.40	0.06811	0.06944	0.07078	0.07215	0.07353	0.07493	0.07636	0.07780	0.07927	0.08076	-1.40
-1.30	0.08226	0.08379	0.08534	0.08691	0.08851	0.09012	0.09176	0.09342	0.09510	0.09680	-1.30
-1.20	0.09853	0.10027	0.10204	0.10383	0.10565	0.10749	0.10935	0.11123	0.11314	0.11507	-1.20
-1.10	0.11702	0.11900	0.12100	0.12302	0.12507	0.12714	0.12924	0.13136	0.13350	0.13567	-1.10
-1.00	0.13786	0.14007	0.14231	0.14457	0.14686	0.14917	0.15151	0.15386	0.15625	0.15866	-1.00
-0.90	0.16109	0.16354	0.16602	0.16853	0.17106	0.17361	0.17619	0.17879	0.18141	0.18406	-0.90
-0.80	0.18673	0.18943	0.19215	0.19489	0.19766	0.20045	0.20327	0.20611	0.20897	0.21186	-0.80
-0.70	0.21476	0.21770	0.22065	0.22363	0.22663	0.22965	0.23270	0.23576	0.23885	0.24196	-0.70
-0.60	0.24510	0.24825	0.25143	0.25463	0.25785	0.26109	0.26435	0.26763	0.27093	0.27425	-0.60
-0.50	0.27760	0.28096	0.28434	0.28774	0.29116	0.29460	0.29806	0.30153	0.30503	0.30854	-0.50
-0.40	0.31207	0.31561	0.31918	0.32276	0.32636	0.32997	0.33360	0.33724	0.3409	0.34458	-0.40
-0.30	0.34827	0.35197	0.35569	0.35942	0.36317	0.36693	0.37070	0.37448	0.37828	0.38209	-0.30
-0.20	0.38591	0.38974	0.39358	0.39743	0.40129	0.40517	0.40905	0.41294	0.41683	0.42074	-0.20
-0.10	0.42465	0.42858	0.43251	0.43644	0.44038	0.44433	0.44828	0.45224	0.45620	0.46017	-0.10
-0.00	0.46414	0.46812	0.47210	0.47608	0.48006	0.48405	0.48803	0.49202	0.49601	0.50000	-0.00

Standard Normal Table (continued)
Areas Under the Standard Normal Curve



z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09	z
0.00	0.50000	0.50399	0.50798	0.51197	0.51595	0.51994	0.52392	0.52790	0.53188	0.53586	0.00
0.10	0.53983	0.54380	0.54776	0.55172	0.55567	0.55962	0.56356	0.56749	0.57142	0.57535	0.10
0.20	0.57926	0.58317	0.58706	0.59095	0.59483	0.59871	0.60257	0.60642	0.61026	0.61409	0.20
0.30	0.61791	0.62172	0.62552	0.62930	0.63307	0.63683	0.64058	0.64431	0.64803	0.65173	0.30
0.40	0.65542	0.65910	0.66276	0.66640	0.67003	0.67364	0.67724	0.68082	0.68439	0.68793	0.40
0.50	0.69146	0.69497	0.69847	0.70194	0.70540	0.70884	0.71226	0.71566	0.71904	0.72240	0.50
0.60	0.72575	0.72907	0.73237	0.73565	0.73891	0.74215	0.74537	0.74857	0.75175	0.75490	0.60
0.70	0.75804	0.76115	0.76424	0.76730	0.77035	0.77337	0.77637	0.77935	0.78230	0.78524	0.70
0.80	0.78814	0.79103	0.79389	0.79673	0.79955	0.80234	0.80511	0.80785	0.81057	0.81327	0.80
0.90	0.81594	0.81859	0.82121	0.82381	0.82639	0.82894	0.83147	0.83398	0.83646	0.83891	0.90
1.00	0.84134	0.84375	0.84614	0.84849	0.85083	0.85314	0.85543	0.85769	0.85993	0.86214	1.00
1.10	0.86433	0.86650	0.86864	0.87076	0.87286	0.87493	0.87698	0.87900	0.88100	0.88298	1.10
1.20	0.88493	0.88686	0.88877	0.89065	0.89251	0.89435	0.89617	0.89796	0.89973	0.90147	1.20
1.30	0.90320	0.90490	0.90658	0.90824	0.90988	0.91149	0.91309	0.91466	0.91621	0.91774	1.30
1.40	0.91924	0.92073	0.92220	0.92364	0.92507	0.92647	0.92785	0.92922	0.93056	0.93189	1.40
1.50	0.93319	0.93448	0.93574	0.93699	0.93822	0.93943	0.94062	0.94179	0.94295	0.94408	1.50
1.60	0.94520	0.94630	0.94738	0.94845	0.94950	0.95053	0.95154	0.95254	0.95352	0.95449	1.60
1.70	0.95543	0.95637	0.95728	0.95818	0.95907	0.95994	0.96080	0.96164	0.96246	0.96327	1.70
1.80	0.96407	0.96485	0.96562	0.96638	0.96712	0.96784	0.96856	0.96926	0.96995	0.97062	1.80
1.90	0.97128	0.97193	0.97257	0.97320	0.97381	0.97441	0.97500	0.97558	0.97615	0.97670	1.90
2.00	0.97725	0.97778	0.97831	0.97882	0.97932	0.97982	0.98030	0.98077	0.98124	0.98169	2.00
2.10	0.98214	0.98257	0.98300	0.98341	0.98382	0.98422	0.98461	0.98500	0.98537	0.98574	2.10
2.20	0.98610	0.98645	0.98679	0.98713	0.98745	0.98778	0.98809	0.98840	0.98870	0.98899	2.20
2.30	0.98928	0.98956	0.98983	0.99010	0.99036	0.99061	0.99086	0.99111	0.99134	0.99158	2.30
2.40	0.99180	0.99202	0.99224	0.99245	0.99266	0.99286	0.99305	0.99324	0.99343	0.99361	2.40
2.50	0.99379	0.99396	0.99413	0.99430	0.99446	0.99461	0.99477	0.99492	0.99506	0.99520	2.50
2.60	0.99534	0.99547	0.99560	0.99573	0.99585	0.99598	0.99609	0.99621	0.99632	0.99643	2.60
2.70	0.99653	0.99664	0.99674	0.99683	0.99693	0.99702	0.99711	0.99720	0.99728	0.99736	2.70
2.80	0.99744	0.99752	0.99760	0.99767	0.99774	0.99781	0.99788	0.99795	0.99801	0.99807	2.80
2.90	0.99813	0.99819	0.99825	0.99831	0.99836	0.99841	0.99846	0.99851	0.99856	0.99861	2.90
3.00	0.99865	0.99869	0.99874	0.99878	0.99882	0.99886	0.99889	0.99893	0.99896	0.9990	3.00
3.10	0.99903	0.99906	0.99910	0.99913	0.99916	0.99918	0.99921	0.99924	0.99926	0.99929	3.10
3.20	0.99931	0.99934	0.99936	0.99938	0.99940	0.99942	0.99944	0.99946	0.99948	0.99950	3.20
3.30	0.99952	0.99953	0.99955	0.99957	0.99958	0.99960	0.99961	0.99962	0.99964	0.99965	3.30
3.40	0.99966	0.99968	0.99969	0.99970	0.99971	0.99972	0.99973	0.99974	0.99975	0.99976	3.40
3.50	0.99977	0.99978	0.99978	0.99979	0.99980	0.99981	0.99981	0.99982	0.99983	0.99983	3.50


4.8 The t-Distribution:

- Student's t distribution.
- t-distribution is a **distribution of a continuous random variable**.
- t-distribution with $(n-1)$ degrees of freedom ($df = v = n - 1$), and we write :


$$T \sim t(v) \quad \text{or} \quad T \sim t(n - 1)$$

Note : (Characteristics of the t-Distribution)

- t-distribution is a continuous distribution.
- The value of t random variable range from $-\infty$ to $+\infty$ (that is , $-\infty < t < \infty$)
- The mean of t distribution is 0.
- It is symmetric about the mean 0.
- The shape of t-distribution is similar to the shape of the standard normal distribution.
- t-distribution $\longrightarrow$ Standard normal distribution as $n \longrightarrow \infty$.



Notation (t_α) :



- t_α = The t-value under which we find an area equal to α
= The t-value that leaves an area of α to the left.
- The value t_α satisfies: $P(T < t_\alpha) = \alpha$.

$$t_\alpha \gg P(T < t_\alpha) = \alpha$$

• Since the curve of the pdf of $T \sim t(v)$ is symmetric about 0, we have

$$t_{\alpha} = -t_{1-\alpha}$$

for example $t_{0.1} = -t_{1-0.1} = -t_{0.9}$
 $t_{0.975} = -t_{1-0.975} = -t_{0.025}$

• Values of t_{α} are tabulated in a special table for several values of α and several values of degrees of freedom (df=v =n-1).

degrees of freedom

Critical Values of the t-distribution (t_{α})



t_{α}

v=df	$t_{0.90}$	$t_{0.95}$	$t_{0.975}$	$t_{0.99}$	$t_{0.995}$
1	3.078	1.638	1.960	3.182	3.078
2	1.886	1.638	1.960	2.920	1.886
3	1.638	1.638	1.960	2.353	1.638
4	1.533	1.638	1.960	2.132	1.533
5	1.476	1.638	1.960	2.015	1.476
6	1.440	1.638	1.960	1.943	1.440
7	1.415	1.638	1.960	1.895	1.415
8	1.397	1.638	1.960	1.860	1.397
9	1.383	1.638	1.960	1.833	1.383
10	1.372	1.638	1.960	1.812	1.372
11	1.363	1.638	1.960	1.796	1.363
12	1.356	1.638	1.960	1.782	1.356
13	1.350	1.638	1.960	1.771	1.350
14	1.345	1.638	1.960	1.761	1.345
15	1.341	1.638	1.960	1.753	1.341

Example:

Find the t-value with $v=14$ (df) that leaves an area of:


- (a) 0.95 to the left.
- (b) 0.95 to the right.

Solution:

$v = 14$ (df) ; $T \sim t(14)$

(a) The t-value that leaves an area of **0.95 to the left** is:

$$t_{0.95} = 1.761$$




Critical Values of the t-distribution (t_α)

$v=df$	$t_{0.90}$	$t_{0.95}$	$t_{0.975}$	$t_{0.99}$	$t_{0.995}$
1	3.078	6.314	12.706	31.821	63.657
2	1.886	2.920	4.303	6.965	9.925
3	1.638	2.353	3.182	4.541	5.841
4	1.533	2.132	2.776	3.747	4.604
5	1.476	2.015	2.571	3.365	4.032
6	1.440	1.943	2.447	3.143	3.707
7	1.415	1.895	2.365	2.998	3.499
8	1.397	1.860	2.306	2.896	3.355
9	1.383	1.833	2.262	2.821	3.250
10	1.372	1.812	2.228	2.764	3.169
11	1.363	1.796	2.201	2.718	3.106
12	1.356	1.782	2.179	2.681	3.055
13	1.350	1.771	2.160	2.650	3.012
14	1.345	1.761	2.145	2.624	2.977
15	1.341	1.753	2.131	2.602	2.947
16	1.337	1.746	2.120	2.583	2.921
17	1.333	1.740	2.110	2.567	2.898
18	1.330	1.734	2.101	2.552	2.878
19	1.328	1.729	2.093	2.539	2.861
20	1.325	1.725	2.086	2.528	2.845
21	1.323	1.721	2.080	2.518	2.831
22	1.321	1.717	2.074	2.508	2.819
23	1.319	1.714	2.069	2.500	2.807
24	1.318	1.711	2.064	2.492	2.797
25	1.316	1.708	2.060	2.485	2.787



(b) The t-value that leaves an area of 0.95 to the right is :

$$t_{0.05} = -t_{1-0.05} = -t_{0.95} = -1.761$$



Critical Values of the t-distribution (t_{α})

v=df	$t_{0.90}$	$t_{0.95}$	$t_{0.975}$	$t_{0.99}$	$t_{0.995}$
1	3.078	6.314	12.706	31.821	63.657
2	1.886	2.920	4.303	6.965	9.925
3	1.638	2.353	3.182	4.541	5.841
4	1.533	2.132	2.776	3.747	4.604
5	1.476	2.015	2.571	3.365	4.032
6	1.440	1.943	2.447	3.143	3.707
7	1.415	1.895	2.365	2.998	3.499
8	1.397	1.860	2.306	2.896	3.355
9	1.383	1.833	2.262	2.821	3.250
10	1.372	1.812	2.228	2.764	3.169
11	1.363	1.796	2.201	2.718	3.106
12	1.356	1.782	2.179	2.681	3.055
13	1.350	1.771	2.160	2.650	3.012
14	1.345	1.761	2.145	2.624	2.977
15	1.341	1.753	2.131	2.602	2.947
16	1.337	1.746	2.120	2.583	2.921
17	1.333	1.740	2.110	2.567	2.898
18	1.330	1.734	2.101	2.552	2.878
19	1.328	1.729	2.093	2.539	2.861
20	1.325	1.725	2.086	2.528	2.845
21	1.323	1.721	2.080	2.518	2.831
22	1.321	1.717	2.074	2.508	2.819
23	1.319	1.714	2.069	2.500	2.807
24	1.318	1.711	2.064	2.492	2.797
25	1.316	1.708	2.060	2.485	2.787



Note:

Some t-tables contain values of α that are greater than or equal to 0.90. When we search for small values of α in these tables, we may use the fact that:

$$t_{\alpha} = -t_{1-\alpha}$$

Example:

For $\nu = 10$ degrees of freedom (df), find $t_{0.93}$ and $t_{0.07}$

Solution

(from the table)

$$t_{0.93} = \frac{(1.372+1.812)}{2} = 1.592$$

(using the rule: $t_{\alpha} = -t_{1-\alpha}$)

$$t_{0.07} = -t_{1-0.07} = -t_{0.93} = -1.592$$


Critical Values of the t-distribution (t_{α})

$\nu=df$	$t_{0.90}$	$t_{0.95}$	$t_{0.975}$	$t_{0.99}$	$t_{0.995}$
1	3.078	6.314	12.706	31.821	63.657
2	1.886	2.920	4.303	6.965	9.925
3	1.638	2.353	3.182	4.541	5.841
4	1.533	2.132	2.776	3.747	4.604
5	1.476	2.015	2.571	3.365	4.032
6	1.440	1.943	2.447	3.143	3.707
7	1.415	1.895	2.365	2.998	3.499
8	1.397	1.860	2.306	2.896	3.355
9	1.383	1.833	2.262	2.821	3.250
10	1.372	1.812	2.228	2.764	3.169
11	1.363	1.796	2.201	2.718	3.106
12	1.356	1.782	2.179	2.681	3.055
13	1.350	1.771	2.160	2.650	3.012




The t-value that leaves an area of 0.975 to the left (use $\nu = 12$) is :

$$t_{0.975} = 2.179$$




Critical Values of the t-distribution (t_{α})



$\nu=df$	$t_{0.90}$	$t_{0.95}$	$t_{0.975}$	$t_{0.99}$	$t_{0.995}$
1	3.078	6.314	12.706	31.821	63.657
2	1.886	2.920	4.303	6.965	9.925
3	1.638	2.353	3.182	4.541	5.841
4	1.533	2.132	2.776	3.747	4.604
5	1.476	2.015	2.571	3.365	4.032
6	1.440	1.943	2.447	3.143	3.707
7	1.415	1.895	2.365	2.998	3.499
8	1.397	1.860	2.306	2.896	3.355
9	1.383	1.833	2.262	2.821	3.250
10	1.372	1.812	2.228	2.764	3.169
11	1.363	1.796	2.201	2.718	3.106
12	1.356	1.782	2.179	2.681	3.055
13	1.350	1.771	2.160	2.650	3.012

The t-value that leaves an area of 0.90 to the **right** (use $\nu = 16$) is :

$$t_{0.10} = - t_{1-0.10} = - t_{0.90} = -1.337$$




Critical Values of the t-distribution (t_{α})



$\nu=df$	$t_{0.90}$	$t_{0.95}$	$t_{0.975}$	$t_{0.99}$	$t_{0.995}$
1	3.078	6.314	12.706	31.821	63.657
2	1.886	2.920	4.303	6.965	9.925
3	1.638	2.353	3.182	4.541	5.841
4	1.533	2.132	2.776	3.747	4.604
5	1.476	2.015	2.571	3.365	4.032
6	1.440	1.943	2.447	3.143	3.707
7	1.415	1.895	2.365	2.998	3.499
8	1.397	1.860	2.306	2.896	3.355
9	1.383	1.833	2.262	2.821	3.250
10	1.372	1.812	2.228	2.764	3.169
11	1.363	1.796	2.201	2.718	3.106
12	1.356	1.782	2.179	2.681	3.055
13	1.350	1.771	2.160	2.650	3.012
14	1.345	1.761	2.145	2.624	2.977
15	1.341	1.753	2.131	2.602	2.947
16	1.337	1.746	2.120	2.583	2.921
17	1.333	1.740	2.110	2.567	2.898

- The t-value that leaves an area of 0.025 to the right (use $v = 8$) is :
 $t_{0.975} = 2.306$



Critical Values of the t-distribution (t_{α})

$v=df$	$t_{0.90}$	$t_{0.95}$	$t_{0.975}$	$t_{0.99}$	$t_{0.995}$
1	3.078	6.314	12.706	31.821	63.657
2	1.886	2.920	4.303	6.965	9.925
3	1.638	2.353	3.182	4.541	5.841
4	1.533	2.132	2.776	3.747	4.604
5	1.476	2.015	2.571	3.365	4.032
6	1.440	1.943	2.447	3.143	3.707
7	1.415	1.895	2.365	2.998	3.499
8	1.397	1.858	2.306	2.896	3.355
9	1.383	1.833	2.262	2.821	3.250
10	1.372	1.812	2.228	2.764	3.169



Examples

- The t-value that leaves an area of 0.025 to the left (use $\nu = 8$) is :

$$t_{0.025} = - t_{1-0.025} = - t_{0.975} = -2.306$$

- The t-value that leaves an area of 0.93 to the left (use $\nu = 10$) is :


$$t_{0.93} = \frac{t_{0.90} + t_{0.95}}{2} = \frac{1.372 + 1.812}{2} = 1.592$$

- The t-value that leaves an area of 0.07 to the left (use $\nu = 10$) is :

$$t_{0.70} = - t_{0.93} = - \left(\frac{t_{0.90} + t_{0.95}}{2} \right) = -1.592$$

- $P(T \leq K) = 0.90, df = 10$
- $P(T \leq K) = 0.90 \gg t_{0.90}$
- $K=1.375$

Critical Values of the t-distribution (t_{α})



v=df	$t_{0.90}$	$t_{0.95}$	$t_{0.975}$	$t_{0.99}$	$t_{0.995}$
1	3.078	6.314	12.706	31.821	63.657
2	1.886	2.920	4.303	6.965	9.925
3	1.638	2.353	3.182	4.541	5.841
4	1.533	2.132	2.776	3.747	4.604
5	1.476	2.015	2.571	3.365	4.032
6	1.440	1.943	2.447	3.143	3.707
7	1.415	1.895	2.365	2.998	3.499
8	1.397	1.860	2.306	2.896	3.355
9	1.383	1.833	2.262	2.821	3.250
10	1.372		2.228	2.764	3.169
11	1.363		2.201	2.718	3.106

Value of t

- $P(T \geq K) = 0.95$, $df = 15$
- $P(T \geq K) = 0.95 \gg P(T \leq K) = 1 - 0.95$
- $\gg P(T \leq K) = 0.05$
- $\gg t_{0.05} = -t_{1-0.05}$
- $\gg -t_{0.95}$

$K = -1.753$

Critical Values of the t-distribution (t_{α})




v=df	$t_{0.90}$	$t_{0.95}$	$t_{0.975}$	$t_{0.99}$	$t_{0.995}$
1	3.078	6.314	12.706	31.821	63.657
2	1.886	2.920	4.303	6.965	9.925
3	1.638	2.353	3.182	4.541	5.841
4	1.533	2.132	2.776	3.747	4.604
5	1.476	2.015	2.571	3.365	4.032
6	1.440	1.943	2.447	3.143	3.707
7	1.415	1.895	2.365	2.998	3.499
8	1.397	1.860	2.306	2.896	3.355
9	1.383	1.833	2.262	2.821	3.250
10	1.372	1.812	2.228	2.764	3.169
11	1.363	1.796	2.201	2.718	3.106
12	1.356	1.782	2.179	2.681	3.055
13	1.350	1.771	2.160	2.650	3.012
14	1.345	1.761	2.145	2.624	2.977
15	1.341	1.753	2.131	2.602	2.947
16	1.337	1.746	2.120	2.583	2.921
17	1.333	1.740	2.110	2.567	2.898

$$P(T \leq 2.110) = ? , df=17$$

$$P(T \leq 2.110) = 0.975$$

Critical Values of the t-distribution (t_{α})



v=df	$t_{0.90}$	$t_{0.95}$	$t_{0.975}$	$t_{0.99}$	$t_{0.995}$
1	3.078	6.314	1.996	31.821	63.657
2	1.886	2.920	1.960	6.965	9.925
3	1.638	2.353	1.932	4.541	5.841
4	1.533	2.132	1.915	3.747	4.604
5	1.476	2.015	1.901	3.365	4.032
6	1.440	1.943	1.897	3.143	3.707
7	1.415	1.895	1.894	2.998	3.499
8	1.397	1.860	1.891	2.896	3.355
9	1.383	1.833	1.888	2.821	3.250
10	1.372	1.812	1.885	2.764	3.169
11	1.363	1.796	1.882	2.718	3.106
12	1.356	1.782	1.879	2.681	3.055
13	1.350	1.771	1.876	2.650	3.012
14	1.345	1.761	1.873	2.624	2.977
15	1.341	1.753	1.871	2.602	2.947
16	1.337	1.746	1.868	2.583	2.921
17	1.333	1.740	2.110	2.567	2.898

$$P(T \leq 2.718) = ? , df=11$$


$$P(T \leq 2.718) = 0.99$$

Critical Values of the t-distribution (t_{α})




v=df	$t_{0.90}$	$t_{0.95}$	$t_{0.975}$	$t_{0.99}$	$t_{0.995}$
1	3.078	6.314	12.706	31.821	63.657
2	1.886	2.920	4.303	6.965	9.925
3	1.638	2.353	3.182	4.541	5.841
4	1.533	2.132	2.776	3.747	4.604
5	1.476	2.015	2.571	3.365	4.032
6	1.440	1.943	2.447	3.143	3.707
7	1.415	1.895	2.365	2.998	3.499
8	1.397	1.860	2.306	2.896	3.355
9	1.383	1.833	2.262	2.821	3.250
10	1.372	1.812	2.228	2.764	3.169
11	1.363	1.796	2.201	2.718	3.106
12	1.356	1.782	2.179	2.681	3.055
13	1.350	1.771	2.160	2.650	3.012

Critical Values of the t-distribution (t_{α})



v=df	$t_{0.90}$	$t_{0.95}$	$t_{0.975}$	$t_{0.99}$	$t_{0.995}$
1	3.078	6.314	12.706	31.821	63.657
2	1.886	2.920	4.303	6.965	9.925
3	1.638	2.353	3.182	4.541	5.841
4	1.533	2.132	2.776	3.747	4.604
5	1.476	2.015	2.571	3.365	4.032
6	1.440	1.943	2.447	3.143	3.707
7	1.415	1.895	2.365	2.998	3.499
8	1.397	1.860	2.306	2.896	3.355
9	1.383	1.833	2.262	2.821	3.250
10	1.372	1.812	2.228	2.764	3.169
11	1.363	1.796	2.201	2.718	3.106
12	1.356	1.782	2.179	2.681	3.055
13	1.350	1.771	2.160	2.650	3.012
14	1.345	1.761	2.145	2.624	2.977
15	1.341	1.753	2.131	2.602	2.947
16	1.337	1.746	2.120	2.583	2.921
17	1.333	1.740	2.110	2.567	2.898
18	1.330	1.734	2.101	2.552	2.878
19	1.328	1.729	2.093	2.539	2.861
20	1.325	1.725	2.086	2.528	2.845
21	1.323	1.721	2.080	2.518	2.831
22	1.321	1.717	2.074	2.508	2.819
23	1.319	1.714	2.069	2.500	2.807
24	1.318	1.711	2.064	2.492	2.797
25	1.316	1.708	2.060	2.485	2.787

Critical Values of the t -distribution (t_α)



$v=df$	$t_{0.90}$	$t_{0.95}$	$t_{0.975}$	$t_{0.99}$	$t_{0.995}$
26	1.315	1.706	2.056	2.479	2.779
27	1.314	1.703	2.052	2.473	2.771
28	1.313	1.701	2.048	2.467	2.763
29	1.311	1.699	2.045	2.462	2.756
30	1.310	1.697	2.042	2.457	2.750
35	1.3062	1.6896	2.0301	2.4377	2.7238
40	1.3030	1.6840	2.0210	2.4230	2.7040
45	1.3006	1.6794	2.0141	2.4121	2.6896
50	1.2987	1.6759	2.0086	2.4033	2.6778
60	1.2958	1.6706	2.0003	2.3901	2.6603
70	1.2938	1.6669	1.9944	2.3808	2.6479
80	1.2922	1.6641	1.9901	2.3739	2.6387
90	1.2910	1.6620	1.9867	2.3685	2.6316
100	1.2901	1.6602	1.9840	2.3642	2.6259
120	1.2886	1.6577	1.9799	2.3578	2.6174
140	1.2876	1.6558	1.9771	2.3533	2.6114
160	1.2869	1.6544	1.9749	2.3499	2.6069
180	1.2863	1.6534	1.9732	2.3472	2.6034
200	1.2858	1.6525	1.9719	2.3451	2.6006
∞	1.282	1.645	1.960	2.326	2.576

Chapter 5:

Probabilistic Features of the Distributions of Certain Sample Statistics

5.1 Introduction

5.2 sampling Distribution

5.3 Distribution of the Sample Mean $\bar{X}$

5.4 Distribution of the Difference Between Two Sample Means $\bar{X}_1 - \bar{X}_2$

5.5 Distribution of the Sample Proportion $\hat{P}$

5.6 Distributions of the difference between two sample proportions $\hat{P}_1 - \hat{P}_2$

Introduction

In this Chapter we will discuss the probability distributions of some statistics.

As we mention earlier, a statistic is a measure computed from the random sample. As the sample values vary from sample to sample, the value of the statistic varies accordingly.

A statistic is a random variable; it has a probability distribution, a mean and a variance.

	Sample	Population
Size	n	N
Mean	$\bar{X}$	μ
Variance	S^2	σ^2
Standard deviation	S	σ
Proportion	$\hat{P}$	P
Difference between two means	$\bar{X}_1 - \bar{X}_2$	$\mu_1 - \mu_2$
Difference between two proportions	$\hat{P}_1 - \hat{P}_2$	$P_1 - P_2$

Statistics

Parameter

5.2 sampling Distribution :

The probability distribution of a statistic is called the sampling distribution of that statistic.

The sampling distribution of the statistic is used to make statistical inference about the unknown parameter.

5.3 Distribution of the Sample Mean ($\bar{X}$) :

Suppose that we have a population with mean μ and variance σ^2 .

Suppose that $x_1, x_2, \dots, x_n$ is a random sample of size (n) selected randomly from this population. We know that the sample mean is:

$$\bar{X} = \frac{\sum_{i=1}^n x_n}{n}$$

Suppose that we select several random samples of size $n = 5$:

	1 st sample	2 nd sample	3 rd sample	...	Last sample
Sample value	28	31	14	...	17
	30	20	31	...	32
	34	31	25	...	29
	34	40	27	...	31
	17	28	32	...	30
Sample Mean $\bar{x}$	28.4	29.9	25.8	...	27.8

- The value of the sample mean $\bar{X}$ varies from random sample to another.
- The value of $\bar{X}$ is random and it depends on the random sample.
- The sample mean $\bar{X}$ is a random variable.
- The probability distribution of $\bar{X}$ is called the sampling distribution of the sample mean $\bar{X}$.

•Questions :

- What is the sampling distribution of the sample mean $\bar{X}$?
- What is the mean of the sample mean $\bar{X}$?
- What is the variance of the sample mean $\bar{X}$?

Result (1) : (Mean & variance of $\bar{X}$)

If $x_1, x_2, \dots, x_n$ is a random sample of size (n) from any distribution with mean μ and variance σ^2 , then:

1. The mean of $\bar{X}$ is : $\mu_{\bar{X}} = \mu$
2. The variance of $\bar{X}$ is : $\sigma_{\bar{X}}^2 = \frac{\sigma^2}{n}$
3. The standard deviation of $\bar{X}$ is called the standard error and is defined by : $\sigma_{\bar{X}} = \sqrt{\sigma_{\bar{X}}^2} = \frac{\sigma}{\sqrt{n}}$

Result (2): (Sampling from normal population)

If $x_1, x_2, \dots, x_n$ is a random sample of size (n) from a normal population with mean μ and variance σ^2 , that is $\text{Normal}(\mu, \sigma^2)$, then the sample mean $\bar{X}$ has a normal distribution with mean μ and variance $\frac{\sigma^2}{n}$, that is :

1. $\bar{X} \sim \text{Normal}(\mu, \frac{\sigma^2}{n})$
2. $Z = \frac{\bar{X} - \mu}{\frac{\sigma}{\sqrt{n}}} \sim \text{Normal}(0, 1)$

We use this result when sampling from **normal distribution with known variance σ^2** .

Result (3): (Central Limit Theorem: Sampling from Non-normal population)

Suppose that $x_1, x_2, \dots, x_n$ is a random sample of size (n) from a non-normal population with mean μ and variance σ^2 . if the sample size n is large ($n \geq 30$), then the sample mean $\bar{X}$ has approximately a normal distribution with mean μ and variance $\frac{\sigma^2}{n}$, that is :

1. $\bar{X} \approx \text{Normal}(\mu, \frac{\sigma^2}{n})$ (approximately)
2. $Z = \frac{\bar{X} - \mu}{\frac{\sigma}{\sqrt{n}}} \approx \text{Normal}(0, 1)$ (approximately)

Notes :

- “ $\approx$ ” means “approximately distributed”.
- We use this result when sampling from non-normal distribution with known variance σ^2 and with large sample size ($n \geq 30$).

Result (4): (used when σ^2 is unknown + normal distribution)

If $x_1, x_2, \dots, x_n$ is a random sample of size (n) from a normal distribution with mean μ and variance σ^2 , that is $\text{Normal}(\mu, \sigma^2)$, then the statistic :

$$1. \quad \bar{X} \sim t(n - 1)$$

$$2. \quad T = \frac{\bar{X} - \mu}{S/\sqrt{n}} \sim t(n-1)$$

Notation : degrees of freedom = $df = v = n - 1$

We use this result when sampling from normal distribution with Unknown variance σ^2 and with small sample size ($n < 30$).

Application:

Example: (Sampling distribution of the sample mean)

Suppose that the time duration of a minor surgery is approximately normally distributed with mean equal to 800 seconds and a standard deviation of 40 seconds. Find the probability that a random sample of 16 surgeries will have average time duration of less than 775 seconds.

Solution:

X = the duration of the surgery

$$\mu = 800, \sigma = 40, \sigma^2 = 40^2 = 1600, n = 16$$

$$X \sim \text{Normal}(\mu = 800, \sigma^2 = 1600)$$

Calculating mean, variance, and standard error (standard deviation) of the sample mean $\bar{X}$:

The mean of $\bar{X}$: $\mu_{\bar{X}} = \mu = 800$

The variance of $\bar{X}$: $\sigma_{\bar{X}}^2 = \frac{\sigma^2}{n} = \frac{1600}{16} = 100$

The standard deviation of $\bar{X}$: $\sigma_{\bar{X}} = \sqrt{\sigma_{\bar{X}}^2} = \frac{\sigma}{\sqrt{n}} = \frac{40}{\sqrt{16}} = 10$

The sampling distribution of $\bar{X}$ is :

mean $\mu_{\bar{X}} = 800$ and variance $\sigma_{\bar{X}}^2 = 100$. That is :

$$\bar{X} \sim \text{Normal}(800, 100)$$

$$Z = \frac{\bar{X} - \mu}{\frac{\sigma}{\sqrt{n}}} = \frac{\bar{X} - 800}{10} \sim \text{Normal}(0, 1)$$

Result (2):
normal distribution with
known variance σ^2

The probability that a random sample of 16 surgeries will have average time duration of less than 775 seconds equals to :


$$P(\bar{X} < 775) = P\left(\frac{\bar{X} - 800}{10} < \frac{775 - 800}{10}\right) = P(Z < -2.5) = 0.00621$$

$$\bar{X} \sim N\left(\mu, \frac{\sigma^2}{n}\right) = N(800, 100)$$

$$\mu_{\bar{X}} = 800$$

$$\sigma_{\bar{X}}^2 = \frac{\sigma^2}{n} = 100$$

$$\sigma_{\bar{X}} = \frac{\sigma}{\sqrt{n}} = 10$$



Example :

If the mean and standard deviation of serum iron values for healthy men are 120 and 15 microgram/100ml, respectively, what is the probability that a random sample of size 50 normal men will yield a mean between 115 and 125 microgram/100ml ?

Solution :

X = the serum iron value

$$\mu = 120, \sigma = 15, \sigma^2 = 15^2 = 225, n = 50$$

$$X \sim \text{Normal} (\mu = 120, \sigma^2 = 225)$$

Calculating mean, variance, and standard error (standard deviation) of the sample mean $\bar{X}$:

The mean of $\bar{X}$: $\mu_{\bar{X}} = \mu = 120$

The variance of $\bar{X}$: $\sigma_{\bar{X}}^2 = \frac{\sigma^2}{n} = \frac{225}{50} = 4.5$

The standard deviation of $\bar{X}$: $\sigma_{\bar{X}} = \sqrt{\sigma_{\bar{X}}^2} = \frac{\sigma}{\sqrt{n}} = \frac{15}{\sqrt{50}} = 2.1213$

Using the central limit theorem $\bar{X}$ has a normal distribution

with mean $\mu_{\bar{X}} = 120$ and variance $\sigma_{\bar{X}}^2 = 4.5$

That is :

$$\bar{X} \sim \text{Normal}(120, 4.5)$$

$$Z = \frac{\bar{X} - \mu}{\frac{\sigma}{\sqrt{n}}} = \frac{\bar{X} - 120}{2.12} \sim \text{Normal}(0, 1)$$

Result (3):
non-normal distribution with
known variance σ^2 and with
large sample size ($n \geq 30$).

The probability that a random sample of size 50 men will yield a mean between 115 and 125 microgram/100ml equals to :


$$\begin{aligned} P(115 < \bar{X} < 125) &= P\left(\frac{115-120}{2.1213} < \frac{\bar{X}-120}{2.12} < \frac{125-120}{2.1213}\right) \\ &= P(-2.36 < Z < 2.36) \\ &= P(Z < 2.36) - P(Z < -2.36) \\ &= 0.99086 - 0.00914 \\ &= 0.98172 \end{aligned}$$

5.4 Distribution of the Difference Between Two Sample Means ($\bar{X}_1 - \bar{X}_2$) :

Suppose that we have two populations:

- 1-st population with mean μ_1 and variance σ_1^2 .
- 2-nd population with mean μ_2 and variance σ_2^2 .
- We are interested in comparing m_1 and m_2 , or equivalently, making inferences about the difference between the means ($\mu_1 - \mu_2$) .
- We independently select a random sample of size n_1 from the 1-st population and another random sample of size n_2 from the 2-nd population .

- Let $\bar{X}_1$ and S_1^2 be the sample mean and the sample variance of the 1-st sample .
- Let $\bar{X}_2$ and S_2^2 be the sample mean and the sample variance of the 2-nd sample
- The sampling distribution of $\bar{X}_1 - \bar{X}_2$ is used to make inferences about $\mu_1 - \mu_2$.



The sampling distribution of $\bar{X}_1 - \bar{X}_2$:

Result (1) : The mean , variance and the standard deviation of $\bar{X}_1 - \bar{X}_2$ are :

1. The mean of $\bar{X}_1 - \bar{X}_2$ is : $\mu_{\bar{X}_1 - \bar{X}_2} = \mu_1 - \mu_2$
2. The variance of $\bar{X}_1 - \bar{X}_2$ is : $\sigma_{\bar{X}_1 - \bar{X}_2}^2 = \frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}$
3. The standard deviation of $\bar{X}_1 - \bar{X}_2$ is called the standard error and is defined by : $\sigma_{\bar{X}_1 - \bar{X}_2} = \sqrt{\sigma_{\bar{X}_1 - \bar{X}_2}^2} = \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}$

• **Note:** Square roots distribute over multiplication or division, but not addition or subtraction.

$$\sqrt{a + b} \neq \sqrt{a} + \sqrt{b}$$

• **In general:** $Z = \frac{\text{value} - \text{Mean}}{\text{Standard deviation}}$

Result (2) :

If the two random samples were selected from normal distributions (or non-normal distributions with large sample sizes) with known variances σ_1^2 and σ_2^2 , then the difference between the sample means $\bar{X}_1 - \bar{X}_2$ has a normal distribution with mean $\mu_1 - \mu_2$ and variance $\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}$, that is :

$$1. \quad \bar{X}_1 - \bar{X}_2 \sim \text{Normal} (\mu_1 - \mu_2, \frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2})$$

$$2. \quad Z = \frac{(\bar{X}_1 - \bar{X}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}} \sim \text{Normal} (0,1)$$

We use this result when sampling :

1-from normal distribution with known variance σ_1^2, σ_2^2 .

2-from non-normal distribution with known variance σ_1^2, σ_2^2 ,and with large sample size ($n_1, n_2 \geq 30$).

Example:

Suppose it has been established that for a certain type of client (type A) the average length of a home visit by a public health nurse is 45 minutes with standard deviation of 15 minutes, and that for second type (type B) of client the average home visit is 30 minutes long with standard deviation of 20 minutes. If a nurse randomly visits 35 clients from the first type and 40 clients from the second type, what is the probability that the average length of home visit of first type will be greater than the average length of home visit of second type by 20 or more minutes ?

Solution :

Type	Mean (Average)	Standard deviation	Variance	Sample size
A (First type)	$\mu_1=45$	$\sigma_1=15$	$\sigma_1^2=225$	$n_1=35$ (large)
B (Second type)	$\mu_2=30$	$\sigma_2=20$	$\sigma_2^2=400$	$n_2=40$ (large)

The mean, the variance and the standard deviation of $\bar{X}_1 - \bar{X}_2$ are:

1. The mean of $\bar{X}_1 - \bar{X}_2$ is : $\mu_{\bar{X}_1 - \bar{X}_2} = \mu_1 - \mu_2 = 45 - 30 = 15$
2. The variance of $\bar{X}_1 - \bar{X}_2$ is : $\sigma_{\bar{X}_1 - \bar{X}_2}^2 = \frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2} = \frac{225}{35} + \frac{400}{40} = 16.4286$
3. The standard deviation of $\bar{X}_1 - \bar{X}_2$ is called the standard error and is defined by : $\sigma_{\bar{X}_1 - \bar{X}_2} = \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}} = \sqrt{16.4286} = 4.0532$

The sampling distribution of $\bar{X}_1 - \bar{X}_2$ is :

$$\bar{X}_1 - \bar{X}_2 \sim \text{Normal} (15 , 16.4286)$$

$$Z = \frac{(\bar{X}_1 - \bar{X}_2) - 15}{\sqrt{16.4286}} \sim \text{Normal} (0,1)$$


Result :
non-normal distribution with
known variance and with
large sample size .

the probability that the average length of home visit of first type will be greater than the average length of home visit of second type by 20 or more minutes equals to :

$$\begin{aligned} P(\bar{X}_1 > \bar{X}_2 + 20) &= \\ &= P(\bar{X}_1 - \bar{X}_2 > 20) \\ &= P\left(Z > \frac{20-15}{4.0532} \right) \\ &= P(Z > 1.23) \\ &= 1 - P(Z < 1.23) \\ &= 1 - 0.89065 \\ &= 0.10935 \end{aligned}$$

5.5 Distribution of the Sample Proportion ($\hat{P}$):

The sampling distribution of $\hat{P}$ is used to make inferences about p .



- **For the population:**

$N(A)$ = number of elements in the population with a specified characteristic "A"

N = total number of elements in the population (population size)

The population proportion is :

$$P = \frac{N(A)}{N} \quad (P \text{ is a parameter})$$

- **For the sample:**

$n(A)$ = number of elements in the sample with the same characteristic "A"

n = sample size

The sample proportion is :

$$\hat{P} = \frac{n(A)}{n} \quad (\hat{P} \text{ is a statistic})$$

Result (1):

- The mean of the sample proportion $\hat{P}$ is the population proportion (P) , that is: $\mu_{\hat{P}} = P$
- The variance of the sample proportion $\hat{P}$ is : $\sigma_{\hat{P}}^2 = \frac{P(1-P)}{n} = \frac{pq}{n}$ (where $q=1-p$)

The standard error (standard deviation) of the sample proportion $\hat{P}$ is : $\sigma_{\hat{P}} = \sqrt{\frac{pq}{n}}$

Result (2):

- For large sample size ($n \geq 30, np > 5, nq > 5$), the sample proportion $\hat{P}$ has approximately a normal distribution with $\mu_{\hat{P}} = P$ and $\sigma_{\hat{P}}^2 = \frac{pq}{n}$, that is:

$$\hat{P} \approx \text{Normal} \left(P, \frac{pq}{n} \right) \quad (\text{approximately})$$

$$Z = \frac{\hat{P} - P}{\sqrt{\frac{pq}{n}}} \sim \text{Normal} (0,1)$$

Example:

Suppose that 45% of the patients visiting a certain clinic are females. If a sample of 35 patients was selected at random, find the probability that:

1. The proportion of females in the sample will be greater than 0.4.
2. The proportion of females in the sample will be between 0.4 and 0.5.

Solution:

$n = 35$ (large)

p = The population proportion of females = $45\% = \frac{45}{100} = 0.45$ ($q = 1 - p = 0.55$)

$\hat{P}$ = The sample proportion (proportion of females in the sample)

- The mean of the sample proportion $\hat{P}$ is : $P=0.45$
- The variance of the sample proportion $\hat{P}$ is : $\frac{pq}{n} = \frac{(0.45)(0.55)}{35} = 0.0071$
- The standard error (standard deviation) of the sample proportion $\hat{P}$ is : $\sqrt{\frac{pq}{n}} = \sqrt{0.0071} = 0.084$

- $n = 35 \geq 30$, $np = 35(0.45) = 15.75 > 5$, $nq = 35(0.55) = 19.25 > 5$

The distribution of $\hat{P}$ is: $\hat{P} \approx \text{Normal}(P=0.45, \frac{pq}{n} = 0.0071)$


- The probability that the proportion of females in the sample will be greater than 0.4 :

$$\begin{aligned} P(\hat{P} > 0.4) &= P(Z > \frac{0.4-0.45}{\sqrt{0.0071}}) \\ &= P(Z > -0.59) = 1 - P(Z < -0.59) \\ &= 1 - 0.2776 \\ &= 0.7224 \end{aligned}$$

- The probability that the proportion of females in the sample will be between 0.4 and 0.5 :

$$\begin{aligned} P(0.4 < \hat{P} < 0.5) &= P(\frac{0.4-0.45}{\sqrt{0.0071}} < Z < \frac{0.5-0.45}{\sqrt{0.0071}}) \\ &= P(-0.59 < Z < 0.59) \\ &= P(Z < 0.59) - P(Z < -0.59) \\ &= 0.7224 - 0.2776 \\ &= 0.4448 \end{aligned}$$

5.6 Distributions of the difference between two sample proportions ($\hat{P}_1 - \hat{P}_2$):



Suppose that we have two populations:

- P_1 = proportion of elements of type (A) in the 1-st population.
- P_2 = proportion of elements of type (A) in the 2-nd population.
- We are interested in comparing P_1 and P_2 , or equivalently, making inferences about $P_1 - P_2$.
- We independently select a random sample of size n_1 from the 1-st population and another random sample of size n_2 from the 2-nd population:
- Let X_1 = no. of elements of type (A) in the 1-st sample.
- Let X_2 = no. of elements of type (A) in the 2-nd sample.
- $\hat{P}_1 = \frac{X_1}{n_1}$ = sample proportion of the 1-st sample .
- $\hat{P}_2 = \frac{X_2}{n_2}$ = sample proportion of the 2-nd sample .
- The sampling distribution of $\hat{P}_1 - \hat{P}_2$ is used to make inferences about $P_1 - P_2$.

The sampling distribution of $\hat{P}_1 - \hat{P}_2$:

Result (1) :

The mean, the variance and the standard error (standard deviation) of $\hat{P}_1 - \hat{P}_2$ are :

1. The mean of $\hat{P}_1 - \hat{P}_2$: $\mu_{\hat{P}_1 - \hat{P}_2} = P_1 - P_2$

2. The variance of $\hat{P}_1 - \hat{P}_2$ is : $\sigma_{\hat{P}_1 - \hat{P}_2}^2 = \frac{p_1 q_1}{n_1} + \frac{p_2 q_2}{n_2}$

3. The standard error (standard deviation) of $\hat{P}_1 - \hat{P}_2$ is : $\sigma_{\hat{P}_1 - \hat{P}_2} = \sqrt{\frac{p_1 q_1}{n_1} + \frac{p_2 q_2}{n_2}}$

Where : $q_1 = 1 - p_1$ and $q_2 = 1 - p_2$.

Result (2) :

For large samples sizes ($n_1 \geq 30, n_2 \geq 30, n_1 p_1 > 5, n_1 q_1 > 5, n_2 p_2 > 5, n_2 q_2 > 5$) , we have :

$$\hat{P}_1 - \hat{P}_2 \approx \text{Normal} \left(P_1 - P_2, \frac{p_1 q_1}{n_1} + \frac{p_2 q_2}{n_2} \right) \quad (\text{approximately})$$

$$Z = \frac{(\hat{P}_1 - \hat{P}_2) - (P_1 - P_2)}{\sqrt{\frac{p_1 q_1}{n_1} + \frac{p_2 q_2}{n_2}}} \sim \text{Normal} (0,1)$$

Example :

Suppose that 40% of Non-Saudi residents have medical insurance and 30% of Saudi residents have medical insurance in a certain city. We have randomly and independently selected a sample of 130 Non-Saudi residents and another sample of 120 Saudi residents. What is the probability that the difference between the sample proportions, will be between 0.05 and 0.2 ?

P_1 = population proportion of non-Saudi with medical insurance

P_2 = population proportion of Saudi with medical insurance

$\hat{P}_1$ = sample proportion of non-Saudi with medical insurance

$\hat{P}_2$ = sample proportion of Saudi with medical insurance

Type	proportion	q=1-p	Sample size
Non-Saudi residents	$P_1=0.40$	$q_1=0.60$	$n_1=130$ (large)
Saudi residents	$P_2=0.30$	$q_2=0.70$	$n_2=120$ (large)

The mean of $\hat{P}_1 - \hat{P}_2$: $\mu_{\hat{P}_1 - \hat{P}_2} = P_1 - P_2 = 0.4 - 0.3 = 0.1$

The variance of $\hat{P}_1 - \hat{P}_2$: $\sigma_{\hat{P}_1 - \hat{P}_2}^2 = \frac{p_1 q_1}{n_1} + \frac{p_2 q_2}{n_2} = \frac{(0.4)(0.6)}{130} + \frac{(0.3)(0.7)}{120} = 0.0036$

The standard error (standard deviation) of $\hat{P}_1 - \hat{P}_2$: $\sigma_{\hat{P}_1 - \hat{P}_2} = \sqrt{\frac{p_1 q_1}{n_1} + \frac{p_2 q_2}{n_2}} = \sqrt{0.0036} = 0.0599$

The distribution of $\hat{P}_1 - \hat{P}_2$ is: $(n_1 \geq 30, n_2 \geq 30, n_1 p_1 > 5, n_1 q_1 > 5, n_2 p_2 > 5, n_2 q_2 > 5)$

$$\hat{P}_1 - \hat{P}_2 \approx \text{Normal}(0.1, 0.0036)$$

The probability that the difference between the sample proportions, will be between 0.05 and 0.2 is :

$$\begin{aligned} P(0.05 < \hat{P}_1 - \hat{P}_2 < 0.2) &= P\left(\frac{0.05 - (P_1 - P_2)}{\sqrt{\frac{p_1 q_1}{n_1} + \frac{p_2 q_2}{n_2}}} < Z < \frac{0.2 - (P_1 - P_2)}{\sqrt{\frac{p_1 q_1}{n_1} + \frac{p_2 q_2}{n_2}}} \right) \\ &= P\left(\frac{0.05 - 0.1}{0.0599} < Z < \frac{0.2 - 0.1}{0.0599} \right) \\ &= P(-0.83 < Z < 1.67) \\ &= P(Z < 1.67) - P(Z < -0.83) \\ &= 0.95254 - 0.20327 = 0.74927 \end{aligned}$$