

The Fundamental Theorem of Calculus

Bander Almutairi

King Saud University

15 Sept 2013

1 The Fundamental Theorem of Calculus

(i) This theorem is very useful in evaluating definite integral;

- (i) This theorem is very useful in evaluating definite integral;
- (ii) This theorem also tells us the relationship between derivative and definite integral.

- (i) This theorem is very useful in evaluating definite integral;
- (ii) This theorem also tells us the relationship between derivative and definite integral.

The Fundamental Theorem of Calculus:

- (i) This theorem is very useful in evaluating definite integral;
- (ii) This theorem also tells us the relationship between derivative and definite integral.

The Fundamental Theorem of Calculus: Suppose f is continuous on the closed interval $[a, b]$.

- (i) This theorem is very useful in evaluating definite integral;
- (ii) This theorem also tells us the relationship between derivative and definite integral.

The Fundamental Theorem of Calculus: Suppose f is continuous on the closed interval $[a, b]$.

1 If $G(x) = \int_a^x f(t)dt$, for every $x \in [a, b]$,

- (i) This theorem is very useful in evaluating definite integral;
- (ii) This theorem also tells us the relationship between derivative and definite integral.

The Fundamental Theorem of Calculus: Suppose f is continuous on the closed interval $[a, b]$.

- 1 If $G(x) = \int_a^x f(t)dt$, for every $x \in [a, b]$, then $G(x)$ is an **antiderivative** of f on $[a, b]$.

- (i) This theorem is very useful in evaluating definite integral;
- (ii) This theorem also tells us the relationship between derivative and definite integral.

The Fundamental Theorem of Calculus: Suppose f is continuous on the closed interval $[a, b]$.

- 1 If $G(x) = \int_a^x f(t)dt$, for every $x \in [a, b]$, then $G(x)$ is an **antiderivative** of f on $[a, b]$.
- 2 If $F(x)$ is any antiderivative of f on $[a, b]$,

- (i) This theorem is very useful in evaluating definite integral;
- (ii) This theorem also tells us the relationship between derivative and definite integral.

The Fundamental Theorem of Calculus: Suppose f is continuous on the closed interval $[a, b]$.

- 1 If $G(x) = \int_a^x f(t)dt$, for every $x \in [a, b]$, then $G(x)$ is an **antiderivative** of f on $[a, b]$.
- 2 If $F(x)$ is any antiderivative of f on $[a, b]$, then

$$\int_a^b f(x)dx = F(b) - F(a).$$

Theorem (1): $\int_a^b f(x)dx = \left[\int f(x)dx \right]_a^b.$

Theorem (1): $\int_a^b f(x)dx = \left[\int f(x)dx \right]_a^b$.

Theorem (2): Let f be a continuous on $[a, b]$.

Theorem (1): $\int_a^b f(x)dx = \left[\int f(x)dx \right]_a^b$.

Theorem (2): Let f be a continuous on $[a, b]$. If
 $G(x) = \int_c^x f(t)dt$, where $a \leq c \leq b$,

Theorem (1): $\int_a^b f(x)dx = \left[\int f(x)dx \right]_a^b$.

Theorem (2): Let f be a continuous on $[a, b]$. If $G(x) = \int_c^x f(t)dt$, where $a \leq c \leq b$, then for every $x \in [a, b]$,

Theorem (1): $\int_a^b f(x)dx = \left[\int f(x)dx \right]_a^b$.

Theorem (2): Let f be a continuous on $[a, b]$. If $G(x) = \int_c^x f(t)dt$, where $a \leq c \leq b$, then for every $x \in [a, b]$,

$$G'(x)$$

Theorem (1): $\int_a^b f(x)dx = \left[\int f(x)dx \right]_a^b$.

Theorem (2): Let f be a continuous on $[a, b]$. If $G(x) = \int_c^x f(t)dt$, where $a \leq c \leq b$, then for every $x \in [a, b]$,

$$G'(x) = \frac{d}{dx} \left[\int_c^x f(t)dt \right]$$

Theorem (1): $\int_a^b f(x)dx = \left[\int f(x)dx \right]_a^b$.

Theorem (2): Let f be a continuous on $[a, b]$. If $G(x) = \int_c^x f(t)dt$, where $a \leq c \leq b$, then for every $x \in [a, b]$,

$$G'(x) = \frac{d}{dx} \left[\int_c^x f(t)dt \right] = f(x).$$

Theorem (1): $\int_a^b f(x)dx = \left[\int f(x)dx \right]_a^b$.

Theorem (2): Let f be a continuous on $[a, b]$. If $G(x) = \int_c^x f(t)dt$, where $a \leq c \leq b$, then for every $x \in [a, b]$,

$$G'(x) = \frac{d}{dx} \left[\int_c^x f(t)dt \right] = f(x).$$

Examples:

Theorem (1): $\int_a^b f(x)dx = \left[\int f(x)dx \right]_a^b$.

Theorem (2): Let f be a continuous on $[a, b]$. If $G(x) = \int_c^x f(t)dt$, where $a \leq c \leq b$, then for every $x \in [a, b]$,

$$G'(x) = \frac{d}{dx} \left[\int_c^x f(t)dt \right] = f(x).$$

Examples:

- (i) If $G(x) = \int_1^x \frac{1}{t} dt$ and $x > 0$, find $G'(x)$ without integrating.

Theorem (1): $\int_a^b f(x)dx = \left[\int f(x)dx \right]_a^b$.

Theorem (2): Let f be a continuous on $[a, b]$. If $G(x) = \int_c^x f(t)dt$, where $a \leq c \leq b$, then for every $x \in [a, b]$,

$$G'(x) = \frac{d}{dx} \left[\int_c^x f(t)dt \right] = f(x).$$

Examples:

- (i) If $G(x) = \int_1^x \frac{1}{t} dt$ and $x > 0$, find $G'(x)$ without integrating.
- (ii) Find $\frac{d}{dx} \int_0^x \frac{1}{1+t} dt$ without integrating.

Solution:

Solution:

(i) If $G(x) = \int_1^x \frac{1}{t} dt$ and $x > 0$, find $G'(x)$ without integrating.

Solution:

(i) If $G(x) = \int_1^x \frac{1}{t} dt$ and $x > 0$, find $G'(x)$ without integrating.

We apply theorem (2) with $c = 1$

Solution:

(i) If $G(x) = \int_1^x \frac{1}{t} dt$ and $x > 0$, find $G'(x)$ without integrating.

We apply theorem (2) with $c = 1$ and

Solution:

(i) If $G(x) = \int_1^x \frac{1}{t} dt$ and $x > 0$, find $G'(x)$ without integrating.

We apply theorem (2) with $c = 1$ and $f(x) = \frac{1}{x}$.

Solution:

(i) If $G(x) = \int_1^x \frac{1}{t} dt$ and $x > 0$, find $G'(x)$ without integrating.

We apply theorem (2) with $c = 1$ and $f(x) = \frac{1}{x}$. Choose

$$0 < a \leq c \leq b,$$

Solution:

(i) If $G(x) = \int_1^x \frac{1}{t} dt$ and $x > 0$, find $G'(x)$ without integrating.

We apply theorem (2) with $c = 1$ and $f(x) = \frac{1}{x}$. Choose $0 < a \leq c \leq b$, then f continuous on $[a, b]$.

Solution:

(i) If $G(x) = \int_1^x \frac{1}{t} dt$ and $x > 0$, find $G'(x)$ without integrating.

We apply theorem (2) with $c = 1$ and $f(x) = \frac{1}{x}$. Choose

$0 < a \leq c \leq b$, then f continuous on $[a, b]$. Hence, by theorem (2), for $x \in [a, b]$

Solution:

(i) If $G(x) = \int_1^x \frac{1}{t} dt$ and $x > 0$, find $G'(x)$ without integrating.

We apply theorem (2) with $c = 1$ and $f(x) = \frac{1}{x}$. Choose

$0 < a \leq c \leq b$, then f continuous on $[a, b]$. Hence, by theorem (2), for $x \in [a, b]$

$$G'$$

Solution:

(i) If $G(x) = \int_1^x \frac{1}{t} dt$ and $x > 0$, find $G'(x)$ without integrating. We apply theorem (2) with $c = 1$ and $f(x) = \frac{1}{x}$. Choose $0 < a \leq c \leq b$, then f continuous on $[a, b]$. Hence, by theorem (2), for $x \in [a, b]$

$$G' = \frac{d}{dx} \left[\int_c^x \frac{1}{t} dt \right]$$

Solution:

(i) If $G(x) = \int_1^x \frac{1}{t} dt$ and $x > 0$, find $G'(x)$ without integrating. We apply theorem (2) with $c = 1$ and $f(x) = \frac{1}{x}$. Choose $0 < a \leq c \leq b$, then f continuous on $[a, b]$. Hence, by theorem (2), for $x \in [a, b]$

$$G' = \frac{d}{dx} \left[\int_c^x \frac{1}{t} dt \right] = \frac{1}{x}.$$

Solution:

(i) If $G(x) = \int_1^x \frac{1}{t} dt$ and $x > 0$, find $G'(x)$ without integrating. We apply theorem (2) with $c = 1$ and $f(x) = \frac{1}{x}$. Choose $0 < a \leq c \leq b$, then f continuous on $[a, b]$. Hence, by theorem (2), for $x \in [a, b]$

$$G' = \frac{d}{dx} \left[\int_c^x \frac{1}{t} dt \right] = \frac{1}{x}.$$

(ii) Find $\frac{d}{dx} \int_0^x \frac{1}{1+t} dt$ without integrating.

Solution:

(i) If $G(x) = \int_1^x \frac{1}{t} dt$ and $x > 0$, find $G'(x)$ without integrating. We apply theorem (2) with $c = 1$ and $f(x) = \frac{1}{x}$. Choose $0 < a \leq c \leq b$, then f continuous on $[a, b]$. Hence, by theorem (2), for $x \in [a, b]$

$$G' = \frac{d}{dx} \left[\int_c^x \frac{1}{t} dt \right] = \frac{1}{x}.$$

(ii) Find $\frac{d}{dx} \int_0^x \frac{1}{1+t} dt$ without integrating.

Take $c = 0$

Solution:

(i) If $G(x) = \int_1^x \frac{1}{t} dt$ and $x > 0$, find $G'(x)$ without integrating. We apply theorem (2) with $c = 1$ and $f(x) = \frac{1}{x}$. Choose $0 < a \leq c \leq b$, then f continuous on $[a, b]$. Hence, by theorem (2), for $x \in [a, b]$

$$G' = \frac{d}{dx} \left[\int_c^x \frac{1}{t} dt \right] = \frac{1}{x}.$$

(ii) Find $\frac{d}{dx} \int_0^x \frac{1}{1+t} dt$ without integrating.
Take $c = 0$, $f(x) = \frac{1}{1+x}$

Solution:

(i) If $G(x) = \int_1^x \frac{1}{t} dt$ and $x > 0$, find $G'(x)$ without integrating. We apply theorem (2) with $c = 1$ and $f(x) = \frac{1}{x}$. Choose $0 < a \leq c \leq b$, then f continuous on $[a, b]$. Hence, by theorem (2), for $x \in [a, b]$

$$G' = \frac{d}{dx} \left[\int_c^x \frac{1}{t} dt \right] = \frac{1}{x}.$$

(ii) Find $\frac{d}{dx} \int_0^x \frac{1}{1+t} dt$ without integrating.

Take $c = 0$, $f(x) = \frac{1}{1+x}$ and $G(x) = \int_0^x \frac{1}{1+t} dt$.

Solution:

(i) If $G(x) = \int_1^x \frac{1}{t} dt$ and $x > 0$, find $G'(x)$ without integrating. We apply theorem (2) with $c = 1$ and $f(x) = \frac{1}{x}$. Choose $0 < a \leq c \leq b$, then f continuous on $[a, b]$. Hence, by theorem (2), for $x \in [a, b]$

$$G' = \frac{d}{dx} \left[\int_c^x \frac{1}{t} dt \right] = \frac{1}{x}.$$

(ii) Find $\frac{d}{dx} \int_0^x \frac{1}{1+t} dt$ without integrating.

Take $c = 0$, $f(x) = \frac{1}{1+x}$ and $G(x) = \int_0^x \frac{1}{1+t} dt$. Choose a and b such that

Solution:

(i) If $G(x) = \int_1^x \frac{1}{t} dt$ and $x > 0$, find $G'(x)$ without integrating. We apply theorem (2) with $c = 1$ and $f(x) = \frac{1}{x}$. Choose $0 < a \leq c \leq b$, then f continuous on $[a, b]$. Hence, by theorem (2), for $x \in [a, b]$

$$G' = \frac{d}{dx} \left[\int_c^x \frac{1}{t} dt \right] = \frac{1}{x}.$$

(ii) Find $\frac{d}{dx} \int_0^x \frac{1}{1+t} dt$ without integrating.

Take $c = 0$, $f(x) = \frac{1}{1+x}$ and $G(x) = \int_0^x \frac{1}{1+t} dt$. Choose a and b such that $-1 < a \leq c \leq b$,

Solution:

(i) If $G(x) = \int_1^x \frac{1}{t} dt$ and $x > 0$, find $G'(x)$ without integrating. We apply theorem (2) with $c = 1$ and $f(x) = \frac{1}{x}$. Choose $0 < a \leq c \leq b$, then f continuous on $[a, b]$. Hence, by theorem (2), for $x \in [a, b]$

$$G' = \frac{d}{dx} \left[\int_c^x \frac{1}{t} dt \right] = \frac{1}{x}.$$

(ii) Find $\frac{d}{dx} \int_0^x \frac{1}{1+t} dt$ without integrating.

Take $c = 0$, $f(x) = \frac{1}{1+x}$ and $G(x) = \int_0^x \frac{1}{1+t} dt$. Choose a and b such that $-1 < a \leq c \leq b$, so f is continuous on $[a, b]$.

Solution:

(i) If $G(x) = \int_1^x \frac{1}{t} dt$ and $x > 0$, find $G'(x)$ without integrating. We apply theorem (2) with $c = 1$ and $f(x) = \frac{1}{x}$. Choose $0 < a \leq c \leq b$, then f continuous on $[a, b]$. Hence, by theorem (2), for $x \in [a, b]$

$$G' = \frac{d}{dx} \left[\int_c^x \frac{1}{t} dt \right] = \frac{1}{x}.$$

(ii) Find $\frac{d}{dx} \int_0^x \frac{1}{1+t} dt$ without integrating.

Take $c = 0$, $f(x) = \frac{1}{1+x}$ and $G(x) = \int_0^x \frac{1}{1+t} dt$. Choose a and b such that $-1 < a \leq c \leq b$, so f is continuous on $[a, b]$. By theorem (2)

Solution:

(i) If $G(x) = \int_1^x \frac{1}{t} dt$ and $x > 0$, find $G'(x)$ without integrating. We apply theorem (2) with $c = 1$ and $f(x) = \frac{1}{x}$. Choose $0 < a \leq c \leq b$, then f continuous on $[a, b]$. Hence, by theorem (2), for $x \in [a, b]$

$$G' = \frac{d}{dx} \left[\int_c^x \frac{1}{t} dt \right] = \frac{1}{x}.$$

(ii) Find $\frac{d}{dx} \int_0^x \frac{1}{1+t} dt$ without integrating.

Take $c = 0$, $f(x) = \frac{1}{1+x}$ and $G(x) = \int_0^x \frac{1}{1+t} dt$. Choose a and b such that $-1 < a \leq c \leq b$, so f is continuous on $[a, b]$. By theorem (2)

$$G'(x)$$

Solution:

(i) If $G(x) = \int_1^x \frac{1}{t} dt$ and $x > 0$, find $G'(x)$ without integrating. We apply theorem (2) with $c = 1$ and $f(x) = \frac{1}{x}$. Choose $0 < a \leq c \leq b$, then f continuous on $[a, b]$. Hence, by theorem (2), for $x \in [a, b]$

$$G' = \frac{d}{dx} \left[\int_c^x \frac{1}{t} dt \right] = \frac{1}{x}.$$

(ii) Find $\frac{d}{dx} \int_0^x \frac{1}{1+t} dt$ without integrating.

Take $c = 0$, $f(x) = \frac{1}{1+x}$ and $G(x) = \int_0^x \frac{1}{1+t} dt$. Choose a and b such that $-1 < a \leq c \leq b$, so f is continuous on $[a, b]$. By theorem (2)

$$G'(x) = \frac{d}{dx} \left[\int_c^x \frac{1}{t+1} dt \right]$$

Solution:

(i) If $G(x) = \int_1^x \frac{1}{t} dt$ and $x > 0$, find $G'(x)$ without integrating. We apply theorem (2) with $c = 1$ and $f(x) = \frac{1}{x}$. Choose $0 < a \leq c \leq b$, then f continuous on $[a, b]$. Hence, by theorem (2), for $x \in [a, b]$

$$G' = \frac{d}{dx} \left[\int_c^x \frac{1}{t} dt \right] = \frac{1}{x}.$$

(ii) Find $\frac{d}{dx} \int_0^x \frac{1}{1+t} dt$ without integrating.

Take $c = 0$, $f(x) = \frac{1}{1+x}$ and $G(x) = \int_0^x \frac{1}{1+t} dt$. Choose a and b such that $-1 < a \leq c \leq b$, so f is continuous on $[a, b]$. By theorem (2)

$$G'(x) = \frac{d}{dx} \left[\int_c^x \frac{1}{t+1} dt \right] = \frac{1}{x+1}.$$

Example:

Example: Find $\frac{d}{dx} \left[\int_0^x \frac{1}{\sqrt{1-t^2}} dt \right]$, $|x| < 1$ without integrating.

Example: Find $\frac{d}{dx} \left[\int_0^x \frac{1}{\sqrt{1-t^2}} dt \right]$, $|x| < 1$ without integrating.

Solution:

Example: Find $\frac{d}{dx} \left[\int_0^x \frac{1}{\sqrt{1-t^2}} dt \right]$, $|x| < 1$ without integrating.

Solution:

Take $c = 0$,

Example: Find $\frac{d}{dx} \left[\int_0^x \frac{1}{\sqrt{1-t^2}} dt \right]$, $|x| < 1$ without integrating.

Solution:

Take $c = 0$, $f(x) = \frac{1}{\sqrt{1-x^2}}$

Example: Find $\frac{d}{dx} \left[\int_0^x \frac{1}{\sqrt{1-t^2}} dt \right]$, $|x| < 1$ without integrating.

Solution:

Take $c = 0$, $f(x) = \frac{1}{\sqrt{1-x^2}}$ and $G(x) = \int_0^x \frac{1}{\sqrt{1-t^2}} dt$.

Example: Find $\frac{d}{dx} \left[\int_0^x \frac{1}{\sqrt{1-t^2}} dt \right]$, $|x| < 1$ without integrating.

Solution:

Take $c = 0$, $f(x) = \frac{1}{\sqrt{1-x^2}}$ and $G(x) = \int_0^x \frac{1}{\sqrt{1-t^2}} dt$. Note that $f(x)$ is not continuous at $x = \pm 1$.

Example: Find $\frac{d}{dx} \left[\int_0^x \frac{1}{\sqrt{1-t^2}} dt \right]$, $|x| < 1$ without integrating.

Solution:

Take $c = 0$, $f(x) = \frac{1}{\sqrt{1-x^2}}$ and $G(x) = \int_0^x \frac{1}{\sqrt{1-t^2}} dt$. Note that $f(x)$ is not continuous at $x = \pm 1$. Choose $-1 < a \leq c \leq b < 1$, so $f(x)$ is continuous on $[a, b]$.

Example: Find $\frac{d}{dx} \left[\int_0^x \frac{1}{\sqrt{1-t^2}} dt \right]$, $|x| < 1$ without integrating.

Solution:

Take $c = 0$, $f(x) = \frac{1}{\sqrt{1-x^2}}$ and $G(x) = \int_0^x \frac{1}{\sqrt{1-t^2}} dt$. Note that $f(x)$ is not continuous at $x = \pm 1$. Choose $-1 < a \leq c \leq b < 1$, so $f(x)$ is continuous on $[a, b]$. Now by theorem (2), we get

Example: Find $\frac{d}{dx} \left[\int_0^x \frac{1}{\sqrt{1-t^2}} dt \right]$, $|x| < 1$ without integrating.

Solution:

Take $c = 0$, $f(x) = \frac{1}{\sqrt{1-x^2}}$ and $G(x) = \int_0^x \frac{1}{\sqrt{1-t^2}} dt$. Note that $f(x)$ is not continuous at $x = \pm 1$. Choose $-1 < a \leq c \leq b < 1$, so $f(x)$ is continuous on $[a, b]$. Now by theorem (2), we get

$$G'(x)$$

Example: Find $\frac{d}{dx} \left[\int_0^x \frac{1}{\sqrt{1-t^2}} dt \right]$, $|x| < 1$ without integrating.

Solution:

Take $c = 0$, $f(x) = \frac{1}{\sqrt{1-x^2}}$ and $G(x) = \int_0^x \frac{1}{\sqrt{1-t^2}} dt$. Note that

$f(x)$ is not continuous at $x = \pm 1$. Choose

$-1 < a \leq c \leq b < 1$, so $f(x)$ is continuous on $[a, b]$. Now by theorem (2), we get

$$G'(x) = \frac{d}{dx} \left[\int_0^x \frac{1}{\sqrt{1-t^2}} dt \right]$$

Example: Find $\frac{d}{dx} \left[\int_0^x \frac{1}{\sqrt{1-t^2}} dt \right]$, $|x| < 1$ without integrating.

Solution:

Take $c = 0$, $f(x) = \frac{1}{\sqrt{1-x^2}}$ and $G(x) = \int_0^x \frac{1}{\sqrt{1-t^2}} dt$. Note that

$f(x)$ is not continuous at $x = \pm 1$. Choose

$-1 < a \leq c \leq b < 1$, so $f(x)$ is continuous on $[a, b]$. Now by theorem (2), we get

$$G'(x) = \frac{d}{dx} \left[\int_0^x \frac{1}{\sqrt{1-t^2}} dt \right] = \frac{1}{\sqrt{1-x^2}}.$$

Exercise:

Example: Find $\frac{d}{dx} \left[\int_0^x \frac{1}{\sqrt{1-t^2}} dt \right]$, $|x| < 1$ without integrating.

Solution:

Take $c = 0$, $f(x) = \frac{1}{\sqrt{1-x^2}}$ and $G(x) = \int_0^x \frac{1}{\sqrt{1-t^2}} dt$. Note that $f(x)$ is not continuous at $x = \pm 1$. Choose

$-1 < a \leq c \leq b < 1$, so $f(x)$ is continuous on $[a, b]$. Now by theorem (2), we get

$$G'(x) = \frac{d}{dx} \left[\int_0^x \frac{1}{\sqrt{1-t^2}} dt \right] = \frac{1}{\sqrt{1-x^2}}.$$

Exercise: Find

$$\frac{d}{dx} \left[\int_0^1 \frac{x^5 \sqrt{x^4 + 3}}{\sqrt{x^{10} + x^6 + 2}} dx \right].$$

without integrating.

Theorem: If g and k are differentiable and f is a continuous function,

Theorem: If g and k are differentiable and f is a continuous function, then

Theorem: If g and k are differentiable and f is a continuous function, then

$$\frac{d}{dx} \int_{k(x)}^{g(x)} f(t) dt$$

Theorem: If g and k are differentiable and f is a continuous function, then

$$\frac{d}{dx} \int_{k(x)}^{g(x)} f(t) dt = f(g(x))$$

Theorem: If g and k are differentiable and f is a continuous function, then

$$\frac{d}{dx} \int_{k(x)}^{g(x)} f(t) dt = f(g(x))g'(x)$$

Theorem: If g and k are differentiable and f is a continuous function, then

$$\frac{d}{dx} \int_{k(x)}^{g(x)} f(t) dt = f(g(x))g'(x) - f(k(x))$$

Theorem: If g and k are differentiable and f is a continuous function, then

$$\frac{d}{dx} \int_{k(x)}^{g(x)} f(t) dt = f(g(x))g'(x) - f(k(x))k'(x).$$

Theorem: If g and k are differentiable and f is a continuous function, then

$$\frac{d}{dx} \int_{k(x)}^{g(x)} f(t) dt = f(g(x))g'(x) - f(k(x))k'(x).$$

Example: Find $\frac{d}{dx} \int_{3x}^{x^2} (t^3 + 1)^{10} dt$, without integrating.

Example: Find $\frac{d}{dx} \int_{3x}^{x^2} (t^3 + 1)^{10} dt$, without integrating.

Solution:

Example: Find $\frac{d}{dx} \int_{3x}^{x^2} (t^3 + 1)^{10} dt$, without integrating.

Solution: Take $k(x) = 3x$

Example: Find $\frac{d}{dx} \int_{3x}^{x^2} (t^3 + 1)^{10} dt$, without integrating.

Solution: Take $k(x) = 3x$, $g(x) = x^2$

Example: Find $\frac{d}{dx} \int_{3x}^{x^2} (t^3 + 1)^{10} dt$, without integrating.

Solution: Take $k(x) = 3x$, $g(x) = x^2$ and $f(x) = (x^3 + 1)^{10}$.

Example: Find $\frac{d}{dx} \int_{3x}^{x^2} (t^3 + 1)^{10} dt$, without integrating.

Solution: Take $k(x) = 3x$, $g(x) = x^2$ and $f(x) = (x^3 + 1)^{10}$.
The function k and g are differentiable

Example: Find $\frac{d}{dx} \int_{3x}^{x^2} (t^3 + 1)^{10} dt$, without integrating.

Solution: Take $k(x) = 3x$, $g(x) = x^2$ and $f(x) = (x^3 + 1)^{10}$.
The function k and g are differentiable and f is continuous

Example: Find $\frac{d}{dx} \int_{3x}^{x^2} (t^3 + 1)^{10} dt$, without integrating.

Solution: Take $k(x) = 3x$, $g(x) = x^2$ and $f(x) = (x^3 + 1)^{10}$.
The function k and g are differentiable and f is continuous,
therefore, by the previous theorem we get

Example: Find $\frac{d}{dx} \int_{3x}^{x^2} (t^3 + 1)^{10} dt$, without integrating.

Solution: Take $k(x) = 3x$, $g(x) = x^2$ and $f(x) = (x^3 + 1)^{10}$.
The function k and g are differentiable and f is continuous,
therefore, by the previous theorem we get

$$f(g(x))g'(x)$$

Example: Find $\frac{d}{dx} \int_{3x}^{x^2} (t^3 + 1)^{10} dt$, without integrating.

Solution: Take $k(x) = 3x$, $g(x) = x^2$ and $f(x) = (x^3 + 1)^{10}$.
The function k and g are differentiable and f is continuous,
therefore, by the previous theorem we get

$$f(g(x))g'(x) = f(x^2)2x$$

Example: Find $\frac{d}{dx} \int_{3x}^{x^2} (t^3 + 1)^{10} dt$, without integrating.

Solution: Take $k(x) = 3x$, $g(x) = x^2$ and $f(x) = (x^3 + 1)^{10}$.
The function k and g are differentiable and f is continuous,
therefore, by the previous theorem we get

$$f(g(x))g'(x) = f(x^2)2x = 2x((x^2)^3 + 1)^{10}$$

Example: Find $\frac{d}{dx} \int_{3x}^{x^2} (t^3 + 1)^{10} dt$, without integrating.

Solution: Take $k(x) = 3x$, $g(x) = x^2$ and $f(x) = (x^3 + 1)^{10}$.
The function k and g are differentiable and f is continuous,
therefore, by the previous theorem we get

$$f(g(x))g'(x) = f(x^2)2x = 2x((x^2)^3 + 1)^{10}$$
$$f(k(x))k'(x)$$

Example: Find $\frac{d}{dx} \int_{3x}^{x^2} (t^3 + 1)^{10} dt$, without integrating.

Solution: Take $k(x) = 3x$, $g(x) = x^2$ and $f(x) = (x^3 + 1)^{10}$.
The function k and g are differentiable and f is continuous,
therefore, by the previous theorem we get

$$f(g(x))g'(x) = f(x^2)2x = 2x((x^2)^3 + 1)^{10}$$

$$f(k(x))k'(x) = f(3x).3$$

Example: Find $\frac{d}{dx} \int_{3x}^{x^2} (t^3 + 1)^{10} dt$, without integrating.

Solution: Take $k(x) = 3x$, $g(x) = x^2$ and $f(x) = (x^3 + 1)^{10}$.
The function k and g are differentiable and f is continuous,
therefore, by the previous theorem we get

$$f(g(x))g'(x) = f(x^2)2x = 2x((x^2)^3 + 1)^{10}$$

$$f(k(x))k'(x) = f(3x).3 = 3(3x^3 + 1)^{10} = 3(27x^3 + 1)^{10}.$$

Example: Find $\frac{d}{dx} \int_{3x}^{x^2} (t^3 + 1)^{10} dt$, without integrating.

Solution: Take $k(x) = 3x$, $g(x) = x^2$ and $f(x) = (x^3 + 1)^{10}$.
The function k and g are differentiable and f is continuous,
therefore, by the previous theorem we get

$$f(g(x))g'(x) = f(x^2)2x = 2x((x^2)^3 + 1)^{10}$$

$$f(k(x))k'(x) = f(3x).3 = 3(3x^3 + 1)^{10} = 3(27x^3 + 1)^{10}.$$

Hence,

Example: Find $\frac{d}{dx} \int_{3x}^{x^2} (t^3 + 1)^{10} dt$, without integrating.

Solution: Take $k(x) = 3x$, $g(x) = x^2$ and $f(x) = (x^3 + 1)^{10}$. The function k and g are differentiable and f is continuous, **therefore**, by the previous theorem we get

$$f(g(x))g'(x) = f(x^2)2x = 2x((x^2)^3 + 1)^{10}$$

$$f(k(x))k'(x) = f(3x).3 = 3(3x^3 + 1)^{10} = 3(27x^3 + 1)^{10}.$$

Hence,

$$\frac{d}{dx} \left[\int_{3x}^{x^2} (t^3 + 1)^{10} dt \right]$$

Example: Find $\frac{d}{dx} \int_{3x}^{x^2} (t^3 + 1)^{10} dt$, without integrating.

Solution: Take $k(x) = 3x$, $g(x) = x^2$ and $f(x) = (x^3 + 1)^{10}$.
The function k and g are differentiable and f is continuous,
therefore, by the previous theorem we get

$$\begin{aligned} f(g(x))g'(x) &= f(x^2)2x = 2x((x^2)^3 + 1)^{10} \\ f(k(x))k'(x) &= f(3x).3 = 3(3x^3 + 1)^{10} = 3(27x^3 + 1)^{10}. \end{aligned}$$

Hence,

$$\frac{d}{dx} \left[\int_{3x}^{x^2} (t^3 + 1)^{10} dt \right] = f(g(x))g'(x) - f(k(x))k'(x)$$

Example: Find $\frac{d}{dx} \int_{3x}^{x^2} (t^3 + 1)^{10} dt$, without integrating.

Solution: Take $k(x) = 3x$, $g(x) = x^2$ and $f(x) = (x^3 + 1)^{10}$. The function k and g are differentiable and f is continuous, **therefore**, by the previous theorem we get

$$f(g(x))g'(x) = f(x^2)2x = 2x((x^2)^3 + 1)^{10}$$

$$f(k(x))k'(x) = f(3x).3 = 3(3x^3 + 1)^{10} = 3(27x^3 + 1)^{10}.$$

Hence,

$$\begin{aligned} \frac{d}{dx} \left[\int_{3x}^{x^2} (t^3 + 1)^{10} dt \right] &= f(g(x))g'(x) - f(k(x))k'(x) \\ &= 2x((x^2)^3 + 1)^{10} - 3(27x^3 + 1)^{10}. \end{aligned}$$

Exercises:

(a) Find

$$\frac{d}{dx} \int_0^{x^2} \sqrt[3]{t^4 + 1} dt.$$

Exercises:

(a) Find

$$\frac{d}{dx} \int_0^{x^2} \sqrt[3]{t^4 + 1} dt.$$

(b) Find

$$\frac{d}{dx} \int_{6x-1}^0 \sqrt{4t + 9} dt.$$

Exercises:

(a) Find

$$\frac{d}{dx} \int_0^{x^2} \sqrt[3]{t^4 + 1} dt.$$

(b) Find

$$\frac{d}{dx} \int_{6x-1}^0 \sqrt{4t + 9} dt.$$

(c) If $F(x) = (x^2) \int_2^x [t + 3F'(t)] dt$, find $F'(2)$.

Exercises:

(a) Find

$$\frac{d}{dx} \int_0^{x^2} \sqrt[3]{t^4 + 1} dt.$$

(b) Find

$$\frac{d}{dx} \int_{6x-1}^0 \sqrt{4t + 9} dt.$$

(c) If $F(x) = (x^2) \int_2^x [t + 3F'(t)] dt$, find $F'(2)$.

(d) If

$$g(x) = \frac{\int_{\frac{\pi}{2}}^x \sin(t) dt}{x},$$

Exercises:

(a) Find

$$\frac{d}{dx} \int_0^{x^2} \sqrt[3]{t^4 + 1} dt.$$

(b) Find

$$\frac{d}{dx} \int_{6x-1}^0 \sqrt{4t + 9} dt.$$

(c) If $F(x) = (x^2) \int_2^x [t + 3F'(t)] dt$, find $F'(2)$.

(d) If

$$g(x) = \frac{\int_{\frac{\pi}{2}}^x \sin(t) dt}{x},$$

find $g'(x)$.