

Accepted Manuscript

Original article

Isolation of Antimicrobial Peptides from *Apis florea* and *Apis carnica* in Saudi Arabia and Investigation of the Antimicrobial Properties of Natural Honey Samples

Tahany H. Ayaad, Ghada H. Shaker, Amal M. Almuhaa

PII: S1018-3647(11)00003-6
DOI: [10.1016/j.jksus.2011.01.002](https://doi.org/10.1016/j.jksus.2011.01.002)
Reference: JKUSUS 142

To appear in: *Journal of King Saud University - Science*

Received Date: 6 November 2010
Accepted Date: 16 January 2011

Please cite this article as: T.H. Ayaad, G.H. Shaker, A.M. Almuhaa, Isolation of Antimicrobial Peptides from *Apis florea* and *Apis carnica* in Saudi Arabia and Investigation of the Antimicrobial Properties of Natural Honey Samples, *Journal of King Saud University - Science* (2011), doi: [10.1016/j.jksus.2011.01.002](https://doi.org/10.1016/j.jksus.2011.01.002)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Isolation of Antimicrobial Peptides from *Apis florae* and *Apis carnica* in Saudi Arabia and Investigation of the Antimicrobial Properties of Natural Honey Samples

Tahany, H. Ayaad; *^aGhada, H. Shaker; ^bAmal, M. Almuhnaa^c

^{a,c}Department of Zoology, Faculty of Science, King Saud University.

^bDepartment of Pharmaceutics - Microbiology, Faculty of Pharmacy, King Saud University.
P.O.Box11495, Riyadh11452, Saudi Arabia. E-mail: ayaad_t@hotmail.com

ACCEPTED MANUSCRIPT

Isolation of Antimicrobial Peptides from *Apis floreae* and *Apis carnica* in Saudi Arabia and Investigation of the Antimicrobial Properties of Natural Honey Samples

Abstract:

As part of the ongoing search for novel antimicrobial agents and their use in singular or combined drug therapy, peptide fractions of molecular weights about 14.500 and 15.00 KDa were isolated from the hemolymph of wild (*Apis floreae*) and carniolan (*Apis carnica*) bees of Saudi Arabia obtained from different regions in Riyadh (variable plant sources) during the honey seasons (spring and summer) 2008-2009. Following experimental infection with 1.1×10^6 viable *Escherichia coli* cells (ATCC 25922), the antimicrobial peptides were purified to homogeneity by Reversed - Phase High Performance Liquid Chromatography. Antibacterial activity of the isolated peptide was evaluated *in vitro* by an agar well diffusion method for *Escherichia coli* strain (ATCC 25922) and *Klebsiella pneumoniae* strain (ATCC 11678) the major Gram negative pathogens causing urinary tract infections, and *Staphylococcus aureus* (ATCC 6538) as Gram positive bacteria. A total of ten honey samples collected from bee hives selected arbitrary at different floral areas south Riyadh were also investigated for their antimicrobial activity against the yeast, *Candida albicans* (ATCC 10231) and four standards bacteria strains, *Escherichia coli* (ATCC 25922), *Pseudomonas aeruginosa* (ATCC 27853), *Staphylococcus aureus* (ATCC 6538) and *Bacillus subtilis* (ATCC 6633) using standard antimicrobial assays. The isolated antibacterial peptides and the different honey samples revealed comparable marked variations in antimicrobial activities and their sensitivity might be depending on their variable floral sources.

Key Words: Antimicrobial peptides, Honey, *Apis floreae*, *Apis carnica* , Antimicrobial properties.

Introduction:

Antibiotic-resistant bacteria continue to be of major health concern world-wide. Since the use of antibiotics became widespread over 50 years ago, bacteria have progressively developed resistance (Hsueh *et al.*, 2005). Consequently, scientific efforts have been made to study and develop new compounds to be used beyond conventional antibiotic therapy. Honey has been used since ancient times for the treatment of some respiratory diseases and for the healing of skin wounds. It has been proposed that the healing effect of honey could be due to various physical and chemical properties (Snow and Manley-Harris, 2004). Along with the rapidly increasing interest and research into natural health remedies and supplements, is a resurgence of interest in the therapeutic use of honey. Honey as most natural products, may have a large variance in therapeutic components depending on its origin. Thus, the floral source of honey plays an important role on its biological properties (Molan, 2002). In consequence, it would not be surprising that the provenance of honey could determine its antibacterial properties. Honey produced by honeybees (*Apis floreae*) is one of the oldest traditional medicines considered to be important in the treatment of respiratory, gastrointestinal infection and various other diseases due to the absence of sufficient modern health care system, particularly in rural areas.

Often during pathogenic invasion, the first line of defense involves the innate mechanisms of immunity which in turn is followed by acquired immune responses involving the activation of T and B cells against specific antigens (Fearon and Locksley, 1996; Medzhitov and Janeway, 2000). In contrast to these acquired immune mechanisms, endogenous peptides, which are constitutively expressed or induced, provide a fast and effective means of defense against the pathogen. This group of molecules termed 'antimicrobial peptides' (AMPs) constitutes a primitive immune defense mechanism and is found in a wide range of eukaryotic organisms, from humans, plants and insects (Lehrer and Ganz, 1999). AMPs are an important component of the natural defenses of most living organisms against invading pathogens. During the past two decades several AMPs have been isolated from a wide variety of animals, both vertebrates and invertebrates, and plants as well as from bacteria and fungi. These peptides exhibit broad-spectrum activity against a wide range of microorganisms including Gram-positive and Gram-negative bacteria, protozoa, yeast, fungi and viruses, they have potential to overcome bacterial

resistance makes them promising candidates for therapeutic drugs (Bals, 2000). AMPs are classified based on the three dimensional structural studies carried out with the help of NMR. Most of these peptides are believed to act by disrupting the plasma membrane leading to the lysis of the cell. AMPs have been found to be excellent candidates for developing novel antimicrobial agents and few of these peptides show antimicrobial activity against pathogens causing sexually transmitted infection. Few peptides have already entered clinical trials for the treatment of impetigo, diabetic foot ulcers and gastric helicobacter infections (Reddy *et al.*, 2004). One of the most promising among these antimicrobial peptide families are the cell-free immune repertoire of honeybees (*Apis mellifera*) that are induced by bacterial infection provide broad-spectrum antibacterial defense, such as apidaecin, hymenoptaecin, abaecin, and bee defensin. These peptides represent a viable treatment option for the major pathogens in urinary tract infections, that is, *E. coli* and *K. pneumoniae*, causing 90–95% of all urinary tract infections (Czihal *et al.*, 2007).

The purpose of the present study aimed at the isolation of antimicrobial peptides of wild Saudi Arabian *Apis florae* and carniolan *Apis carnica*. Evaluating scientifically the *in vitro* antimicrobial potential of these peptides and investigating the properties of ten natural honey samples produced by honeybees (*Apis florae*) against standard microorganisms among those commonly involved in causing diseases.

Materials and methods:

Insects:

Two groups of adult honey bee workers (age: 18 – 45 days and weight: 0.25gm ± 0.01) were either collected from natural environmental locations with different floral origins south Riyadh (*Apis florae*) or obtained from Department of Agricultural Extension and Rearing Parcels of Bee Queens in Saudi Arabia (*Apis carnica*). Adult bees were kept in small cages in the laboratory until used for induction by bacteria and isolation of peptides.

Microorganisms:

The standard microorganisms used in this study were the yeast *Candida albicans* (ATCC 10231) and five different bacteria strains, *Escherichia coli* (ATCC 25922), *Pseudomonas aeruginosa* (ATCC 27853), *Klebsiella pneumoniae* (ATCC 11678) as Gram negative and *Staphylococcus aureus* (ATCC 6538), *Bacillus subtilis* (ATCC 6633) as Gram positive bacteria.

Immune induction of honeybees:

Humoral immunity was induced in adult honeybees by puncturing a leg with the tip of a hypodermic needle dipped in the 1.1×10^6 viable *E. coli* cells (ATCC 25922), equivalent to 0.5 McFarland tube suspended in 1 μ l phosphate-buffered saline (PBS, 0.15 M, pH 7.2). Intact and induced adult bees were bled by decapitation and oozing lymph samples (2-4 μ l / bee) were taken after 24hr. post induction, the collected hemolymph was pooled in tubes containing 100 μ l of 2% trifluoroacetic acid (TFA) to prevent proteolytic degradation of the immunoinduced peptides and to precipitate proteins as described by Casteels *et al.*(1994) (for peptide purification). The collected hemolymph was also pooled in ice-cooled Eppendorf tubes containing few crystals of phenylthiourea to prevent melanization of samples (SDS– PAGE of crude hemolymph samples). Hemocytes were centrifuged (10000 g for 10 min.) and the lymph was collected and freeze stored at -70 °C till used.

Purification of antibacterial peptides: (Reversed Phase- High Performance Liquid Chromatography) RP-HPLC.

The lymph samples were heat-treated (100°C/5 min.). The precipitate was spun down and the clear supernatant was acidified with an equal volume of 0.1% TFA, and fractionated by several rounds of high performance liquid chromatography using reversed-phase column supports, all as described by Casteels *et al.*(1993) and Lauth *et al.* (1998).

Sample (50 μ l) aliquots of diluted lymph were taken for RP-HPLC analysis using an ABI 150A system (Applied Biosystems Inc., Ramsey, NJ) with a VYDAC C4 (214 TP54) analytical column (The separations group, Hesperia, CA). Solvent A was 0.1 % TFA (pH 2.0) and solvent B: 70% acetonitrile (MeCN). Fractions were eluted

at 1 ml/min. (70 min. total times). UV detection was done at 214 nm. All differential peaks between control and immune lymph, including peaks 1 and 2, were collected and further purified on VYDAC C18 (218TP54). Collected fractions were lyophilized and re-dissolved in Milli Q water, Promega (nuclease free water) before being tested for biological activity against standard microorganisms, *E. coli* strain (ATCC 25922) and *K. pneumoniae* strain (ATCC 11678). Following this procedure 1.0- 10 µg peptides were routinely purified from each separate batch of *Apis floreae* and *Apis carnica* worker bees.

Sodium Dodecyle Sulphate Polyacrylamide Gel Electrophoresis (SDS- PAGE) for the isolated *Apis floreae* and *Apis carnica* peptides:

SDS- PAGE of control and purified antimicrobial peptides was carried out by the discontinuous buffer system as described by Laemmli, (1970) with some modifications. Electrophoresis was carried out at a constant voltage of 200 V for 90 min. and 12% polyacrylamide gel, under denaturing conditions. The gels were calibrated with standard molecular weight proteins (high and low ranges: 200, 97.4, 68, 29, 18.4 and 8.15 kDa). Protein bands were visualized by Commassie Brilliant Blue dye staining (CBB). M.Wt calculations were determined by regression analysis using the manufacturers' soft-ware.

All Chemicals and buffers are purchased from (Sigma, Chemicals, Co.– Aldrich, Chemicals ,Co., Sweden).

Honey samples:

This study was carried out on ten natural honey samples (1 kg each) collected in Saudi Arabia during the honey seasons (spring and summer) 2008 - 2009. Honey samples collected from *Apis floreae* bee hives were marked randomly as indicated in table (1). Each honey sample was collected in a sterile universal glass container and kept at 2–8°C until tested. Each sample was tested at original concentration 100% and diluted to 70% and 30% of its original concentration using physiological saline PBS, pH 7.2 according to the method described by Nzeako and Hamdi, (2000) with some modifications.

Antimicrobial activity of honey samples and purified antimicrobial peptides:

An agar well diffusion method was used to assess the antimicrobial activity of the honey samples and purified peptide fractions against certain selected standard microorganisms (NCCLS, 2003). Fifty microlitres (50 μ l) of each honey dilution (undiluted, 70% and 30% w/v) were used against *E. coli* (ATCC 25922), *P. aeruginosa* (ATCC 27853), *S. aureus* (ATCC 6538), *B. subtilis* (ATCC 6633) and *C. albicans* (ATCC 10231), separately. While 20 μ l of two fold serial dilution of purified peptide fractions were tested against each of *E. coli* (ATCC 25922), *K. pneumoniae* (ATCC 11678) the major Gram negative pathogens causing urinary tract infections and *S. aureus* (ATCC 6538) as G-positive bacteria, separately. Each of the bacterial strains was inoculated into nutrient broth and incubated overnight at 37°C until growth was 1.5 (optical density at 640 nm). Each of the honey sample concentrations and the purified peptide fractions were added, separately into wells of 5 millimeters (mm) diameter of inoculated Mueller- Hinton agar plates (Oxoid) by selected microorganisms, each dilution was done in triplicate. The plates were incubated at 37°C till the honey seeped into the agar. Zones of growth inhibition were recorded in mm after an overnight incubation at 37°C. The end point of antimicrobial activity of each determination was defined as the highest dilution (lowest concentration) producing an inhibition zone with the tested organisms. The growth after 24 hr. incubation at 37°C was then compared to a control plate that contained no peptide fraction or honey samples. All strains were handled under aseptic conditions and the microorganisms were destroyed by autoclave to ensure bio-safety.

Statistical Analysis:

Data analysis were carried out using SPSS for Windows Ver. 17 .0.

Results:

Major antimicrobial peptides were isolated from two groups of Saudi Arabian *Apis florae* and carniolan bees either unchallenged (intact) or induced. The antimicrobial peptides were initially fractionated under RP- HPLC by using 70% v/v acetonitrile containing 0.1%v/v trifluoroacetic acid recovered as two peaks 1 and 2 (Fig. 1) in the final RP-HPLC step. Gel electrophoresis analysis indicated apparent homogeneity and approximate M.Wt of 14.500 and 15.00 KDa for the isolated peaks 1 and 2 respectively. Differential pattern analysis of described peaks are barely detectable in unchallenged (intact) bees but are strongly induced upon induction. All the peptide fractions obtained from chromatography columns isolated from *Apis florae* and *Apis carnica* groups either (intact or bacterial induced) presented antimicrobial activity against two Gram negative bacteria *E. coli* and *K. pneumoniae*. Fig. (2) represents antibacterial activity of the purified antimicrobial peptide fractions 14.500 and 15.00 KDa for each type of wild and carniolan bees either (induced or intact) against the tested G-negative bacteria. Presented data revealed that the isolated peptides from intact and induced honey bees have significant moderate activities against *E. coli* and *K. pneumoniae*. Overall, the activity of the purified peptide showed comparable antimicrobial activity in both groups of honey bees either *Apis florae* or *Apis carnica*.

On the other hand, no antimicrobial activity was observed against *S. aureus* in case of intact fractions while the isolated antimicrobial peptide from immune bees have significant activities as shown in Fig.(3).

Table (1) Examined natural honey samples and their floral sources

No. of honey sample	Season	Location	Plant cover
1	Summer	Alsomman (plain)	<i>Acacia spp.</i> (Mimosaceae), <i>Ziziphus spina</i> (Rhamnaceae), <i>Rhanterium epapposum</i> (Asteraceae)
2	Summer	Hail (plain)	<i>Acacia spp.</i> (Mimosaceae)
3	Summer	Al-Qasim (plain)	<i>Ziziphus spina</i> (Rhamnaceae)
4	Summer	Horimalaa	<i>Acacia spp.</i> (Mimosaceae), <i>Ziziphus spina</i> (Rhamnaceae)
5	Spring	Roda Eltanhat (plain)	<i>Ziziphus spina</i> (Rhamnaceae), <i>Acacia spp.</i> (Mimosaceae), <i>Peganum harmala</i> (Zygophyllaceae), <i>Rhanterium epapposum</i> (Asteraceae) Wild Shafalah (Mimosaceae)
6	Summer	Horimalaa	<i>Acacia spp.</i> (Mimosaceae)
7	Spring	Toik Mountain	Wild Shaflah (Mimosaceae)
8	Spring	Elkharj	<i>Trifolium spp.</i>
9	Summer	Roda Elkhashm (plain)	<i>Ziziphus spina</i> (Rhamnaceae), <i>Acacia spp.</i> (Mimosaceae)
10	Summer	Horimalaa	<i>Ziziphus spina</i> (Rhamnaceae), <i>Acacia spp.</i> (Mimosaceae)

Flora sources are classified according to Migahid (1996) and Chaudhary (2000)




Figure 1 RP-HPLC purification and SDS-PAGE of antimicrobial peptide fractions from wild *Apis florea* and carniolan *Apis carnica* bees. Heat treated lymph was fractionated, separately, on a VYDAC® C4 and finally C18 VYDAC® columns,. Fractions 1 and 2 developed at a flow rate of 1 ml/min and a gradient of 30-70 % B/70 min. Panels A and B show the HPLC patterns of the isolated peptide fractions from, intact (non-induced) and *E. coli* induced *Apis florea* bees. Panels C and D intact and induced *Apis carnica* bees. The dashed line represents the percentage of solvent B (70% acetonitrile) in solvent A (0.1 % trifluoroacetic acid). Electrophoretic analysis of pure peptide fractions (lanes a and b) and crude adult bee lymph (lane c) were shown in the center of the figure. The molecular weight of standards is shown in lane d ($\times 10^3$). CBB stained.




Figure 2 Antibacterial activity of *Apis floreae* and *Apis carnica* purified peptide fractions against (A) *E.coli* and (B) *K.pneumoniae*. Comparable inhibition zones observed by *Apis floreae* and *Apis carnica* bee peptides whether intact (1-4) or induced (5-8 wells) against both gram negative bacteria.





Figure 3 Antibacterial activity of *Apis floreae* and *Apis carnica* bee purified peptide fractions against *S.aureus*. No activity detected in fractions (1- 4) from intact *Apis floreae* and *Apis carnica* bees. Labels (5, 6 and 7, 8) show inhibition zones of purified fractions from both induced bee types.


Antimicrobial activity of honey samples:

Antimicrobial activity of the ten honey samples were done with three concentrations 30% w/v, 70% w/v and undiluted used in this study are shown in Figure (4). By using the agar-well diffusion method, the growth of all five standard microorganisms *S. aureus*, *B. subtilis*, *P. aeruginosa*, *E. coli* and *C. albicans* were inhibited (Jorgensen and Ferraro, 1998). Antimicrobial activity was evaluated by measuring the zone of inhibition (mm) against the tested microorganisms. At higher concentrations of honey there was a progressive increase in growth inhibition of the microorganisms. It was observed that *S. aureus* was the most inhibited bacterial strain by all honey samples. The average diameter of the inhibition zones produced by the undiluted honeys samples was approximately (35 ± 0.1) mm. Our data show that all honey samples tested have some antibacterial action at 30%, 70% and undiluted concentrations. In general, all the five tested microorganisms were variably sensitive to honey up to 30% concentration.

A



B



C




Figure 4 Mean of microbial inhibition zones by different concentrations (mm) of ten honey samples (30%, 70% and undiluted A,B, C, respectively) against five standard microorganisms (B refers to *Bacillus subtilis*, C, refers to *Candida albicans*, E refers to *Escherichia coli*, P refers to *Pseudomonas aeruginosa* and S refers to *Staphylococcus aureus*).

DISCUSSION:

In recent years there was a dramatically increase in bacteria strains resistance to one or even several antibiotics. Thus, the development of antimicrobial compounds with novel modes of action is a major focus of current pharmaceutical research. A very interesting and promising approach relies on antibacterial peptides, because bacteria do not develop any resistance to these antimicrobial peptide families. One of the most promising among these families are the short, proline-rich antibacterial peptides originally isolated from insects, such as apidaecin, drosocin, formaecin, and pyrrhocoricin. These peptides represent a viable treatment option for the major pathogens in urinary tract infections, that is, *E. coli* and *K. pneumoniae*, causing 90–95% of all urinary tract infections (Czihal *et al.*, 2007).

The exact mechanism of action of AMPs remains a matter of controversy, there is a consensus that these peptides selectively disrupt the cell membranes and the amphipathic structural arrangement of the peptides is believed to play an important role in this mechanism. The phospholipids head group charge on cell membranes and peptide charge distribution appears to play an important role in the peptide membrane interactions (Oren and Shai, 1998; Cudic and Otvos, 2002). There is accumulating

evidence suggesting that the antibacterial or self defense peptides which are usually highly basic, recognize the acidic phospholipids exposed on the surface of the bacterial membrane (Tytler *et al.*, 1995). In the case of microbes, the anionic lipids are present on the outer surface of the membrane whereas for mammalian cells, anionic lipids are present along the cytoplasmic side of the membrane. This feature might account for their preferential activity against bacteria but not against mammalian cells.

Several structure function studies on AMPs have been published (Hanke and Schlue, 1997; Wieprecht *et al.*, 1997; Mor, 2000). It is well documented that biophysical properties such as secondary structure, overall charge and hydrophobicity influence the interaction of AMPs with model membranes and biological cells

An ubiquitous polypeptide fractions were purified from the Saudi Arabian honeybee *Apis florea* and *Apis carnica* . The isolated polypeptides are naturally detected in the adult bees hemolymph, presenting moderate spectrum of antimicrobial activity against *E. coli* and *K.pneumoniae* bacteria the major Gram negative pathogens causing urinary tract infections to human, this is may be due to the nature of G-negative cell wall which characterized by thin layer of peptidoglycans the main target of antimicrobial peptide activity.

On the other hand isolated antimicrobial peptides showed significant inhibition to *S. aureus* growth with induced fractions and non significant with intact one as shown in Fig .3. These results can be explained as G-positive bacteria have thick layer of peptidoglycans in its cell wall which required immune induction for honeybees to produce more peptides.

Apparently an immune induction of the bees increased the polypeptides production as appeared from the purification peaks (RP-HPLC, Fig. 1) and more inhibition of the growth of G-positive bacteria (Fig. 3) during evaluating their antimicrobial activity. .

Similar results was obtained by Casteels *et al.*(1993) for the apidaecin as an increase in the transcription level occurred 4 hr. after experimental infection and very high concentrations were sustained throughout the entire 36 hr. post infection. This suggests that there is mounting evidence that activation of insect peptide antibiotic gene is the endpoint of a signal pathway that has bacteria, or more specifically lipopolysaccharide (LPS), as initiating agent (Girardin *et al.*,2002).

Comparison of the mean inhibition zones of antibacterial activity between the two purified fractions 1 and 2 revealed non significant differences $P \leq 0.05$ indicating that these peptide fractions are functionally identical. These results are also indicative, as revealed from the electrophoretic profile that showed apparent homogeneity and an approximate M.Wt of 14.500 and 15.00 KDa, for both types of bees under investigation, respectively and this may be attributed to the difference in its amino acid modification and the cDNA genes coding the different antimicrobial families as confirmed by Xu *et al.* (2009) in the honey bee *Apis cerana*. Originally the presented data showed that the peptide fractions proved to be effective as an antibacterial agent in the lymph of the intact bees towards the tested standard bacteria (Fig.2). These results are indicative, as in case of the corresponding hymenoptaecin of (Casteels *et al.*, 1993) and defensin of (Chernysh *et al.*, 1996), but unlike apidaecin (Casteels *et al.*,1990). Consequently the isolated peptide holds its place somewhere between the group of peptides that attack G-positive and G-negative bacteria equally well as reported by Boman (1994) and many other antibacterial peptides that seem to have clear preference for either G-negative such as apidaecins (Casteels *et al.*, 1989) and dipterocins (Bulet *et al.*,1995) or G-positive e.g. insect defensin (Lauth *et al.*,1998), lysozyme and royalisin (Fujiwara *et al.*,1990). In addition, as the corresponding results indicates the purified fractions profile proved to be comparably identical for both groups of wild *Apis florae* and carniolan *Apis carnica* bees collected from different floral locations .

The variation in the antimicrobial potential of honey samples used in this study as compared to the previous similar studies highlights that the source of the nectars may have contributed to the difference in the antimicrobial activities of honey that is, the flowers from which bees gathered nectar to produce the honey, since flora source determines many of the attributes of honey, for example flavor, aroma, color and composition. As being a natural product, the composition of honey is highly variable (NHB, 1994). Antimicrobial activity of honey is not dependent alone on its phytochemical nature i.e. tetracycline derivatives, ascorbic acid, peroxidase or amylases, streptomycin, sulfonamides which are claimed as heat labile. On the other hand, the antimicrobial effect of honey is attributed to its phenolic acid, flavonides, benzyl - alcohol, 2-hydroxy benzoic acid which are heat stable and may be active agents but their concentration in honey appears too low to solely responsible

(Heerng, 1998).

The obtained antimicrobial data of ten honey samples collected from hives of bees from different flora were generally consistent with other reports showing that honey has good antibacterial activity (Patricia *et al.*, 2005). Also, Ceyhan and Ugar (2001) tested 84 honeys against eight bacteria and two fungi showing that honey has broad-spectrum activity. In addition, these authors found that the antibacterial activity of honey was greater than that which could be attributed to the sugar content of the honey. Nzeako and Hamdi (2000) in their study of six commercial honeys found that inhibition of *S. aureus*, *E. coli* and *P. aeruginosa* did not occur at honey concentrations 40%, in contrast to the current study where all the tested bacteria showed growth inhibition up to 30% of natural honey concentrations and have shown an excellent activity against *S. aureus*. Interestingly, the obtained results of the ten honey samples under investigation revealed that *C. albicans* sensitivity were less than other bacterial organisms tested and these are consistent with the data proved by Obeseiki and Afonya, (1984) and Nzeako and Hamdi (2000).

The results shown by honey samples in relation to *S. aureus* may be important, given that in recent decades there has been a marked increase in difficult to treat skin and underlying tissue infections associated with *S. aureus* (Halco'n and Milkus, 2004). It has been informed that *S. aureus* has developed resistance against several antibiotics and that it is the principal contaminant agent in many clinical infections (Moreno *et al.*, 2005). Thus, new strategies to treat wounds infected with *S. aureus* are needed, and the possibility to use honey appears as a convenient and less costly treatment option. Poor activity of the honeys against *S. aureus* was unexpected as previous reports by Cooper *et al.* (1999). Part of the explanation for the difference in results from other studies may be due to methodological differences between studies because the agar dilution method used by these authors different from an agar well diffusion method that is used in this study. However it is also likely to be due to variation in the natural floral origin of the honey being produced. Our honey samples also exerted antimicrobial activities on *P. aeruginosa*, which were resistant to some antibiotics.

Conclusions:

Honey and antimicrobial peptide produced by honeybees either *Apis floreae* or *Apis carnica* have antimicrobial activity when tested *in vitro* against standard microorganisms. However, pharmacological standardization and clinical evaluation on the effect of honey and peptides are essential before using them as a preventive and curative measure to common diseases related to the tested bacterial species. In spite of all the positive facts associated with antimicrobial peptides there have been a few problems. Firstly, there are fewer data available on the unknown *in vitro* and *in vivo* toxicities of the peptides. Secondly, the stability of the synthesized compound formulations *in vivo* has not been studied in detail. Lastly, the cost of the production of these peptides on a large scale has been a major obstacle for quite some time.

Hence, further foci should be undertaken to identify more of such novel peptides, redesign the existing peptides to get rid of their toxicity and develop novel recombinant protocols to obtain greater yield of peptides at a lower cost.

Acknowledgments:

The authors wish to express their deep gratitude to the Research Centre of Female Sections for Science and Medical Studies King Saud University, and King Abdul Aziz City for Science and Technology for supporting and funding the present work.

References:

- Allen K., Molan P., and Reid G., 1991. A survey of the antibacterial activity of some New Zealand honeys. *J. Pharm. Pharmacol.*, 43, 817-822.
- Bals R., 2000. Epithelial antimicrobial peptides in host defense against infection. *Resp. Res.* 1, 141-50.
- Boman H., 1994. Cecropins: antibacterial peptides from insects and pigs. In: Hoffmann J.H., Janeway C.A. Natori S., editors. *Phylogenetic perspectives in immunity: the insect host defense*. R.G. Landes & Company, 43-65.
- Bulet P. Hegy G. Lambert J. Van Dorsselaer A. Hoffmann J. and Hetru C., 1995. Insect Immunity. The inducible antibacterial peptide depeptericin carries two *O*-glycans necessary for biological activity. *Biochem.*, 34, 7394-7400.

Casteels P., Ampe C., Jacobs F., and Tempst P., 1993. Functional and chemical characterization of Hymenoptaecin, an antibacterial polypeptide that is infection-inducible in the honeybee *Apis mellifera*, J.Biol.Chem., 268(10), 7044-7054.

Casteels P., Ampe C., Jacobs F., Vaeck M. and Tempst P., 1989, Apiaecins: Antibacterial peptides from honeybee. EMBO J., 8,2387-2391.

Casteels P., Ampe C., Riviere L., Damme J., Elicone C., Fleming M., Jacobs F., Tempst P., 1990. Isolation and characterization of abaecin, a major antimicrobial response peptide in the honey bee (*Apis mellifera*), Eur. J. Biochem., 187,381-386.

Ceyhan N., and Ugar A., 2001, Investigation of in vitro antimicrobial activity of honey. Riv. Biol. Forum, 94,363-372.

Chaudhary S. A., 2000. Flora of the Kingdom of Saudi Arabia. Vol. 2. Ministry of Agriculture, Riyadh.

Chernysh S., Cociancich S., Briand J.-P., Hetru C., and Bulet P., 1996. The inducible anti bacterial peptides of the hemipteran insect *Palomena prasina*: identification of a unique family of Proline-rich peptides and of a novel insect defensin. J. Insect Physiol., 42, 81-89.

Cooper R., Molan P., and Harding K., 1999. Antibacterial activity of honey against strains of *Staphylococcus aureus* from infected wounds, J. R. Soc. Med., 92, 283-285

Cudic M., and Otvos Jr. L., 2002. Intracellular targets of antibacterial peptides. Curr. Drug Targets, 3, 101-106.

Czihal P., Nimptsch A., Möllmann U., Zipfel P., and Hoffmann R., 2007. Antimicrobial activity of apidaecin peptides. Europ. Soci. of Clin. Microbiol. and Infect. Dis., 29(2), S602.

Fearon D., and Locksely R., 1996. The instructive role of innate immunity in the acquired immune response, Science, 272, 50-54.

Fujiwara S., Imai J., Fujiwara J., Yaeshima T., Kawashima T., and Kaobayashi K., 1990. A potent antibacterial protein in royal jelly. J. Biol. Chem., 265,11333-11337.

Girardin S., Sansonetti P., and Phipott D., 2002. Intracellular vs extracellular recognition of pathogens – common concepts in mammals and flies. Trends in Microbiol., 1,193-199.

Halcoń L., and Milkus K., 2004. *Staphylococcus aureus* and wounds: a review of tea tree oil as a promising antimicrobial. Am. J. Infect. Control, 32, 402-408.

Hanke M., and Schlue W.R., 1997. Lipid bilayers: methods and applications. Biological. Techniques series. San Diego, USA, Academic Press.

Heerng W., 1998. Immunochemical screening for antimicrobial drug residue in commercial honey, 123(12), 2759-62.

Hsueh P., Chen, W., and Luh K., 2005., Relationships between antimicrobial use and antimicrobial resistance in Gram-negative bacteria causing nosocomial infections from 1991–2003 at a university hospital in Taiwan. *Int. J. Antimicrob. Agents*, 26, 463–472.

Jorgensen J., and Ferraro M., 1998. Antimicrobial susceptibility testing: general principles and contemporary practices. *Clin Infect Dis.*, 26, 973–980.

Lauth X., Nesin A., Briand J.-P., Roussel J.-P., and Hertu C., 1998. Isolation, characterization and chemical synthesis of a new insect desfensin from *Chironomus plumosus* (Diptera). *Insect Biochem. Mol. Biol.*, 2, 1059 -1066.

Laemmli U.K., 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4, *Nature London*, 227, 680-685.

Lehrer R., and Ganz T. 1999. Antimicrobial peptides in mammalian and insect host defense. *Curr. Opin. Immunol.*, 11, 23-27.

Medzhitov R., and Janeway C., 2000. Advances in immunology: innate immunity. *New Eng. J. Med.*, 343, 338 - 344.

Migahid, A. M., 1996. Flora of Saudi Arabia. Vol. 2, 3rd edition, King Saud University Press, Riyadh.

Molan P., 2002. Not all honeys are the same for wound healing. *Bull. Eur. Tissue Rep. Soc.*, 9, 5–6.

Mor A., 2000. Peptide based antibiotics: a potential answer to raging antimicrobial resistance. *Drug Dev. Res.*, 50, 440–447.

Moreno J., Cruz, C., and Renzoni A., 2005. Tracking methicillin resistant *Staphylococcus aureus* clones Colombian hospitals over 7 years (1996–2003): emergence of a new dominant clone. *Int. J. Antimicrob. Agents*, 25, 457–462.

National Honey Board 1994. Honey Definitions Document. *American bee Journal*, 117– 118.

NCCLS. National Committee for Clinical Laboratory Standards, 2003. Performance standards for antimicrobial susceptibility testing. Fourteenth Informational Supplement, 4(1) Wayne, PA: NCCL

Nzeako B. and Hamdi J., 2000, Antimicrobial potential of honey on some microbial isolates, *SQU. J. Sci. Res. Med. Sci.*, 2, 75–79.

Obeseiki-Ebor E., and Afonya T., 1984. *In-vitro* evaluation of the anti candidiasis activity of honey distillate (HY-1) compared with that of some antimycotic agents. *J Pharm. Pharmacol.*, 36, 283–4.

Oren Z., and Shai Y., 1998., Mode of action of linear amphipathic α - helical antimicrobial peptides. *Biopolymers*, 47,451–63.

Patricia E. , Alexandra L., and Jenny M. , 2005. Bactericidal Activity of Different Honeys against Pathogenic Bacteria . *Archives of Medical Res.*, 36, 464–467

Reddy K., yedery R., and Aranha, C., 2004. Antimicrobial peptides; premises and promises. *Inter. J.of Antimicrob. Agents*, 24: 536-547

Snow M., and Manley-Harris M., 2004. On the nature of non-peroxide antibacterial activity in New Zealand Manuka honey. *Food Chem.*, 84, 145–147.

Tytler E., Anantharamaiah G., Walker D., Mishra V., Palgunachari M., and Segrest J., 1995. Molecular basis for prokaryotic specificity of magainin induced lysis . *Biochemistry*. 34,4393–401.

Wieprecht T., Dathe M., and Beyermann M., 1997. Peptide hydrophobicity controls the activity and selectivity of magainin 2 amide in interaction with membranes. *Biochemistry*, 36, 6124–32.

Xu P. Shi M. and Chen X., 2009. Antimicrobial peptide evolution in the Asiatic honey bee *Apis cerana*. *PLoS One* 4, 1- 9.