

# Newton's Method

Dr. O. Phillips Agboola

# Outline

1. Newton's Method.
2. Example using Newton's Method and Fixed point Iteration.
3. Practical application of Newton's Method

# Newton's Method

## Context

**Newton's** (or the *Newton-Raphson*) **method** is one of the most powerful and well-known numerical methods for solving a root-finding problem.

## Various ways of introducing Newton's method

- Graphically, as is often done in calculus.
- As a technique to obtain faster convergence than offered by other types of functional iteration.

# Newton's Method


$$p \approx p_0 - \frac{f(p_0)}{f'(p_0)} \equiv p_1.$$

## Newton's Method

This sets the stage for Newton's method, which starts with an initial approximation  $p_0$  and generates the sequence  $\{p_n\}_{n=0}^{\infty}$ , by

$$p_n = p_{n-1} - \frac{f(p_{n-1})}{f'(p_{n-1})} \quad \text{for } n \geq 1$$

# Newton's Method: Using Successive Tangents



# Newton's Method

## Stopping Criteria for the Algorithm

- Various stopping procedures can be applied in Step 3.3.
- We can select a tolerance  $\epsilon > 0$  and generate  $p_1, \dots, p_N$  until one of the following conditions is met:

$$|p_N - p_{N-1}| < \epsilon \quad (1)$$

$$\frac{|p_N - p_{N-1}|}{|p_N|} < \epsilon, \quad p_N \neq 0, \quad \text{or} \quad (2)$$

$$|f(p_N)| < \epsilon \quad (3)$$

- Note that none of these inequalities give precise information about the actual error  $|p_N - p|$ .

# Newton's Method as a Functional Iteration Technique

## Functional Iteration

- Newton's Method

$$p_n = p_{n-1} - \frac{f(p_{n-1})}{f'(p_{n-1})} \quad \text{for } n \geq 1$$

- can be written in the form

$$p_n = g(p_{n-1})$$

- with

$$g(p_{n-1}) = p_{n-1} - \frac{f(p_{n-1})}{f'(p_{n-1})} \quad \text{for } n \geq 1$$

# Newton's Method

## Example: Fixed-Point Iteration & Newton's Method


Consider the function

$$f(x) = \cos x - x = 0$$

Approximate a root of  $f$  using (a) a fixed-point method, and (b) Newton's Method



# Newton's Method & Fixed-Point Iteration



# Newton's Method & Fixed-Point Iteration

## (a) Fixed-Point Iteration for $f(x) = \cos x - x$

- A solution to this root-finding problem is also a solution to the fixed-point problem

$$x = \cos x$$

and the graph implies that a single fixed-point  $p$  lies in  $[0, \pi/2]$ .

- The following table shows the results of fixed-point iteration with  $p_0 = \pi/4$ .
- The best conclusion from these results is that  $p \approx 0.74$ .

# Newton's Method & Fixed-Point Iteration

Fixed-Point Iteration:  $x = \cos(x)$ ,  $x_0 = \frac{\pi}{4}$

$n$	$p_{n-1}$	$p_n$	$ p_n - p_{n-1} $	$e_n/e_{n-1}$
1	0.7853982	0.7071068	0.0782914	—
2	0.707107	0.760245	0.053138	0.678719
3	0.760245	0.724667	0.035577	0.669525
4	0.724667	0.748720	0.024052	0.676064
5	0.748720	0.732561	0.016159	0.671826
6	0.732561	0.743464	0.010903	0.674753
7	0.743464	0.736128	0.007336	0.672816

# Newton's Method

## (b) Newton's Method for $f(x) = \cos x - x$

- To apply Newton's method to this problem we need

$$f'(x) = -\sin x - 1$$

- Starting again with  $p_0 = \pi/4$ , we generate the sequence defined, for  $n \geq 1$ , by

$$p_n = p_{n-1} - \frac{f(p_{n-1})}{f'(p_{n-1})} = p_{n-1} - \frac{\cos p_{n-1} - p_{n-1}}{-\sin p_{n-1} - 1}.$$

- This gives the approximations shown in the following table.

# Newton's Method

Newton's Method for  $f(x) = \cos(x) - x$ ,  $x_0 = \frac{\pi}{4}$

$n$	$p_{n-1}$	$f(p_{n-1})$	$f'(p_{n-1})$	$p_n$	$ p_n - p_{n-1} $
1	0.78539816	-0.078291	-1.707107	0.73953613	0.04586203
2	0.73953613	-0.000755	-1.673945	0.73908518	0.00045096
3	0.73908518	-0.000000	-1.673612	0.73908513	0.00000004
4	0.73908513	-0.000000	-1.673612	0.73908513	0.00000000

# Newton's Method

Newton's Method for  $f(x) = \cos(x) - x$ ,  $x_0 = \frac{\pi}{4}$

$n$	$p_{n-1}$	$f(p_{n-1})$	$f'(p_{n-1})$	$p_n$	$ p_n - p_{n-1} $
1	0.78539816	-0.078291	-1.707107	0.73953613	0.04586203
2	0.73953613	-0.000755	-1.673945	0.73908518	0.00045096
3	0.73908518	-0.000000	-1.673612	0.73908513	0.00000004
4	0.73908513	-0.000000	-1.673612	0.73908513	0.00000000

- An excellent approximation is obtained with  $n = 3$ .
- Because of the agreement of  $p_3$  and  $p_4$  we could reasonably expect this result to be accurate to the places listed.

# Newton's Method in Practice

## In a practical application . . .

- an initial approximation is selected
- and successive approximations are generated by Newton's method.
- These will generally either converge quickly to the root,
- or it will be clear that convergence is unlikely.