CONFORMAL AND KILLING VECTOR FIELDS ON REAL SUBMANIFOLDS OF THE CANONICAL COMPLEX SPACE FORM \mathbb{C}^m

HANAN ALOHALI, HAILA ALODAN, AND SHARIEF DESHMUKH

ABSTRACT. In this paper, we find a conformal vector field as well as a Killing vector field on a compact real submanifold of the canonical complex space form $(\mathbb{C}^m, J, \langle , \rangle)$. In particular, using immersion $\psi : M \to \mathbb{C}^m$ of a compact real submanifold M and the complex structure J of the canonical complex space form $(\mathbb{C}^m, J, \langle , \rangle)$, we find conditions under which the tangential component of $J\psi$ is a conformal vector field as well as conditions under which it is a Killing vector field. Finally, we obtain a characterization of n-spheres in the canonical complex space form $(\mathbb{C}^m, J, \langle , \rangle)$.

1. INTRODUCTION

Conformal vector fields and Killing vector fields play a vital role in geometry of a Riemannian manifold (M, q) as well as in physics (cf. [13]). In geometry, these vector fields are used in characterizing spheres among compact or complete Riemannian manifolds (cf. [4]–[12]). A Killing vector field is said to be nontrivial if it is not parallel. The existence of a nontrivial Killing vector field on a compact Riemannian manifold constrains its geometry as well as its topology: it does not allow the Riemannian manifold (M, g) to have nonpositive Ricci curvature and if (M, g) is positively curved, its fundamental group has a cyclic subgroup (cf. [2]). In most of the cases, a conformal vector field or a Killing vector field on a Riemannian manifold (M,q) is derived through treating it as a submanifold of a Euclidean space. For example, a unit sphere S^n admits a conformal vector field that is tangential component of a constant vector field on the ambient Euclidean space R^{n+1} . Similarly, an odd dimensional unit sphere S^{2m-1} with unit normal vector field N as a hypersurface of the canonical complex space form $(\mathbb{C}^m, J, \langle , \rangle)$ admits a Killing vector field $\xi = -JN$, where J is the canonical complex structure on \mathbb{C}^m . Therefore it is an interesting question to find a conformal vector field as well as a Killing vector field on a real submanifold of a canonical complex space form

²⁰¹⁰ Mathematics Subject Classification. 53C20, 53C40.

Key words and phrases. Canonical complex space form; Ricci curvature; Conformal vector fields; Killing vector fields; Real submanifolds.

This research project was supported by a grant from the "Research Center of the Female Scientific and Medical Colleges", Deanship of Scientific Research, King Saud University.

 $(\mathbb{C}^m, J, \langle , \rangle)$. A similar study is taken up in [1] for submanifolds in a Euclidean space. Given an *n*-dimensional real submanifold (M, g) of the canonical complex space form $(\mathbb{C}^m, J, \langle , \rangle)$ with immersion $\psi : M \to \mathbb{C}^m$, we treat ψ as the position vector field of points on M in \mathbb{C}^m , and consequently we have the expression $J\psi = v + \overline{N}$, where v is the tangential component and \overline{N} is the normal component of $J\psi$ on M. This gives a globally defined vector field v on the real submanifold M.

In this paper, we study the above mentioned question for real submanifolds of the canonical complex space form $(\mathbb{C}^m, J, \langle , \rangle)$ and obtain conditions under which the vector field v is a conformal vector field (Theorems 3.1, 3.2) or a Killing vector field (Theorems 4.1, 4.3). We also use this vector field v to find a characterization of a sphere $S^n(c)$ of constant curvature c in the canonical complex space form $(\mathbb{C}^m, J, \langle , \rangle)$ (cf. Theorem 5.1). It is worth noting that the existence of the Killing vector field v not only restricts the geometry and topology of the real submanifold M but also has an influence on the dimensions of both the real submanifold and the ambient canonical complex space form $(\mathbb{C}^m, J, \langle , \rangle)$ (cf. Corollary 4.2). Finally, at the end of this paper, we give an example of a real submanifold of $(\mathbb{C}^m, J, \langle , \rangle)$ on which v is a nontrivial conformal vector field (that is, v is not Killing) and another example of a real submanifold on which v is nontrivial Killing vector field (that is, non-parallel).

2. Preliminaries

Let M be an immersed n-dimensional real submanifold of the canonical complex space form $(\mathbb{C}^m, J, \langle , \rangle)$, J and \langle , \rangle being the canonical complex structure and the Euclidean metric on \mathbb{C}^m respectively. We denote by $\mathfrak{X}(M)$ the Lie algebra of smooth vector fields on M, by $\Gamma(v)$ the space of sections of the normal bundle v of M, and by $\overline{\nabla}$ and ∇ the Riemannian connections on \mathbb{C}^m and on M respectively. Then we have the following Gauss and Weingarten equations for the real submanifold M (cf. [3]):

$$\overline{\nabla}_X Y = \nabla_X Y + h(X, Y), \quad \overline{\nabla}_X N = -A_N X + \nabla_X^{\perp} N, \quad (2.1)$$

 $X, Y \in \mathfrak{X}(M), N \in \Gamma(v)$, where h is the second fundamental form, A_N is the Weingarten map with respect to the normal $N \in \Gamma(v)$, which is related to the second fundamental form h by

$$g(A_NX,Y) = \langle h(X,Y), N \rangle, \quad X,Y \in \mathfrak{X}(M),$$

and ∇^{\perp} is the connection in the normal bundle v. The curvature tensor field R of the real submanifold M is given by

$$R(X,Y) Z = A_{h(Y,Z)} X - A_{h(X,Z)} Y, \quad X,Y,Z \in \mathfrak{X}(M).$$

The Ricci tensor field of the real submanifold M is given by

$$\operatorname{Ric}(X,Y) = ng(h(X,Y),H) - \sum_{i=1}^{n} g(h(X,e_i),h(Y,e_i)),$$

where $\{e_1, \ldots, e_n\}$ is a local orthonormal frame on M and

$$H = \frac{1}{n} \sum_{i=1}^{n} h(e_i, e_i)$$

is the mean curvature vector field of the real submanifold M.

The Ricci operator Q is a symmetric operator defined by

$$\operatorname{Ric}(X,Y) = g(Q(X),Y), \quad X,Y \in \mathfrak{X}(M).$$

Let $\psi: M \to \mathbb{C}^m$ be the immersion of the real submanifold M. Then we set

$$J\psi = v + N$$

where v is the tangential component and \overline{N} is the normal component of $J\psi$.

Now, define skew symmetric tensors φ and G, and the tensors Ψ and F as follows:

$$JX = \varphi X + FX, \quad X \in \mathfrak{X}(M),$$

$$JN = \Psi N + GN, \quad N \in \Gamma(v),$$

where

$$\begin{split} \varphi: \mathfrak{X}(M) &\longrightarrow \mathfrak{X}(M), \quad F: \mathfrak{X}(M) \longrightarrow \Gamma(\upsilon), \\ \Psi: \Gamma(\upsilon) &\longrightarrow \mathfrak{X}(M), \qquad G: \Gamma(\upsilon) \longrightarrow \Gamma(\upsilon), \end{split}$$

that is, φX , ΨN are the tangential components of JX and JN respectively and FX, GN are the normal components of JX and JN respectively.

Define a symmetric tensor C of type (1,1) by $C(X) = A_{\overline{N}}X, X \in \mathfrak{X}(M)$, and a smooth function $E: M \to \mathbb{R}$ on the real submanifold M by $E = \langle H, \overline{N} \rangle$. Then we have

$$\operatorname{tr} C = nE.$$

Lemma 2.1. Let M be an n-dimensional real submanifold of the canonical complex space form $(\mathbb{C}^m, J, \langle , \rangle)$. Then

$$\nabla_X v = \varphi X + C(X)$$
 and $\nabla_X^{\perp} \overline{N} = FX - h(X, v).$

Proof. As J is a complex structure, we have

$$\overline{\nabla}_X J\psi = J\overline{\nabla}_X \psi,$$

which in view of equation (2.1) gives

$$\nabla_X v + h(X, v) + \nabla_X^{\perp} \overline{N} - C(X) = \varphi X + FX, \quad X \in \mathfrak{X}(M).$$

Equating the tangential and the normal components we get the result.

Lemma 2.2. Let M be an n-dimensional real submanifold of the canonical complex space form $(\mathbb{C}^m, J, \langle , \rangle)$. Then for $X, Y \in \mathfrak{X}(M)$ and $N \in \Gamma(v)$, we have $(\nabla \varphi)(X, Y) = A_{F(Y)}X + \Psi(h(X, Y))$, where $(\nabla \varphi)(X, Y) = \nabla_X \varphi Y - \varphi \nabla_X Y$ $(D_X F) Y = G(h(X, Y)) - h(X, \varphi Y)$, where $(D_X F) Y = \nabla_X^{\perp} FY - F(\nabla_X Y)$ $(D_X \Psi) N = A_{G(N)}X - \varphi A_N X$, where $(D_X \Psi) N = \nabla_X \Psi(N) - \Psi(\nabla_X^{\perp} N)$

$$\left(\nabla_X^{\perp}G\right)N = F\left(A_NX\right) - h(X,\Psi(N)), \text{ where } \left(\nabla_X^{\perp}G\right)N = \nabla_X^{\perp}GN - G\left(\nabla_X^{\perp}N\right).$$

Rev. Un. Mat. Argentina, Vol. 60, No. 2 (2019)

Proof. As J is parallel, we have

$$\overline{\nabla}_{X}\left(\varphi Y + F\left(Y\right)\right) = J\left(\nabla_{X}Y + h\left(X,Y\right)\right)$$

which in view of equation (2.1) takes the form

$$\left(\nabla\varphi\right)\left(X,Y\right) + \left(D_XF\right)Y = A_{F(Y)}X + \Psi\left(h\left(X,Y\right)\right) + G\left(h\left(X,Y\right)\right) - h\left(X,\varphi Y\right),$$

which on equating the tangential and the normal components gives the first two relations. Similarly, on using $(\overline{\nabla}_X J) N = 0$, we get the remaining two.

Using Lemma 2.1, we find the divergence of the vector field v as div v = nE and consequently, we have the following:

Lemma 2.3. Let M be an n-dimensional compact real submanifold of the canonical complex space form $(\mathbb{C}^m, J, \langle , \rangle)$. Then

$$\int_{M} E \, dV = 0.$$

The following lemma is an immediate consequence of Lemma 2.1.

Lemma 2.4. Let M be an n-dimensional real submanifold of the canonical complex space form $(\mathbb{C}^m, J, \langle , \rangle)$. Then the tensor C satisfies

(i)
$$(\nabla C)(X,Y) - (\nabla C)(Y,X) = R(X,Y)v + (\nabla \varphi)(Y,X) - (\nabla \varphi)(X,Y),$$

(ii) $\sum_{i=1}^{n} (\nabla C)(e_i, e_i) = n\nabla E + Q(v) + \sum_{i=1}^{n} (\nabla \varphi)(e_i, e_i),$

where $(\nabla C)(X,Y) = \nabla_X C(Y) - C(\nabla_X Y)$, $X, Y \in \mathfrak{X}(M)$, and $\{e_1, \ldots, e_n\}$ is a local orthonormal frame of M.

Lemma 2.5. Let M be an n-dimensional real submanifold of the canonical complex space form $(\mathbb{C}^m, J, \langle , \rangle)$. Then the skew symmetric tensor φ satisfies

- (i) $(\nabla \varphi)(X,Y) (\nabla \varphi)(Y,X) = A_{FY}X A_{FX}Y,$
- (*ii*) $\sum_{i=1}^{n} (\nabla \varphi) (e_i, e_i) = n \Psi (H) + \sum_{i=1}^{n} A_{Fe_i} e_i,$

where $X, Y \in \mathfrak{X}(M)$ and $\{e_1, \ldots, e_n\}$ is a local orthonormal frame of M.

Proof. (i) Using Lemma 2.2, we get

$$(\nabla\varphi) (X,Y) - (\nabla\varphi) (Y,X) = A_{FY}X + \Psi (h (X,Y)) - A_{FX}Y - \Psi (h (Y,X))$$
$$= A_{FY}X - A_{FX}Y, \quad X,Y \in \mathfrak{X}(M).$$

(ii) As $\operatorname{tr} \varphi = 0$, we have

$$\sum_{i=1}^{n} g((\nabla \varphi) (X, e_i), e_i) = 0$$

which gives

$$\sum_{i=1}^{n} \{ g((\nabla \varphi)(e_i, X), e_i) + g(A_{Fe_i}X, e_i) - g(A_{FX}e_i, e_i) \} = 0,$$

that is,

$$\sum_{i=1}^{n} \{ g(-(\nabla \varphi)(e_i, e_i) + A_{Fe_i}e_i, X) + g(n\Psi(H), X) \} = 0.$$

Hence,

$$\sum_{i=1}^{n} (\nabla \varphi) (e_i, e_i) = n \Psi (H) + \sum_{i=1}^{n} A_{Fe_i} e_i.$$

Lemma 2.6. Let M be an n-dimensional compact real submanifold of the canonical complex space form $(\mathbb{C}^m, J, \langle , \rangle)$. Then

$$\int_{M} \left(\text{Ric} (v, v) + \|C\|^{2} - \|\varphi\|^{2} - n^{2} E^{2} \right) dV = 0.$$

Proof. Using Lemmas 2.4 and 2.5, we get

div
$$\varphi v = -\sum_{i=1}^{n} g \left(A_{F(e_i)} e_i, v \right) - ng \left(\Psi \left(H \right), v \right) - \|\varphi\|^2,$$
 (2.2)

div
$$Cv = \operatorname{Ric}(v, v) + nv(E) + ng(\Psi(H), v) + ||C||^2 + \sum_{i=1}^{n} g(A_{Fe_i}e_i, v),$$

and

$$\operatorname{div} Ev = v\left(E\right) + nE^{2}.$$
(2.3)

Using these equations, we conclude that

$$\operatorname{div} Cv = \operatorname{Ric} (v, v) + n \operatorname{div} Ev - n^2 E^2 - \operatorname{div} \varphi v - \|\varphi\|^2 + \|C\|^2,$$

which on integration gives the result. \Box

Lemma 2.7. Let M be an n-dimensional compact real submanifold of the canonical complex space form $(\mathbb{C}^m, J, \langle , \rangle)$. If v satisfies $\Delta v = -\lambda v$ for a constant $\lambda > 0$, where Δ is the Laplace operator acting on smooth vector fields on M, then

$$\int_{M} \left\{ \text{Ric}(v,v) + \lambda \|v\|^{2} - 2 \|\varphi\|^{2} - n^{2} E^{2} \right\} dV = 0$$

Proof. Using the definition of the operator C and Lemma 2.1, we have

$$\begin{aligned} \left(\nabla C\right)\left(X,Y\right) &= \nabla_X CY - C\nabla_X Y \\ &= \nabla_X \left(\nabla_Y v - \varphi Y\right) - \nabla_{\nabla_X Y} v + \varphi \nabla_X Y \\ &= \nabla_X \nabla_Y v - \nabla_{\nabla_X Y} v - \left(\nabla \varphi\right)\left(X,Y\right), \quad X,Y \in \mathfrak{X}(M). \end{aligned}$$

Taking a local orthonormal frame $\{e_1, \ldots, e_n\}$, the above equation leads to

$$\sum_{i=1}^{n} (\nabla C) (e_i, e_i) = \sum_{i=1}^{n} \left(\nabla_{e_i} \nabla_{e_i} v - \nabla_{\nabla_{e_i} e_i} v \right) - \sum_{i=1}^{n} (\nabla \varphi) (e_i, e_i)$$
$$= \Delta v - \sum_{i=1}^{n} (\nabla \varphi) (e_i, e_i)$$
$$= -\lambda v - \sum_{i=1}^{n} (\nabla \varphi) (e_i, e_i),$$

where we used the definition of the Laplace operator acting on smooth vector fields.

Now, using Lemma 2.4 (ii) and Lemma 2.5, we conclude

$$-\lambda \|v\|^{2} = \operatorname{Ric}(v, v) + nv(E) + 2g\left(\sum_{i=1}^{n} A_{F(e_{i})}e_{i}, v\right) + 2ng(\Psi(H), v),$$

and this equation together with equations (2.2) and (2.3) by integration gives

$$\int_{M} \left\{ \text{Ric}(v,v) + \lambda \|v\|^{2} - 2 \|\varphi\|^{2} - n^{2} E^{2} \right\} dV = 0.$$

3. Submanifolds with v as a conformal vector field

Recall that a smooth vector field ξ on a Riemannian manifold (M, g) is said to be a conformal vector field if the flow of ξ consists of conformal transformations of the Riemannian manifold (M, g). Equivalently, a smooth vector field ξ on a Riemannian manifold (M, g) is a conformal vector field if there exists a smooth function ρ on M that satisfies $\pounds_{\xi}g = 2\rho g$, where $\pounds_{\xi}g$ is the Lie derivative of gwith respect to ξ . The smooth function ρ is called the potential function of the conformal vector field ξ . A conformal vector field ξ is said to be a non trivial conformal vector field if the potential function ρ is not a constant. In this section, we find conditions under which the vector field v on the real submanifold M of the canonical complex space form $(\mathbb{C}^m, J, \langle , \rangle)$ is a conformal vector field.

Theorem 3.1. Let M be an n-dimensional compact real submanifold of the canonical complex space form $(\mathbb{C}^m, J, \langle , \rangle)$. If the Ricci curvature $\operatorname{Ric}(v, v)$ of M satisfies

$$\operatorname{Ric}(v, v) \ge n(n-1)E^2 + \|\varphi\|^2$$
,

then v is a conformal vector field on M.

Proof. Using Lemma 2.6, we have

$$\int_{M} \left(\text{Ric}(v,v) - n(n-1)E^{2} - \|\varphi\|^{2} + \|C\|^{2} - nE^{2} \right) dV = 0,$$

which together with the condition in the hypothesis and Schwarz's inequality $\|C\|^2 \ge nE^2$ gives

$$\operatorname{Ric}(v,v) = n(n-1)E^2 + \|\varphi\|^2$$
 and $\|C\|^2 = nE^2$.

The second equality holds if and only if C = EI, and consequently, the first equation in Lemma 2.1 reads

$$\nabla_X v = EX + \varphi X, \quad X \in \mathfrak{X}(M).$$

This equation proves that

$$(\pounds_v g)(X,Y) = 2Eg(X,Y), \quad X,Y \in \mathfrak{X}(M)$$

that is, v is a conformal vector field with potential function E.

Theorem 3.2. Let M be an n-dimensional compact real submanifold of the canonical complex space form $(\mathbb{C}^m, J, \langle , \rangle)$. If the vector field v is an eigenvector of the Laplace operator, $\Delta v = -\lambda v$, and the Ricci curvature $\operatorname{Ric}(v, v)$ satisfies

$$\operatorname{Ric}(v, v) \ge n(n-2)E^2 + \lambda ||v||^2$$
,

then v is a conformal vector field.

Proof. Lemma 2.6 implies

$$-\int_{M} \|\varphi\|^{2} dv = \int_{M} \left(-\operatorname{Ric}(v, v) - \|C\|^{2} + n^{2} E^{2}\right) dV,$$

which in view of Lemma 2.7, gives

$$\int_{M} \left(\text{Ric}(v,v) - \lambda \|v\|^{2} + 2 \|C\|^{2} - n^{2}E^{2} \right) dV = 0,$$

that is,

$$\int_{M} \left(\operatorname{Ric} \left(v, v \right) - \lambda \left\| v \right\|^{2} - n \left(n - 2 \right) E^{2} + 2 \left(\left\| C \right\|^{2} - n E^{2} \right) \right) dV = 0.$$

Thus, using the hypothesis and Schwarz's inequality $||C||^2 \ge nE^2$, we get

Ric
$$(v, v) = n (n - 2) E^2 + \lambda ||v||^2$$
 and $||C||^2 = nE^2$,

that is, C = EI. Hence, by Lemma 2.1, we get that v is a conformal vector field.

4. Submanifolds with v as a Killing vector field

Recall that a smooth vector field ξ on a Riemannian manifold (M, g) is said to be a Killing vector field if the flow of ξ consists of isometries of the Riemannian manifold (M, g). Equivalently, a smooth vector field ξ on a Riemannian manifold (M, g) is a Killing vector field if $\pounds_{\xi}g = 0$. In this section, we find conditions under which the vector field v on the real submanifold M of the canonical complex space form $(\mathbb{C}^m, J, \langle , \rangle)$ is a Killing vector field.

Theorem 4.1. Let M be an n-dimensional compact real submanifold of the canonical complex space form $(\mathbb{C}^m, J, \langle , \rangle)$. Suppose that v satisfies

- (i) v is an eigenvector of the Laplace operator with eigenvalue $-\lambda$,
- (*ii*) $\operatorname{Ric}(v, v) \ge n(n-1)E^2 + \|\varphi\|^2$,
- (iii) $\|\varphi\|^2 \ge \lambda \|v\|^2$.

Then v is a Killing vector field.

Proof. The condition (ii), in view of Theorem 3.1, implies that v is a conformal vector field with C = EI and

$$\operatorname{Ric}(v, v) = n(n-1)E^{2} + \|\varphi\|^{2}.$$
(4.1)

Now, the condition (i), $\Delta v = -\lambda v$, combined with Lemma 2.7 and the above conclusion, gives

$$\int_{M} \left(n(n-1)E^{2} + \|\varphi\|^{2} + \lambda \|v\|^{2} - 2 \|\varphi\|^{2} - n^{2}E^{2} \right) dV = 0,$$

that is,

$$\int_{M} \left((\|\varphi\|^{2} - \lambda \|v\|^{2}) + nE^{2} \right) dV = 0.$$
(4.2)

 \square

Using condition (iii), we conclude that E = 0 and consequently C = 0. Thus, Lemma 2.1 gives

$$\nabla_X v = \varphi X, \quad X \in \mathfrak{X}(M),$$

that is,

$$(\pounds_v g)(X,Y) = 0, \quad X,Y \in \mathfrak{X}(M)$$

Hence, v is a Killing vector field.

Corollary 4.2. Let M be an n-dimensional compact real submanifold of the canonical complex space form $(\mathbb{C}^m, J, \langle , \rangle)$, with positive sectional curvature. Suppose that v satisfies

- (i) v is an eigenvector of the Laplace operator with eigenvalue $-\lambda$, that is, $\Delta v =$ $-\lambda v$,
- (*ii*) Ric $(v, v) \ge n(n-1)E^2 + ||\varphi||^2$, (*iii*) $||\varphi||^2 \ge \lambda ||v||^2$.

Then either n is odd or m > n.

Proof. Notice that n < 2m. Suppose the conditions (i)–(iii) hold. Then equation (4.2) implies E = 0, $\lambda \|v\|^2 = \|\varphi\|^2$, and combining these with equation (4.1), we get

$$\operatorname{Ric}(v,v) = \lambda \|v\|^{2} = \|\varphi\|^{2}.$$
(4.3)

Now, consider the smooth function $f = \frac{1}{2} ||v||^2$, which by Lemma 2.1 and E = 0, gives the gradient $\nabla f = -\varphi v$, and we compute

$$\Delta f = -\sum_{i=1}^{n} g\left(\nabla_{e_i} \varphi v, e_i\right) = -\sum_{i=1}^{n} g\left(\nabla_{e_i} \nabla_v v, e_i\right).$$
(4.4)

Note that E = 0, as in the proof of Theorem 4.1, we get C = 0 and thus, an easy computation on using Lemma 2.1 with E = 0 gives

$$R(X,v)v = \nabla_X \nabla_v v - \varphi^2 X,$$

that is,

$$R(X, v, v, X) = g(\nabla_X \nabla_v v, X) + \|\varphi X\|^2.$$

This equation in view of equation (4.4) implies

$$\operatorname{Ric}(v,v) = -\Delta f + \|\varphi\|^2,$$

which together with equation (4.3) gives $\Delta f = 0$. Hence, f is a constant, that is, v has constant length and consequently, $\varphi v = 0$.

Rev. Un. Mat. Argentina, Vol. 60, No. 2 (2019)

If v = 0, then Lemma 2.1 implies $\varphi = 0$, that is, $J\psi = \overline{N}$, which on taking covariant derivative and using Lemma 2.1 gives JX = FX, $X \in \mathfrak{X}(M)$, and we get that M is a totally real real submanifold of \mathbb{C}^m . Hence, in this case we have $2n \leq 2m$.

If $v \neq 0$, as v is a Killing vector field of constant length $v(p) \neq 0$ for each $p \in M$, and as M is compact connected with positive sectional curvature, then M is odd-dimensional (for on an even-dimensional compact connected manifold of positive sectional curvature a Killing vector field has a zero).

Theorem 4.3. Let M be an n-dimensional compact real submanifold of the canonical complex space form $(\mathbb{C}^m, J, \langle , \rangle)$. Suppose that $v \neq 0$ is not closed and satisfies $\varphi v = 0$, with Ricci curvature

Ric
$$(v, v) \ge n (n-1) E^2 + ||\varphi||^2$$
.

Then v is a Killing vector field of constant length.

Proof. As in Theorem 3.1, the condition $\operatorname{Ric}(v, v) \ge n(n-1)E^2 + \|\varphi\|^2$ implies that v is a conformal vector field and the following hold:

 $\nabla_X v = \varphi X + EX, X \in \mathfrak{X}(M) \quad \text{and} \quad \operatorname{Ric}(v, v) = n(n-1)E^2 + \|\varphi\|^2.$ (4.5)

Using the first equation in (4.5), we get

$$R(X,Y)v = X(E)Y - Y(E)X + (\nabla\varphi)(X,Y) - (\nabla\varphi)(Y,X),$$

which gives

$$\operatorname{Ric}(Y,v) = -(n-1)Y(E) - g\left(Y, \sum_{i=1}^{n} (\nabla \varphi) \left(e_i, e_i\right)\right),$$

that is,

$$\operatorname{Ric}(v,v) = -(n-1)v(E) - g\left(v, \sum_{i=1}^{n} \left(\nabla\varphi\right)(e_i, e_i)\right).$$
(4.6)

Now, taking divergence on both sides of the equation $\varphi v = 0$, in view of equation (4.5), we have

$$-\|\varphi\|^{2} - g\left(v, \sum_{i=1}^{n} (\nabla\varphi) \left(e_{i}, e_{i}\right)\right) = 0, \qquad (4.7)$$

and inserting this equation in (4.6) leads to

$$\operatorname{Ric}(v, v) = -(n-1)v(E) + \|\varphi\|^2,$$

which on comparing with the second equation in (4.5) implies

$$v(E) = -nE^2.$$
 (4.8)

Also, using $\varphi v = 0$ in the first equation in (4.5) gives

$$\nabla_v v = Ev,\tag{4.9}$$

which in view of equations (4.5) and (4.8) leads to

$$R(X, v) v = X(E) v + nE^{2}X - (\nabla\varphi)(v, X) - E\varphi X - \varphi^{2}X$$

which on taking the inner product with v and using $\varphi(\nabla_v v) = 0$ (outcome of equation (4.9)), gives $X(E) \|v\|^2 + nE^2g(X,v) = 0$, that is,

$$||v||^2 \nabla E = -nE^2 v. (4.10)$$

Hence, as $v \neq 0$, we get $\varphi(\nabla E) = 0$, and taking divergence on both sides of this equation leads to div $(\varphi(\nabla E)) = 0$, that is,

$$g\left(\nabla E, \sum_{i=1}^{n} (\nabla \varphi) (e_i, e_i)\right) = 0,$$

which in view of equation (4.10) implies

$$-nE^{2}g\left(v,\sum_{i=1}^{n}\left(\nabla\varphi\right)\left(e_{i},e_{i}\right)\right)=0.$$

Using (4.7) in the above equation, we get

$$nE^2 \left\|\varphi\right\|^2 = 0,$$

and as v is not closed, from above equation, we conclude that E = 0, and thus equation (4.5) reads, $\nabla_X v = \varphi X$, $X \in \mathfrak{X}(M)$, which proves that v is a Killing vector field.

Moreover, if $f = \frac{1}{2} ||v||^2$, then we have

$$X(f) = g(\varphi X, v) = 0, \quad X \in \mathfrak{X}(M),$$

that is, v has constant length.

5. A CHARACTERIZATION OF SPHERES

In this section we consider an *n*-dimensional compact real submanifold M of the canonical complex space form $(\mathbb{C}^m, J, \langle , \rangle)$, and prove the following characterization for the spheres.

Theorem 5.1. Let M be an n-dimensional compact Einstein submanifold of the canonical complex space form $(\mathbb{C}^m, J, \langle , \rangle)$, n > 2. Suppose that v satisfies

(i) v is an eigenvector of the Laplace operator with eigenvalue $-\lambda < \frac{S}{n}$,

(ii) $\operatorname{Ric}(v,v) \ge n(n-1)E^2 + \|\varphi\|^2$, where S is the constant scalar curvature.

Then M is isometric to the sphere $S^{n}(c)$, for a constant c > 0.

Proof. Using Theorem 3.1, we get that v is a conformal vector field on M and equation (4.5) holds. Thus, using the first equation in (4.5), we conclude

$$\left(\nabla\varphi\right)\left(X,Y\right) = \nabla_X\nabla_Y v - \nabla_{\nabla_X Y} v - X\left(E\right)Y, \quad X,Y \in \mathfrak{X}\left(M\right), \tag{5.1}$$

where $(\nabla \varphi)(X, Y) = \nabla_X \varphi y - \varphi \nabla_X Y$. Taking sum in the above equation over a local orthonormal frame $\{e_1, \ldots, e_n\}$ on M and using $\Delta v = -\lambda v$, we get

$$\sum_{i=1}^{n} (\nabla \varphi) (e_i, e_i) = \Delta v - \nabla E = -\lambda v - \nabla E.$$
(5.2)

Rev. Un. Mat. Argentina, Vol. 60, No. 2 (2019)

426

Also, using equation (5.1), we find

$$\left(\nabla\varphi\right)\left(X,Y\right) - \left(\nabla\varphi\right)\left(Y,X\right) = R\left(X,Y\right)v + Y\left(E\right)X - X\left(E\right)Y,$$

which on choosing $X = e_i$ and taking the inner product with e_i and adding these n equations corresponding to a local orthonormal frame $\{e_1, \ldots, e_n\}$ on M, we get

$$-g\left(\sum_{i=1}^{n} \left(\nabla\varphi\right)\left(e_{i}, e_{i}\right), Y\right) = \operatorname{Ric}\left(Y, v\right) + (n-1)Y\left(E\right),$$
(5.3)

where we used the fact that φ is skew-symmetric and consequently $\sum g(\varphi e_i, e_i) = 0$, and that $g((\nabla \varphi)(X, Y), Z) = -g((\nabla \varphi)(X, Z), Y)$. Combining equations (5.2) and (5.3), we arrive at

$$Q(v) = \lambda v - (n-2)\nabla E.$$
(5.4)

Moreover, M being an Einstein manifold, $Q(v) = \frac{S}{n}v$, and thus using equation (5.4) we get

$$\nabla E = -\frac{S - n\lambda}{n(n-2)}v$$

and as S is a constant, we have $\nabla E = -cv$ for a constant c. This leads to

$$\nabla_X \left(\nabla E \right) = -c \nabla_X v = -c \left(EX + \varphi X \right), \tag{5.5}$$

that is, the Hessian of the smooth function E is given by

$$\begin{split} H_{E}\left(X,Y\right) &= -cEg\left(X,Y\right) - cg\left(\varphi X,Y\right)a, \qquad X,Y \in \mathfrak{X}\left(M\right), \\ H_{E}\left(X,Y\right) - H_{E}\left(Y,X\right) &= 2cg\left(\varphi Y,X\right). \end{split}$$

Since the Hessian is symmetric, we get $cg(\varphi Y, X) = 0$, $X, Y \in \mathfrak{X}(M)$. However, condition (i) in the hypothesis does not allow c = 0 (as c = 0 implies $S = n\lambda$); consequently we get $\varphi = 0$, which changes equation (5.5) to

$$\nabla_X (\nabla E) = -cEX, \quad X \in \mathfrak{X} (M),$$

where c is a positive constant by condition (i). Hence, by Obata's Theorem (cf. [11]), we get that M is isometric to $S^{n}(c)$.

6. Examples

In this section, we give two examples of real submanifolds of a canonical complex space form $(\mathbb{C}^m, J, \langle , \rangle)$, one admitting a conformal vector field that is not Killing and other admitting a Killing vector field that is not parallel.

(i) Consider

$$S^{2n}(c) = \left\{ x = (x_1, \dots, x_{2n+1}) \in R^{2n+1} : ||x|| = \frac{1}{\sqrt{c}}, \ c > 1 \right\}$$

and an immersion $\psi: S^{2n}(c) \to C^{n+1}$ defined by

$$\psi(x) = \left(x_1, \dots, x_{2n+1}, \sqrt{1 - \frac{1}{c}}\right),$$

which is clearly a smooth immersion. Observe that

$$T_p(S^{2n}(c)) = \{ X \in \mathbb{R}^{2n+1} : \langle X, p \rangle = 0 \}.$$

The two orthogonal unit normals N_1, N_2 for the real submanifold $S^{2n}(c)$ in C^{n+1} are given by

$$N_1 = \left(-\sqrt{c-1}x_1, \dots, -\sqrt{c-1}x_{2n+1}, \frac{1}{\sqrt{c}}\right)$$

and

$$N_2 = \left(x_1, \dots, x_{2n+1}, \sqrt{1 - \frac{1}{c}}\right).$$

Also, the standard complex structure J on C^{n+1} gives

$$J\psi = \left(-x_{n+2}, \dots, -x_{2n+1}, -\sqrt{1-\frac{1}{c}}, x_1, \dots, x_{n+1}\right)$$
(6.1)

and it is easy to check that

$$\langle J\psi, N_1 \rangle = \sqrt{c} x_{n+1}$$
 and $\langle J\psi, N_2 \rangle = 0.$

Expressing $J\psi = v + \overline{N}$, where $v \in \mathfrak{X}(S^{2n}(c))$, we get

$$v = J\psi - \sqrt{c}x_{n+1} \left(-\sqrt{c-1}x_1, \dots, -\sqrt{c-1}x_{2n+1}, \frac{1}{\sqrt{c}} \right),$$
(6.2)

that is,

$$w = \left(-x_{n+2}, \dots, -x_{2n+1}, -\sqrt{1-\frac{1}{c}}, x_1, \dots, x_{n+1}\right) + \left(\sqrt{c^2 - c}x_1x_{n+1}, \dots, \sqrt{c^2 - c}x_{n+1}x_{2n+1}, -x_{n+1}\right) = \left(\sqrt{c^2 - c}x_1x_{n+1} - x_{n+2}, \dots, \sqrt{c^2 - c}x_{n+1}^2 - \sqrt{1-\frac{1}{c}}, \sqrt{c^2 - c}x_{n+1}x_{n+2} + x_1, \dots, \sqrt{c^2 - c}x_{n+1}x_{2n+1} + x_n, 0\right).$$
(6.3)

Now, using expressions of N_1 and N_2 it is straightforward to show that

 $A_{N_1} = \sqrt{c-1}I \quad \text{and} \quad A_{N_2} = -I,$

and consequently that

$$A_{\overline{N}} = \sqrt{c^2 - c} x_{n+1} I.$$

This proves that the vector field v given by equation (6.3) satisfies

$$\pounds_v g = 2\sqrt{c^2 - c} x_{n+1} g,$$

that is, v is a conformal vector field. Note that this vector field is not a Killing vector field on $S^{2n}(c)$. To verify the last assertion, we see from the last equation that if v is Killing, $x_{n+1} = 0$, and consequently equation (6.2) gives that $v = J\psi$. Moreover, $S^{2n}(c)$ being an even-dimensional compact and connected manifold of

Rev. Un. Mat. Argentina, Vol. 60, No. 2 (2019)

positive sectional curvature, there would exist a point p where $(J\psi)(p) = 0$; using this in equation (6.1) we get c = 1, a contradiction.

(ii) Consider the unit sphere S^{2n-1} in \mathbb{R}^{2n} and an immersion $\psi: S^{2n-1} \to \mathbb{C}^m$, m > n, defined by

$$\psi(x_1,\ldots,x_n,\ldots,x_{2n}) = (x_1,\ldots,x_{2n},c_1,\ldots,c_{2m-2n}),$$

where c_i , $1 \leq i \leq 2m - 2n$, are constants and \mathbb{C}^m is identified with \mathbb{R}^{2m} . A local frame of orthonormal normal vector fields for this immersion is given by $\{N_1, N_2, \ldots, N_{2m-2n+1}\}$, where

$$N_1 = (x_1, \dots, x_{2n}, 0, \dots, 0)$$

and

$$N_{\alpha} = (0, \dots, 0, 1, 0, \dots, 0), 1$$
 at the $(2n + \alpha)^{\text{th}}$ place, $2 \le \alpha \le 2m - 2n + 1$.

Consider a complex structure J on \mathbb{C}^m defined by

$$JE = (-E(x_2), E(x_1), -E(x_4), E(x_3), \dots, -E(x_{2m}), E(x_{2m-1})), \quad E \in \mathfrak{X}(\mathbb{C}^m),$$

which makes $(\mathbb{C}^m, J, \langle, \rangle)$ a Kaehler manifold. Now set $J\psi = v + \overline{N}$, where $v \in \mathfrak{X}(S^{2n-1})$ is the tangential component and \overline{N} is the normal component of $J\psi$. We get

$$J\psi = (-x_2, x_1, \dots, -x_{2n}, x_{2n-1}, -c_2, c_1, \dots, -c_{2m-2n}, c_{2m-2n-1}),$$
(6.4)

$$\langle J\psi, N_1 \rangle = 0, \quad \langle J\psi, N_\alpha \rangle = -(-1)^\alpha c_\alpha, \quad 2 \le \alpha \le 2m - 2n + 1,$$

and consequently,

$$\overline{N} = \sum_{\alpha=1}^{2m-2n+1} \langle \overline{N}, N_{\alpha} \rangle N_{\alpha} = (0, \dots, 0, -c_2, c_1, \dots, -c_{2m-2n}, c_{2m-2n-1}).$$
(6.5)

Thus, equations (6.4) and (6.5) imply

$$v = J\psi - \overline{N} = (-x_2, x_1, \dots, -x_{2n}, x_{2n-1}, 0, \dots, 0).$$
(6.6)

Let $\overline{\nabla}$ and ∇ be the Euclidean connection on \mathbb{C}^m and the Riemannian connection on the real submanifold (S^{2n-1}, g) with respect to the induced metric g. Then using equation (6.6) we get

$$\nabla_X v = \overline{\nabla}_X v - h(X, v)$$

= (-X(x_2), X(x_1), ..., -X(x_{2n}), X(x_{2n-1}), 0, ..., 0) - h(X, v),

 $X \in \mathfrak{X}(S^{2n-1})$, where h is the second fundamental form. Taking the inner product with $Y \in \mathfrak{X}(S^{2n-1})$ in the above equation we arrive at

$$g(\nabla_X v, Y) = -X(x_2)Y(x_1) + \dots - X(x_{2n})Y(x_{2n-1}) + X(x_{2n-1})Y(x_{2n}), \quad (6.7)$$

which leads to

$$g(\nabla_X v, Y) + g(\nabla_Y v, X) = 0, \quad X, Y \in \mathfrak{X}(S^{2n-1}).$$

Thus, the vector field v satisfies

$$\pounds_v g = 0,$$

that is, v is a Killing vector field on S^{2n-1} . That the Killing vector field v is not parallel follows from equation (6.7), that is, v is a nontrivial Killing vector field.

References

- H. Alohali, H. Alodan and S. Deshmukh, Conformal vector fields on submanifolds of a Euclidean space, Publ. Math. Debrecen 91 (2017), no. 1-2, 217–233. MR 3690531.
- [2] V. Berestovskii and Y. Nikonorov, Killing vector fields of constant length on Riemannian manifolds, Siberian Math. J. 49 (2008), no. 3, 395–407. MR 2442533.
- [3] B.Y. Chen, Total Mean Curvature and Submanifolds of Finite Type, World Scientific, Singapore, 1984. MR 0749575.
- S. Deshmukh, Characterizing spheres by conformal vector fields, Ann. Univ. Ferrara Sez. VII Sci. Mat. 56 (2010), no. 2, 231–236. MR 2733411.
- [5] S. Deshmukh, Conformal vector fields and eigenvectors of Laplacian operator, Math. Phys. Anal. Geom. 15 (2012), no. 2, 163–172. MR 2915600.
- [6] S. Deshmukh, A note on hypersurfaces in a sphere, Monatsh. Math. 174 (2014), no. 3, 413–426. MR 3223496.
- [7] S. Deshmukh, Characterizations of Einstein manifolds and odd-dimensional spheres, J. Geom. Phys. 61 (2011), no. 11, 2058–2063. MR 2827109.
- [8] S. Deshmukh, F. Al-Solamy, Conformal gradient vector fields on a compact Riemannian manifold, Colloq. Math. 112 (2008), no. 1, 157–161. MR 2373435.
- [9] A. Lichnerowicz, Géométrie des groupes de transformations, Travaux et Recherches Mathématiques III, Dunod, Paris, 1958. MR 0124009.
- [10] M. Obata, Certain conditions for a Riemannian manifold to be isometric with a sphere, J. Math. Soc. Japan 14 (1962), 333–340 MR 0142086.
- [11] S. Tanno and W. Weber, Closed conformal vector fields, J. Diff. Geom. 3 (1969), 361–366. MR 0261498.
- [12] Y. Tashiro, On conformal and projective transformations in Kählerian manifolds, Tohoku Math. J. 14 (1962), 317–320. MR 0157339.
- [13] Y. Choquet-Bruhat, C. DeWitt-Morette, M. Dillard-Bleick, Analysis, Manifolds and Physics, North-Holland, New York-Oxford, 1977. MR 0467779.

H. Alohali, H. Alodan, S. Deshmukh[⊠] Department of Mathematics, College of Science, King Saud University, P.O. Box-2455, Riyadh-11451, Saudi Arabia halohali@ksu.edu.sa halodan1@ksu.edu.sa shariefd@ksu.edu.sa

Received: November 6, 2018 Accepted: January 22, 2019