

كل عام وأنتن بخير



اسم المقرر : إحصاء عام

رمزه: عرض ٢٦١

المستوى: الثالث

كلية علوم الحاسوب والمعلومات



References



Walpole, R. E., Myers, R. H., and S. L. Myers (2007), Probability and Statistics for Engineers and Scientists, 8th ed., Prentice-Hall, Inc., Upper Saddle River, New Jersey.

Program



د. محمود هندي، د.خلف سلطان،
مفاهيم لطرق التحليل الإحصائي ،
مكتبة الرشد ٤٠٠٢.



The first lecture

We will examine in this lecture :

- **Definition of Statistics.**
- **The difference between the population and the sample.**
- **Branches of Statistics.**
- **Displaying data.**


Introduction to Statistics



The reasons for the appearance of Statistics:

- Census community.
- Inventory of the wealth of individuals.
- Data on births, deaths and production and consumption.


Collection




Organization



Representation



Analysis of data



Drawing of inferences



Definition statistics



Statistics is a science of collecting , organizing , analyzing and interpreting data in order to make decisions.



Basic Definitions

Definition of a population



A **population** is the collection of all outcomes, responses, measurements, or counts that are of interest.



Types of a population

Finite population: the total number of its observations is a finite number.

- The number of students in computer science college.
- The number of cards in a deck.

Infinite population: the total number of its observations is an infinite number.

- The number of stars in sky.
- The observations obtained by measuring the atmospheric pressure every day from the past on into the future .

Definition of a sample



A **sample** is a subset of a population that is representative of the population.



Reasons draw a sample, rather than study a population

- We can not study the population :huge, destinations.
- Preservation from loss
- Less cost.
- Save time.
- More inclusive.
- More accuracy.

Branches of Statistics



Descriptive statistics is the branch of statistics that involves the organization, summarization , and display of data.

Statistical inference is the branch of statistics that involves using a sample to draw conclusions about a population.

Definition of data



Data consist of information coming from observations, counts, measurement, or responses. The singular for data is datum.


Types of data

1. **Quantitative data:** it can be measured in the usual sense like length, weight , and age.

2. **Qualitative data:** it cannot be measured in the usual sense but it can be ordered or ranked. for example:marital status,blood group and eye color



Descriptive statistics



Class Intervals	Frequency
50-59	3
60-69	5
70-79	18
80-89	16
90-99	8
Total	50

51	95	70	74	73	90	71	74	90	67
91	72	83	89	50	80	72	84	85	69
62	82	87	76	91	76	87	75	78	79
71	96	81	88	64	82	73	57	86	70
80	81	75	85	74	90	83	66	77	91



Organization Data

We will learn how to create:

- Frequency table.
- Relative frequency table.
- Percentage frequency table.

Example (1):

the following data represent the level of 60 students in a course:



D	B	E	C	D	B	D	C	E	A
B	E	C	D	B	D	D	A	E	C
C	D	A	C	E	D	C	C	D	B
D	E	D	D	A	D	D	C	D	C
D	A	B	D	B	D	C	D	C	E
D	B	C	C	E	D	C	C	D	A

Example (2):

the following data represent the marks of 50 students in a course:



51	95	70	74	73	90	71	74	90	67
91	72	83	89	50	80	72	84	85	69
62	82	87	76	91	76	87	75	78	79
71	96	81	88	64	82	73	57	86	70
80	81	75	85	74	90	83	66	77	91

Definition of frequency table (frequency distribution)



A **frequency distribution** is a table that shows **classes** or **intervals** of data entries with a count of the number of entries in each class. The **frequency f** of a class is a number of data entries in the class.



Organization qualitative data

Example (1):

the following data represent the level of 60 students in a course:



D	B	E	C	D	B	D	C	E	A
B	E	C	D	B	D	D	A	E	C
C	D	A	C	E	D	C	C	D	B
D	E	D	D	A	D	D	C	D	C
D	A	B	D	B	D	C	D	C	E
D	B	C	C	E	D	C	C	D	A



Level	tally	frequency
A		
B		
C		
D		
E		
Total		

Example (1):

the following data represent the level of 60 students in a course:



D	B	E	C	D	B	D	C	E	A
B	E	C	D	B	D	D	A	E	C
C	D	A	C	E	D	C	C	D	B
D	E	D	D	A	D	D	C	D	C
D	A	B	D	B	D	C	D	C	E
D	B	C	C	E	D	C	C	D	A



Level	tally	frequency
A		6
B		
C		
D		
E		
Total		

Example (1):

the following data represent the level of 60 students in a course:



D	B	E	C	D	B	D	C	E	A
B	E	C	D	B	D	D	A	E	C
C	D	A	C	E	D	C	C	D	B
D	E	D	D	A	D	D	C	D	C
D	A	B	D	B	D	C	D	C	E
D	B	C	C	E	D	C	C	D	A



Level	tally	frequency
A		6
B		8
C		
D		
E		
Total		



Level	tally	frequency
A		6
B		8
C		16
D		22
E		8
Total		60



Table(1): frequency table

Level	frequency
A	6
B	8
C	16
D	22
E	8
Total	60



Table(1): frequency table

Level	A	B	C	D	E	Total
frequency	6	8	16	22	8	60



Organization quantitative data



Example (2):

the following data represent the marks of 50 students in a course:

51	95	70	74	73	90	71	74	90	67
91	72	83	89	50	80	72	84	85	69
62	82	87	76	91	76	87	75	78	79
71	96	81	88	64	82	73	57	86	70
80	81	75	85	74	90	83	66	77	91

Frequency distribution for quantitative data



For large samples, we can't use the simple frequency table to represent the data.

We need to divide the data into groups or intervals or classes.

So, we need to determine:

- First step :the number of intervals (k).
- Second step :the range (R).
- Third step :the Width of the interval (w).

The number of intervals (k)



A small number of intervals are not good because information will be lost.

A large number of intervals are not helpful to summarize the data.

A commonly followed rule is that $5 \leq k \leq 20$ or the following formula may be used, $k=1+3.322(\log n)$.

We select 5 intervals in our example.

The range (R)



It is the difference between the maximum and the minimum observation (entries) in the data set.

R = the maximum entry - the minimum entry

$$\mathbf{R = 96 - 50}$$

$$\mathbf{= 46}$$



Example (2):

the following data represent the marks of 50 students in a course:

51	95	70	74	73	90	71	74	90	67
91	72	83	89	50	80	72	84	85	69
62	82	87	76	91	76	87	75	78	79
71	96	81	88	64	82	73	57	86	70
80	81	75	85	74	90	83	66	77	91

The range (R)



It is the difference between the maximum and the minimum observation(entries) in the data set.

$$R = X_{\max} - X_{\min}$$

$$R = 96 - 50$$

$$= 46$$



The Width of the interval (w)

Class intervals generally should be of the same width. Thus, if we want k intervals, then w is chosen such that $w \geq R/k$.

$$w = \frac{46}{5} = 9.4 \approx 10$$

Forth step:

Choose the minimum observation to be the lower limit of the first interval and add the width of interval to get the lower limit of the second interval and so on

the lower limit of the second interval

$$50+10=60$$

the lower limit of the third interval

$$60+10=70$$

the lower limit of the fourth interval

$$70+10=80$$

the lower limit of the fifth interval

$$80+10=90$$

Fifth step:

To find the upper limit of any interval add the following to the lower limit of interval :

$$W-1=10-1$$

$$=9$$

the upper limit of first interval

$$50+9=59$$

the upper limit of second interval

$$60+9=69$$

the upper limit of third interval

$$70+9=79$$

the upper limit of fourth interval

$$80+9=89$$

the upper limit of fifth interval

$$90+9=99$$



Class interval	tally	frequency
50-59		
60-69		
70-79		
80-89		
90-99		
Total		

Example (2):

the following data represent the marks of 50 students in a course:



51	95	70	74	73	90	71	74	90	67
91	72	83	89	50	80	72	84	85	69
62	82	87	76	91	76	87	75	78	79
71	96	81	88	64	82	73	57	86	70
80	81	75	85	74	90	83	66	77	91



Class interval	tally	frequency
50-59		3
60-69		
70-79		
80-89		
90-99		
Total		



Example (2):
the following data represent the marks of 50 students in a course:

51	95	70	74	73	90	71	74	90	67
91	72	83	89	50	80	72	84	85	69
62	82	87	76	91	76	87	75	78	79
71	96	81	88	64	82	73	57	86	70
80	81	75	85	74	90	83	66	77	91



Class interval	tally	frequency
50-59		3
60-69		5
70-79		18
80-89		16
90-99		8
Total		50

Table(2): frequency table of students' marks



Class interval	frequency
50-59	3
60-69	5
70-79	18
80-89	16
90-99	8
Total	50



Table(2): frequency table of students' marks

Class interval	50-59	60-69	70-79	80-89	90-99	Total
frequency	3	5	18	16	8	50

Definition of the true class intervals



the true lower unit = the lower limit - 0.5

the true upper unit = the upper limit + 0.5



Class interval	True class interval	frequency
50-59	49.5-59.5	3
60-69	59.5-69.5	5
70-79	69.5-79.5	18
80-89	79.5-89.5	16
90-99	89.5-99.5	8
Total		50

Definition of the Mid-interval (Midpoints)



the Mid-interval(Midpoints)

= $(\text{the lower limit} + \text{the upper limit})/2$

$$\frac{69.5 + 59.5}{2}$$

$$\frac{59.5 + 49.5}{2}$$



True class interval	Midpoints	frequency
49.5-59.5	54.5	3
59.5-69.5	64.5	5
69.5-79.5	74.5	18
79.5-89.5	84.5	16
89.5-99.5	94.5	8
Total		50

Definition of the relative frequency



the relative frequency of interval=the frequency of interval/the sum of frequencies (n)

the relative frequency table



Class interval	frequency	the relative frequency
50-59	3	0.06
60-69	5	0.10
70-79	18	0.36
80-89	16	0.32
90-99	8	0.16
Total	50	1

Definition of the percentage frequency



the percentage frequency = the relative frequency $\times 100$



the percentage frequency table

Class interval	the relative frequency	the percentage frequency
50-59	0.06	6 ← 0.06×100
60-69	0.10	10 ← 0.10×100
70-79	0.36	36
80-89	0.32	32
90-99	0.16	16
Total	1	100



Class interval	True class interval	Midpoints	frequency	the relative frequency	the percentage frequency
50-59	49.5-59.5	54.5	3	0.06	6
60-69	59.5-69.5	64.5	5	0.10	10
70-79	69.5-79.5	74.5	18	0.36	36
80-89	79.5-89.5	84.5	16	0.32	32
90-99	89.5-99.5	94.5	8	0.16	16
Total			50	1	100



Example

Class interval	frequency
16-20	100
21-25	122
26-30	900
31-35	207
36-40	795
41-45	568
46-50	322

Find from the table:

- **The Width of the interval**
- **The midpoints**
- **True class intervals**
- **The relative frequency of intervals.**
- **The percentage frequency of intervals.**

**In these lecture we
create:**

- **frequency table**
- **the percentage frequency
table**
- **the relative frequency table**



Summary of lecture