

Geometric Relaxation Time of Induced Polarization Fractal Dimension For Characterizing Shajara Reservoirs of The Shajara Formation of the Permo-Carboniferous Unayzah Group, Saudi Arabia

*Khalid Elyas Mohamed Elameen Alkhidir

**Department of Petroleum and Natural Gas Engineering, College of Engineering, King Saud University, Saudi Arabia*

Abstract

Sandstone samples were collected from the surface type section of the Shajara Formation of the Permo-Carboniferous Unayzah Group for detailed reservoir characterization. Capillary pressure experiment was performed to contact porosity and permeability was derived from the Data. Geometric relaxation time of induced polarization was calculated from the distribution of pores and the fractal dimension was proven from the relationship between water saturation and geometric relaxation time of induced polarization. In addition to field observation and obtained results of fractal dimension, the Shajara reservoirs of the Shajara Formation of the permo-Carboniferous Unayzah Group were divided here into three fractal dimension units. The Units from base to top are: Lower Shajara Geometric Relaxation Time Fractal dimension Unit, Middle Shajara Geometric Relaxation Time Fractal Dimension Unit, and Upper Shajara Geometric Relaxation Time Fractal Dimension Unit. These units were also proved by arithmetic relaxation time of induced polarization fractal dimension. It was found that the geometric relaxation time fractal dimension is similar to the arithmetic relaxation time fractal dimension of induced polarization. It was also reported that the obtained fractal dimension speeds with increasing permeability and relaxation time due to an increase in pore connectivity.

Keywords

Shajara Formation; Unayzah Group; Geometric Relaxation Time Fractal Dimension; Arithmetic Relaxation Time Fractal Dimension

Introduction

The phenomenon of electrical induced polarization (IP) in water-bearing sediments was reported by [1]. Laboratory study of induced polarization in sandstone rocks with varying salinity and shalyness was investigated by [2]. An increase of induced polarization effect with increasing permeability, increasing porosity, and decreasing shalyness was examined by [2]. An increase of permeability with increasing geometric and arithmetic induced polarization relaxation time and increasing porosity was confirmed by [3].

A new integrated model with relaxation time spectra, porosity and formation factor was proposed to predict permeability was recognized by [4]. Relationships between induced polarization relaxation time and hydraulic properties of sandstone was investigated by [5]. They reported a positive logarithmic relationship between

the relaxation time and the pore throat diameter. They also observed an increase of permeability with increasing characteristic relaxation time. Electrical conductivity, induced polarization, and permeability of the Fontainebleau sandstone was studied by [6]. They reported a positive correlation between cole-cole relaxation time and pore size.

***Corresponding author:** Khalid Elyas Mohamed Elameen Alkhidir, Department of Petroleum and Natural Gas Engineering, College of Engineering, King Saud University, Saudi Arabia. E-mail: kalkhidir@ksu.edu.sa; Tel: +966 11 4679118

Received February 18, 2018; **Accepted** February 28, 2018; **Published** March 10, 2018

Citation: Khalid Elyas Mohamed Elameen Alkhidir (2018) Geometric Relaxation Time of Induced Polarization Fractal Dimension For Characterizing Shajara Reservoirs of The Shajara Formation of the Permo-Carboniferous Unayzah Group, Saudi Arabia. SF J Petroleum 2:1.

Copyright: © 2018 Khalid Elyas Mohamed Elameen Alkhidir. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Bimodal Pore Size behavior of the Shajara Formation reservoirs of the permo-carboniferous Unayzah group was investigated by [7]. Subdivision of the Shajara reservoirs into three units based on thermodynamic fractal dimension approach and 3-D fractal geometry model of mercury intrusion technique was reported by [8]. New Finding reported from King Saud University Describe advances in Geoscience was reported by [9]. Subdivision of the Shajara reservoirs into three units: Lower Shajara Differential Capacity Fractal Dimension Unit, Middle Shajara Differential Capacity Fractal Dimension Unit, Upper Shajara Differential Capacity Fractal Dimension Unit was described by [10]. Pressure head fractal dimension for characterizing Shajara Reservoirs of the Shajara Formation of the Permo-Carboniferous Unayzah Group, Saudi Arabia was studied by [11].

Method

The geometric relaxation time of induced polarization can be scaled as

$$S_w = \left[\frac{IPTg^{1.57^{\frac{1}{2}}}}{IPTgmax^{1.57^{\frac{1}{2}}}} \right]^{3-Df} \quad 1$$

Where S_w the water saturation, $IPTg$ the geometric relaxation time of induced polarization, $IPTgmax$ the maximum geometric relaxation time of induced polarization, and Df the fractal dimension. Equation 1 can be proofed from

$$k=9.6*(IPTg*\Phi^4)^{1.57} \quad 2$$

Where k the permeability in millidarcy, 9.6 constant, $IPTg$ geometric relaxation time of induced polarization in milliseconds; Φ the porosity; and 1.57 constant.

The maximum permeability can be scaled as

$$Kmax=9.6*(IPTgmax*\Phi^4)^{1.57} \quad 3$$

Divide equation 2 by equation 3

$$\left[\frac{K}{Kmax} \right] = \frac{\left[9.6* \left[IPTg*\Phi^4 \right]^{1.57} \right]}{\left[9.6* \left[IPTgmax*\Phi^4 \right]^{1.57} \right]} \quad 4$$

Equation 4 after simplification will become

$$\left[\frac{k}{kmax} \right] = \frac{[IPTg]^{1.57}}{[IPTgmax]^{1.57}} \quad 5$$

Take the square root of equation 5

$$\sqrt{\left[\frac{k}{kmax} \right]} = \sqrt{\frac{[IPTg]^{1.57}}{[IPTgmax]^{1.57}}} \quad 6$$

Equation 6 can also be written as

$$\left[\frac{k^{\frac{1}{2}}}{kmax^{\frac{1}{2}}} \right] = \left[\frac{IPTg^{1.57^{\frac{1}{2}}}}{IPTgmax^{1.57^{\frac{1}{2}}}} \right] \quad 7$$

Take the Logarithm of equation 7

$$\log \left[\frac{k^{\frac{1}{2}}}{kmax^{\frac{1}{2}}} \right] = \log \left[\frac{IPTg^{1.57^{\frac{1}{2}}}}{IPTgmax^{1.57^{\frac{1}{2}}}} \right] \quad 8$$

$$\text{But; } \log \left[\frac{k^{\frac{1}{2}}}{kmax^{\frac{1}{2}}} \right] = \log \frac{S_w}{3-Df} \quad 9$$

Insert equation 9 into equation 8

$$\log \frac{S_w}{3-Df} = \log \left[\frac{IPTg^{1.57^{\frac{1}{2}}}}{IPTgmax^{1.57^{\frac{1}{2}}}} \right] \quad 10$$

If we remove the Log from equation 10

$$S_w = \left[\frac{IPTg^{1.57^{\frac{1}{2}}}}{IPTgmax^{1.57^{\frac{1}{2}}}} \right]^{3-Df} \quad 11$$

Equation 11 the proof of equation 1 which relates the water saturation, the geometric relaxation time of induced polarization, the maximum geometric relaxation time of induced polarization, and the fractal dimension. The arithmetic relaxation time of induced polarization can be scaled as

$$S_w = \left[\frac{IPTa^{(1.74)^{\frac{1}{2}}}}{IPTamax^{(1.74)^{\frac{1}{2}}}} \right]^{3-Df} \quad 12$$

Where Sw the water saturation, IPTa the induced polarization relaxation time arithmetic , IPTamax the maximum induced polarization relaxation time arithmetic , and Df the fractal dimension.

Equation 12 can be proofed from

$$K=11.1(IPTa * \Phi^4)^{1.74} \quad 13$$

Where k the permeability in millidarcy, 11.1 constant, IPTa the induced polarization relaxation time in milliseconds, Φ the porosity, and 1.74 constant.

The maximum permeability can be scaled as

$$K_{max}=11.1(IPT_{max} * \Phi^4)^{1.74} \quad 14$$

Divide equation 13 by equation 14

$$\left[\frac{K}{K_{max}} \right] = \left[\frac{11.1(IPTa * \Phi^4)^{1.74}}{11.1(IPT_{max} * \Phi^4)^{1.74}} \right] \quad 15$$

Equation 15 after simplification will become

$$\left[\frac{k}{k_{max}} \right] = \left[\frac{IPTa^{1.74}}{IPT_{max}^{1.74}} \right] \quad 16$$

Take the square root of equation 16

$$\sqrt{\left[\frac{k}{k_{max}} \right]} = \sqrt{\left[\frac{IPT^{1.74}}{IPT_{max}^{1.74}} \right]} \quad 17$$

Equation 17 can also be written as

$$\left[\frac{k^{\frac{1}{2}}}{k_{max}^{\frac{1}{2}}} \right] = \left[\frac{IPTa^{(1.74)\frac{1}{2}}}{IPT_{max}^{(1.74)\frac{1}{2}}} \right] \quad 18$$

Take the logarithm of equation 18

$$\text{Log} \left[\frac{k^{\frac{1}{2}}}{k_{max}^{\frac{1}{2}}} \right] = \text{Log} \left[\frac{IPTa^{(1.74)\frac{1}{2}}}{IPT_{max}^{(1.74)\frac{1}{2}}} \right] \quad 19$$

$$\text{But; Log} \left[\frac{k^{\frac{1}{2}}}{k_{max}^{\frac{1}{2}}} \right] = \frac{\text{LogSw}}{3 - Df} \quad 20$$

Insert equation 20 into equation 19

$$\frac{\text{LogSw}}{3 - Df} = \text{Log} \left[\frac{IPTa^{(1.74)\frac{1}{2}}}{IPT_{max}^{(1.74)\frac{1}{2}}} \right] \quad 21$$

If we remove the Log from equation 21

$$Sw = \left[\frac{IPTa^{(1.74)\frac{1}{2}}}{IPT_{max}^{(1.74)\frac{1}{2}}} \right] \quad 22$$

Equation 22 the proof of equation 12 which relates the water saturation Sw. the arithmetic relaxation time of induced polarization IPTa, the maximum arithmetic relaxation time of induced polarization IPTa max, and the fractal dimension Df.

Porosity was measured on collected sandstone samples from the surface type section of the Shajara formation of the permo-carboniferous Unayzah Group. Permeability was calculated from the measured capillary pressure data. The objective of this paper is to calculate the geometric relaxation time and arithmetic relaxation time of induced polarization from the distribution of pores and to calculate the fractal dimension from the relaxation time.

Results and Discussion

Based on field observation the Shajara Reservoirs of the Shajara Formation of the Permo-Carboniferous Unayzah Group were divided into three units as described in figure 1. These units from base to top are: Lower Shajara Reservoir, Middle Shajara reservoir, and Upper Shajara Reservoir.

Their acquired results of the geometric relaxation time fractal dimension and arithmetic relaxation time fractal dimension of induced polarization are displayed in table 1. Based on the attained results it was found that the geometric relaxation time fractal dimension is equal to the arithmetic relaxation time fractal dimension. The maximum value of the fractal dimension was found to be 2.7872 assigned to sample SJ13 from the Upper Shajara Reservoir as verify in Table 1. Whereas the minimum value of the fractal dimension was reported from sample SJ3 from the Lower Shajara reservoir as displayed in table 1. The geometric relaxation time induced polarization fractal dimension and arithmetic relaxation time fractal dimension were observed to increase with increasing permeability as proofed in table 1 owing to the possibility

of having interconnected channels.

The Lower Shajara reservoir was denoted by four sandstone samples out of six, label as SJ1, SJ2, SJ3 and SJ4 as confirmed in figure1. Their geometric fractal dimension and arithmetic time fractal dimension values are proofed in table 1. As we proceed from sample SJ2 to SJ3 a pronounced reduction in permeability due to

compaction was reported from 1955 md to 56 md which reflects decrease in geometric and arithmetic relaxation time fractal dimension from 2.7748 to 2.4379 as specified in table 1. Again, an increase in grain size and permeability was recorded from sample SJ4 whose geometric fractal dimension and arithmetic fractal dimension was found to be 2.6843 as described in table 1.

Table 1: Petrophysical Model Showing the Three Shajara Reservoir Units with their Corresponding Values of Geometric and Arithmetic Relaxation Time Fractal Dimension of Induced Polarization

Reservoir	Sample	Φ %	K (md)	IPTg fractal dimension	IPTa fractal dimension
Upper Shajara Reservoir	SJ13	25	973	2.7872	2.7872
	SJ12	28	1440	2.7859	2.7859
	SJ11	36	1197	2.7586	2.7586
Middle Shajara Reservoir	SJ9	31	1394	2.7786	2.7786
	SJ8	32	1344	2.7752	2.7752
	SJ7	35	1472	2.7683	2.7683
Lower Shajara Reservoir	SJ4	30	176	2.6843	2.6843
	SJ3	34	56	2.4379	2.4379
	SJ2	35	1955	2.7748	2.7748
	SJ1	29	1680	2.7859	2.7859

In contrast, the Middle Shajara reservoir which is separated from the Lower Shajara reservoir by an unconformity surface as shown in figure 1. It was designated by three samples out of four, namely SJ7, SJ8, and SJ9 as illustrated in figure 1. Their geometric relaxation time induced polarization fractal dimensions (IPTg) and arithmetic relaxation time fractal dimensions (IPTa) show similarities as delineated in Table 1. Their fractal dimensions are higher than those of samples SJ3 and SJ4 from the Lower Shajara Reservoir due to an increase in their permeability as explained in table 1.

On the other hand, the Upper Shajara reservoir is separated from the Middle Shajara reservoir by yellow green mudstone as revealed in figure 1. It is defined by three samples so called SJ11, SJ12, SJ13 as explained in table 1. Moreover, their geometric relaxation time fractal dimension and arithmetic relaxation time fractal dimension are also higher than those of sample SJ3 and SJ4 from the Lower Shajara Reservoir due to an increase in their permeability as clarified in table 1.


Overall a plot of geometric relaxation time fractal dimension versus arithmetic relaxation time fractal dimension as shown in figure 2 reveals three permeable zones of varying Petrophysical properties. The higher

fractal dimension zone with fractal dimension higher than 2.75 (Figure 2) corresponds to the complete Upper Shajara Reservoir, total Middle Shajara Reservoir and Sample SJ1 and SJ2 from the lower Shajara Reservoir. The middle fractal dimension zone with a value of about 2.68 (Figure 2) resembles sample SJ4 from the lower Shajara reservoir. The lower fractal dimension value 2.43 allocates to sample SJ3 from the Lower Shajara reservoir as shown in (Figure 2).

Figure 1: Surface type section of the Shajara reservoir of the Shajara formation of the Permo-Carboniferous Unayzah Group, Wadi Shajara, Qusayba area, al Qassim district, Saudi Arabia, N 2652 17.4, E 4336 18.

AGE	Fm.	Mbr.	unit	LITHO-LOGY	DESCRIPTION		
Late Permian	Khuff Formation	Hagay1 Member			Limestone : Cream, dense, burrowed, thickness 6.56' Sub-Khuff unconformity.		
Late Carboniferous - Permian	Shajara Formation	Upper Shajara Member	Upper Shajara mudstone		Mudstone : Yellow, thickness 17.7'		
			Upper Shajara Reservoir	SJ13▲	Sandstone : Light brown, cross-bedded, coarse-grained, poorly sorted, porous, friable, thickness 6.5'		
				SJ12▲	Sandstone : Yellow, medium-grained, very coarse-grained, poorly, moderately sorted, porous, friable, thickness 13.1'		
			Middle Shajara Member	Middle Shajara mudstone	SJ11▲	Mudstone : Yellow-green, thickness 11.8'	
						Mudstone : Yellow, thickness 1.3'	
					Mudstone : Brown, thickness 4.5'		
		Middle Shajara Reservoir		SJ10▲	Sandstone : Light brown, medium-grained, moderately sorted, porous, friable, thickness 3.6'		
				SJ9▲ SJ8▲	Sandstone : Yellow, medium-grained, moderately well sorted, porous, friable, thickness 0.9'		
		Lower Shajara Member	Lower Shajara Reservoir	SJ7▲	Sandstone : Red, coarse-grained, medium-grained, moderately well sorted, porous, friable, thickness 13.4'		
				SJ6▲	Sandstone : White with yellow spots, fine-grained, hard, thickness 2.6'		
				SJ5▲ SJ4▲	Sandstone : Limonite, thickness 1.3' Sandstone : White, coarse-grained, very poorly sorted, thickness 4.5'		
				SJ3▲ SJ2▲	Sandstone : White-pink, poorly sorted, thickness 1.6' Sandstone : Yellow, medium-grained, well sorted, porous, friable, thickness 3.9'		
				SJ1▲	Sandstone : Red, medium-grained, moderately well sorted, porous, friable, thickness 11.8'		
		Early Devonian	Tasil Formation				Sub-Unayzah unconformity. Sandstone : White, fine-grained.

Figure 2: Arithmetic relaxation time fractal dimension versus geometric relaxation time fractal dimension of induced polarization showing three permeable zones.



Conclusions

- The sandstones of the Shajara Reservoirs of the Shajara formation of the permo-Carboniferous Unayzah group were divided here into three units based on geometric relaxation time fractal dimension of induced polarization.
- The Units from base to top are: Lower Shajara Geometric Relaxation time Fractal dimension Unit, Middle Shajara Relaxation Time Fractal Dimension Unit, and Upper Shajara Relaxation time Fractal Dimension Unit.
- These units were also proved by arithmetic relaxation time fractal dimension of induced polarization.
- The fractal dimension was found to increase with increasing grain size and permeability.

Acknowledgement

The author would like to thank King Saud University, college of Engineering, Department of Petroleum and natural gas engineering, Department of Chemical Engineering, Research Centre at College of Engineering, and King Abdullah Institute for research and consulting Studies for their supports.

References

1. Schlumberger C (1920) Etude sur la prospection électrique des sous-sols Gauthier-Villars. Paris.
2. Schoepel RJ, Thrasher JE (1966) Laboratory study of induced polarization in sandstone rocks with varying salinity and shaliness, SPWLA 7th Annual Logging Symposium. Oklahoma.
3. Maosong T, Weinan W, Li L, et al. (2004) Estimation of permeability of shaly sand reservoir from induced polarization relaxation time spectra. J Petrol Sci Eng 45: 1-10.
4. Liu X, Kong Li, Zhang P (2014) Permeability estimation using relaxation time spectra derived from differential evolution inversion. J Geophys Eng 11: 1-8.
5. Konstantin Titov, Andrey Tarasov, Yuri Ilyin, et al. (2010) Relationships between induced polarization relaxation time and hydraulic properties of sandstone. Geophys J Int 180: 1095-1106.
6. André Revil, Pauline Kessouri, Carlos Torres Verdín (2014) Electrical conductivity, induced polarization, and permeability of the Fontainebleau sandstone. Geophysics 79: D301-D318.

7. Al Khidir KE, Al Laboun A, Al Qurishi A, et al. (2011) Bimodal Pore Size behavior of the Shajara formation reservoirs of the Permo-Carboniferous Unayzah group, Saudi Arabia. J Pet Explor Prod Technol 1: 1-9.
8. Al Khidir KE, Benzagouta M, Al Qurishi A, et al. (2013) Characterization of heterogeneity of the Shajara Reservoirs of the Shajara formation of the Permo-Carboniferous Unayzah group. ARAB J GEOSCI 6: 3989-3995.
9. Geoscience (2013) New findings reported from King Saud. University describe advances in geoscience. Al Khidir et al. science letter 359.
10. Al Khidir KE, Benzagouta MS, AlQuraishi AA, et al. (2013) Differential capacity fractal dimension and water saturation fractal dimension as parameters for reservoir characterization: Shajara formation of the Permo-Carboniferous Unayzah Group as a case study. 10th Meeting of the Saudi Society for Geoscience Geosciences for Sustainable Development KFUPM Campus. Dhahran. Saudi Arabia.
11. Al Khidir KE (2017) Pressure Head Fractal Dimension for Characterizing Shajara Reservoirs of the Shajara Formation of the Permo-Carboniferous Unayzah Group, Saudi Arabia. Arch Pet Environ Biotechnol 2.

Citation: Khalid Elyas Mohamed Elameen Alkhidir (2018) Geometric Relaxation Time of Induced Polarization Fractal Dimension For Characterizing Shajara Reservoirs of The Shajara Formation of the Permo-Carboniferous Unayzah Group, Saudi Arabia. SF J Petroleum 2:1.