


AUTHOR QUERY FORM

 ELSEVIER	Journal: NEFRO	Please e-mail your responses and any corrections to:
	Article Number: 240	E-mail: corrections.eses@elsevier.thomsondigital.com

Dear Author,

Please check your proof carefully and mark all corrections at the appropriate place in the proof (e.g., by using on-screen annotation in the PDF file) or compile them in a separate list. Note: if you opt to annotate the file with software other than Adobe Reader then please also highlight the appropriate place in the PDF file. To ensure fast publication of your paper please return your corrections within 48 hours.

For correction or revision of any artwork, please consult <http://www.elsevier.com/artworkinstructions>.

Any queries or remarks that have arisen during the processing of your manuscript are listed below and highlighted by flags in the proof. Click on the 'Q' link to go to the location in the proof.

Location in article	Query / Remark: click on the Q link to go Please insert your reply or correction at the corresponding line in the proof
<p>Q1</p> <p>Q2</p> <p>Q3</p>	<p>Reference(s) given here were noted in the reference list but are missing from the text – please position each reference in the text or delete it from the list.</p> <p>Please confirm that given name and surname are correctly identified. The different colors indicate whether tagged as first or last name. Please note that proper identification is key for correct indexing of the article.</p> <p>Please check and confirm the text in 'Conflicts of interest'.</p> <p>Uncited reference: This section comprises reference that occurs in the reference list but not in the body of the text. Please cite this reference in the text or, alternatively, delete it.</p>
	<div style="border: 1px solid black; padding: 10px;"> <p>Please check this box or indicate your approval if you have no corrections to make to the PDF file</p> <div style="float: right; text-align: center;"> <input type="checkbox"/> </div> </div>

Thank you for your assistance.



Review

Reno-protective effects of propolis on gentamicin-induced acute renal toxicity in swiss albino mice

Q1 **Badr Abdullah Aldahmash^{a,*}, Doaa Mohamed El-Nagar^b, Khalid Elfakki Ibrahim^a**

^a King Saud University, College of Science, Department of Zoology, Riyadh, Saudi Arabia

^b Ain Shams University, College of Girls for Science, Arts and Education, Department of Zoology, Cairo, Egypt

ARTICLE INFO

Article history:

Received 27 July 2015

Accepted 13 June 2016

Available online xxx

Keywords:

Gentamicin

Propolis

Kidney

Apoptosis

Oxidative stress

ABSTRACT

Background: Kidney is a vital organ which plays an important and irreplaceable role in detoxification and removal of xenobiotics. And therefore is vulnerable to develop various forms of injuries. Hence, making it immensely important to search for natural reno-protective compounds.

Objectives: This study therefore, aims to evaluate the reno-protective properties of propolis against gentamicin induced renal toxicity in mice.

Methods: Three groups of 10 male mice each were used for this study. First group served as control, the second group (Gm group) was administered orally 80 mg/kg body weight gentamicin for 7 days, and the third group (GmP group) was administered same dose of gentamicin with propolis (500 mg/kg body weight) for 7 days. Various parameters were used to study the renal toxicity.

Results: Gentamicin caused significant renal damage as evident by the rise in BUN levels, diminished glomeruli hypocellularity, moderately dilated tubules, and mild loss of brush border, severe infiltration, extensive tubular degeneration and presence of tubular cast. Histochemistry results show presence of collagen and reticular fibres. Immunohistochemical reactions show kidney injury (Kim-1 gene-expression), oxidative stress (MDA gene-expression), and an increase in apoptosis (caspase-3 gene-expression). Co-administration of propolis with gentamicin showed significant decrease in BUN levels, appearance of healthy glomeruli with normal cellularity, reduction of tubular injury, decrease of collagen and reticular fibres deposition, reduction of apoptosis, kidney injury and oxidative stress.

Conclusion: Results presented in this study clearly show the reno-protective role of propolis against gentamicin-induced toxicity on mice kidney.

© 2016 Published by Elsevier España, S.L.U. on behalf of Sociedad Española de Nefrología.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author.

E-mail address: badr.zool.ksu@gmail.com (B.A. Aldahmash).

<http://dx.doi.org/10.1016/j.nefro.2016.06.004>

0211-6995/© 2016 Published by Elsevier España, S.L.U. on behalf of Sociedad Española de Nefrología. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Efectos renoprotectores del propóleo sobre la toxicidad renal aguda inducida por gentamicina en ratones albinos suizos

R E S U M E N

Palabras clave:

Gentamicina
Propóleo
Riñón
Apoptosis
Estrés oxidativo

Antecedentes: El riñón es un órgano vital que desempeña una función importante e insustituible en la desintoxicación y la eliminación de los xenobióticos y, por lo tanto, es vulnerable a desarrollar diversas formas de lesión. Esto hace muy importante la búsqueda de compuestos renoprotectores naturales.

Objetivos: Este estudio tiene como objetivo evaluar las propiedades renoprotectoras del propóleo contra la toxicidad renal inducida por gentamicina en ratones.

Métodos: Para este estudio se utilizaron 3 grupos de 10 ratones macho en cada uno. El primer grupo sirvió como control, el segundo grupo (grupo Gm) recibió 80 mg/kg de peso corporal de gentamicina por vía oral durante 7 días y el tercer grupo (grupo GmP) recibió la misma dosis de gentamicina con propóleo (500 mg/kg de peso corporal) durante 7 días. Se utilizaron varios parámetros para estudiar la toxicidad renal.

Resultados: La gentamicina causó daño renal significativo, como demostró el aumento de los niveles de nitrógeno ureico en sangre, la disminución de la hipocelularidad glomerular, los túbulos moderadamente dilatados y la pérdida leve del borde en cepillo, la infiltración grave, la degeneración tubular extensa y la presencia de cilindros tubulares. Los resultados de la histoquímica muestran presencia de colágeno y fibras reticulares. Las reacciones inmuno-histoquímicas muestran lesión renal (expresión del gen Kim-1), estrés oxidativo (expresión del gen MDA) y un aumento de la apoptosis (expresión del gen caspasa-3). La administración concomitante de propóleo con gentamicina mostró disminución significativa de los niveles de nitrógeno ureico en la sangre, aspecto de glomérulos sanos con celularidad normal, reducción de la lesión tubular, disminución de colágeno y deposición de fibras reticulares, reducción de la apoptosis, daño renal y estrés oxidativo.

Conclusión: Los resultados presentados en este estudio muestran claramente la función renoprotectora del propóleo contra la toxicidad inducida por gentamicina en el riñón de los ratones.

© 2016 Publicado por Elsevier España, S.L.U. en nombre de Sociedad Española de Nefrología. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/s/by-nc-nd/4.0/>).

Introduction

Gentamicin is commonly used aminoglycoside antibiotic for the treatment of various bacterial infections. The recommended routes of administration of gentamicin are intravenous, intramuscular, intraperitoneal or topical as it is not sufficiently absorbed by the intestinal tract.^{1,2} However, potential clinical use of gentamicin is limited due to gentamicin-induced toxicity, even at doses slightly higher than recommended doses. Gentamicin can cause tissue injury such as nephrotoxicity, ototoxicity^{3,4} and liver toxicity,⁵ possibly through generation of free oxygen radicals. Nephrotoxicity of gentamicin arises due to its accumulation in renal cortical tubular epithelial cells.² Although the pathogenesis of gentamicin-induced acute kidney injury (AKI) has been the focus of a large number of studies, the underlying mechanisms are not yet fully elucidated. Recent studies suggest that gentamicin nephrotoxicity is a complex and multifaceted process in which gentamicin triggers cellular responses involving multiple pathways that culminate in renal damage and necrosis.^{6,7} Therefore, a number of different molecular markers are being used to assess the kidney injury including Kidney Injury Molecule-1 (KIM-1), markers for apoptosis and oxidative stress.^{8–10}

Several agents and strategies have been attempted to ameliorate gentamicin nephrotoxicity^{11–13} with main focus on

the use of various antioxidant agents including the extracts from medicinal plants with antioxidant properties.¹¹ However, none of these have been found safe/suitable for clinical practice due to known and unexplored side effects. Propolis is a gum like substance gathered by bees from various plants and varies in colour from light yellow to dark brown,¹⁴ possesses a broad spectrum of biological activities such as anti-hepatitis and anti-arthritis, and is also known to enhance immune system.^{15–17} This biological activity may be attributed to its constituents obtained from plants, mainly phenolic compounds such as flavonoids. Flavonoids are well-known antioxidant possessing free radical scavenging and metal chelating activity.¹⁸ At least 38 different flavonoids have been reported in propolis.¹⁹ Some components of the propolis are absorbed and circulate in the blood and behave as hydrophilic antioxidant and save vitamin C.²⁰ The present study therefore evaluates the potential of propolis when administered orally to protect the kidney against the harmful effects and acute nephrotoxicity of gentamicin in swiss albino mice.

Materials and methods

Animals

Swiss albino male mice weighing 25±g were used for the experiment. These animals were acclimated to 22±1°C and

were maintained under 12-h periods of light and dark each, with free access to clean water and commercial mice food. The animals were housed in polypropylene cages inside a well-ventilated room.

Experimental design

Mice were randomly distributed into three groups, each containing 10 mice. Group 1 mice received saline and served as control group while group 2 mice received intraperitoneal injection of gentamicin at dose of 80 mg/kg for 7 consecutive days and this group was marked as Gm group. Mice in group 3 were treated as group 2 and were additionally co-administered with 500 mg/kg of propolis one hour-post gentamicin injection and this group was marked as GmP group.

Kidney index

Following treatments as described above, each mouse was weighed; kidneys were removed and weighed. Finally, the kidney index was calculated by dividing the left kidney weight by the body weight and then multiplying by 100 and the results were statistically analyzed by SPSS software (SPSS Inc.).

Biochemical analysis

Blood samples for the measurement of blood chemistry were drawn into prechilled tubes containing EDTA, and immediately placed on ice. Serum in the samples was separated by centrifugation at 3000 rpm and stored at -80°C until assay. Serums were used for the estimation of blood urea nitrogen (BUN) and creatinine.

Histopathological analysis

Histopathological preparation

Kidneys were collected and cut into small pieces, fixed in 10% neutral buffered formalin. Following fixation, specimens were dehydrated, embedded in wax, and then sectioned to $5\mu\text{m}$ thicknesses. Sections were stained with haematoxylin and eosin, Masson's Trichrome stain and Gomori silver technique. Digital images of kidneys tissues were obtained using a light microscope at a magnification of $400\times$.

Gene-expression localization studies

Paraffin embedded kidney sections were deparaffinized in xylene and rehydrated in descending grades of alcohol and finally distilled water. Sections were then heated in citrate buffer (pH 6) in microwave for 5 min, washed with PBS buffer for 5 min and were incubated in peroxidase blocking solution for 10 min. After blocking sections were incubated overnight at 4°C with diluted primary antibody (anti-caspase3 ab13585, anti-Kim-1, rabbit polyclonal antibody ab78494, anti-malondialdehyde ab194225). Sections were then incubated with biotinylated goat anti-mouse secondary antibody (ab128976) for 30 min, followed by incubation in avidin-biotin complex for 30 min. Finally DAB (ab64238) was used as chromogenic substrate for the detection of Ab binding. Stained sections were counter stained with Mayer's haematoxylin, and

dehydrated within ascending grades of alcohol and cleared with two changes of xylene, mounted with cover slip based on DPX mountant, (all reagents from Abcam). Kidney sections were examined under microscope for brown immunoreactivity colour and photos at $400\times$ magnification.

Renal pathology analysis

Formalin-fixed kidney sections ($5\mu\text{m}$) were stained with haematoxylin and eosin to distinguish cell nuclei and digital images of glomeruli were recorded at $400\times$ magnification using a light microscope. Glomerular tuft areas were measured by microscopy computer system (Motic-2000), while, glomerular cellularity was determined by counting the number of nuclei in 20 hilar glomerular tuft cross-sections per animal.

Pathological score for tubular injury

For determining pathological score haematoxylin eosin stained preparations were evaluated under light microscope. Dilated tubules, loss of brush border, tubular casts, leukocytic infiltration and tubular degeneration in the cortical area were scored as described by Biswas et al.²¹ The scoring system used is described as follows. Kidneys showing no tubular injury were marked 0. While, kidneys exhibiting mild tubular injury $\leq 10\%$ were given a score of 1. Similarly, kidneys showing mild (10–25%), moderate (26–50%), extensive ($=51-75\%$) and severe ($\geq 75\%$) tubular injuries were assigned a score of 2, 3, 4 and 5, respectively. Tubular cast scored as 0 = negative and 1 = positive.

Histochemical and immunohistochemical analysis

Kidney sections stained with Mason's Trichrome, Gomori silver technique, Caspase 3 in glomeruli and tubules, Kim-1 in glomeruli and tubules, and malondialdehyde gene-expressions by ABC method were quantitatively scored as – = none, + = little, ++ = mild and +++ = intense.

Statistical analysis

Statistical evaluation was carried out by using one-way ANOVA test and SPSS (16.0 software), all values were expressed as mean \pm SD. Values of $p < 0.05$ were accepted as significant.

Results

Kidney index and biochemical analysis

Kidney index showed insignificant difference between control and gentamicin experimental groups (Table 1). Blood urea nitrogen (BUN) levels increased significantly ($p < 0.05$) in gentamicin (Gm) and gentamicin with propolis (GmP) groups compared to control group. The Kidney index for Control, Gm and GmP group was 22, 41 and 38, respectively (Table 1). It is important to note that there was an insignificant decrease of Kidney index in GmP group (38) compared to Gm group (41). Creatinine levels showed insignificant difference in gentamicin experimental groups compared to control group (Table 1).

Table 1 – Change in kidney index, BUN blood serum and creatinine in blood serum following, treatment with gentamicin alone and along with propolis, in mice.

Parameters	Control	Gentamicin (Gm group)	Gentamicin + propolis (GmP group)
Kidney index	0.60 (± 0.03)	0.62 (± 0.08), +3.3% [†]	0.65 (± 0.07), +8.3%
BUN (mg/dl)	22.6 (± 0.6)	41 (± 2.0), +81.4%	38 (± 3.0), +68.1%
Creatinine (mg/dl)	0.4 (± 0.03)	0.36 (± 0.05), –10%	0.33 (± 0.02), –17.5%

Values presented in parenthesis as mean \pm SD (standard deviation).

* Significant difference (p value < 0.5) compared to control group.

[†] Values in parenthesis show % increase (+), or decrease (–), when compared to control.

Table 2 – Glomerular areas and glomerular cellularity of Control, Gm and GmP mice groups.

Parameters	Control	Gentamicin (Gm group)	Gentamicin + propolis (GmP group)
Glomerular area (μm^3)	4.3 (± 2.8)	2.4 (± 1.5), –44% [†]	3.8 (± 2), –11%
Glomerular cellularity (cells/gcs)	30 (± 1.2)	20 (± 0.9), –33%	27 (± 1.2), –10%

Values in parenthesis are mean \pm SD (standard deviation).

* Significant difference (p value < 0.5) compared to control group.

[†] Values in parenthesis show % decrease (–), as compared to control.

Histopathological analysis


Glomerular analysis

Control kidney exhibit normal glomeruli score ($4.3 \mu\text{m}^3$), glomerular area, and (30 C/gcs) cells (Table 2) with abundant podocytes, mesangial cells with healthy mesangial matrix in between and normal capsular space (Fig. 1a). Kidney sections of Gm mice group showed diminished glomeruli that scored significant decrease in area ($2.4 \mu\text{m}^3$) and cellularity (20 C/gcs) compared to control group $p < 0.05$, in addition to severe degeneration in mesangial matrix (Fig. 1b and c). Whereas, GmP mice revealed relatively healthy glomeruli evident from large podocytes, abundant mesangial cells and healthy mesangial matrix (Fig. 1d), scoring $3.8 \mu\text{m}^3$ glomerular area and (27 C/gcs) glomerular cells with insignificant difference

compared to control group and significant increase compared to gentamicin group (Table 2).

Control kidney sections stained with Masson's Trichrome showed abundant glomerular cells without any depositions of collagenous fibres inside glomeruli or in between cortical tubules (– to collagenous fibres) (Table 4, Fig. 2a). Whereas, kidney sections of Gm mice showed intense depositions of collagenous fibres and stained blue by Masson's Trichrome in the glomeruli and also in between cortical tubules (+++) (Table 4, Fig. 2b and c). Kidney sections of GmP mice show no collagenous fibres depositions in glomeruli or in between tubules (–) (Table 4, Fig. 2d).

Control kidney sections stained with Gomori silver technique showed no deposition of reticular fibres (–) (Fig. 3a). Whereas, kidney sections of Gm showed mild depositions of

**Fig. 1 – Glomerular analysis of kidney from control (a), Gm (b, c) and GmP (d) group of mice.**




Fig. 2 – Showing depositions of collagenous fibres in control (a), Gm (b and c), and GmP mice group.




Fig. 3 – Showing reticular fibres in control, Gm and GmP group of mice.

brown reticular fibres (++) in necrotic areas (Fig. 3b). While, kidney sections of GmP mice show no reticular fibres depositions (–) (Table 4, Fig. 3c).

Kidney sections stained by Avidin Biotin Complex (ABC) immunohistochemistry method for caspase-3 gene-expression show no immunoreactivity (–) in the kidney sections of control mice group (Fig. 4a) and in kidney sections from GmP mice (Fig. 4c). Whereas, kidney of Gm show intense brown immunoprecipitation (+++) inside the glomeruli (Fig. 4b, Table 4), indicating apoptosis.

Similarly, Kim-1 gene-expression shows no immunoreactivity (–) in control sections (Fig. 5a). Whereas, an intense immunoprecipitation was observed in glomeruli and cortical tubules in sections of Gm mice kidney (+++) (Fig. 5b). A slight ameliorative effect of propolis was evident from weak brownish immunoprecipitation observed in sections of GmP mice kidney (+) (Table 4, Fig. 5c). Kidney sections stained for Malondialdehyde (oxidative stress Marker) show no immunoprecipitation (–ve) in untreated control sections (Fig. 6a) almost similar immunoreaction was observed in GmP mice (Fig. 6c). Whereas, intense immunoprecipitation was observed

in glomeruli sections of Kidney from Gm mice group (+++) (Fig. 6b, Table 4).

Tubular analysis

Control kidney sections showed normal tubules without dilatation and proximal tubules appeared filled because of the long microvilli of the brush border and aggregates of small plasma proteins bound to this structure, by contrast lumens of distal tubules appeared empty (Fig. 7a). Sections of Gm mice kidney showed mild dilatation with a pathological score of 2 with empty lumens of proximal tubules score (3), moderate loss of pathological score (Fig. 7b). Whereas, sections of GmP mice scored 1, with mild injuries, dilatation and loss of brush borders (Table 3, Fig. 7c).

Control sections show (score 0) no leucocytic infiltration, tubular degeneration and tubular cast (Fig. 8a). While sections of Gm mice kidney show severe leucocytic infiltration (score 5, Fig. 8b), extensive tubular degeneration (score 4, Fig. 8c) and presence of tubular cast (Score 1, Fig. 8d, Table 3). Whereas, sections of GmP scored show mild leucocytic infiltration (Score




Fig. 4 – Caspase-3 gene-expression in control (a), Gm (b) and GmP (c) mice group.




Fig. 5 – Immunoreaction of Kim-1 gene in control (a), Gm (b) and GmP (c) treated mice.




Fig. 6 – Malondialdehyde immunoreaction in control (a), Gm (b) and GmP (c) treated mice.





Fig. 7 – Tubular analysis of control (a), Gm (b), GmP (c) and treated mice.

Table 3 – Pathological score of tubular injury in control, Gm and GmP experimental group of mice.

Parameters	Control	Gentamicin (Gm group)	Gentamicin + propolis (GmP group)
Dilated tubules	0	3 (± 0.1)	1 (± 0.1)
Loss of brush border	0	2 (± 0.3)	1 (± 0.1)
Leucocytic infiltration	0	5 (± 0.09)	1 (± 0.1)
Tubular degeneration	0	4 (± 0.1)	1 (± 0.1)
Tubular cast	0	1 (± 0.09)	0.4 (± 0.1)

The data presented in parenthesis are \pm SD (standard deviation).

**Fig. 8 – Leucocytic infiltration and tubular degeneration in control (a), gentamicin administered (Gm group; b, c) and GmP mice group.**

1), and tubular degeneration but do not show tubular cast (score 0, Fig. 8e).

Immunohistochemical analysis of control mice shows no immunoreactivity in control sections (–) for caspase 3 (Fig. 9a1). Whereas, mild (++) immunoprecipitates were seen in tubules kidney of Gm mice (Fig. 9a2). In GmP mice group however, there was a significant decrease in the intensity of immunoprecipitation (+) (Table 4, Fig. 9a3).

Kim-1 gene-expression also shows no immunoreactivity in control sections (–) (Fig. 9b1). Whereas, intense (+++) immunoprecipitation was observed in tubules of Gm mice kidney (Fig. 9b2). Moreover, in GmP mice (little, + Table 4) the intensity of Kim-1 gene immunoprecipitation was very low (Fig. 9b3). Kidney sections stained for Malondialdehyde (oxidative stress Marker) gene-expression showed no immunoreactivity in control sections (–) (Fig. 9c1). Whereas, intense (+++) brownish immunoprecipitates were seen in tubules of Gm mice kidney (Fig. 9c2), and very low intensity of (+, Table 4) immunoprecipitates was found in the tubules of Gmp mice kidney (Fig. 9c3).

Discussion

Results presented in this study confirmed that gentamicin administration caused marked changes in kidney tubules may be due to gentamicin reabsorption in proximal

convoluted tubules, causing degeneration and necrosis of the epithelial cells of the tubules. These changes are manifested by dilated tubules, loss of brush border, severe leucocytic infiltrations, tubular degeneration and presence of tubular casts. These findings are in agreement with previous studies.^{2,23,24} Co-administration of propolis with gentamicin revealed significant improvement in kidney tubules marked by the absence of tubular casts, reduction of infiltration, degeneration and tubular dilatation. Azab et al.²⁵ also reported similar effect of propolis, wherein co-administration of propolis with gentamicin, resulted in normal epithelial lining with brush borders in proximal convoluted tubules. However, some tubules appeared regenerating with disrupted brush borders.

Han et al.²⁶ has shown the activation of proapoptotic proteins in kidneys exhibiting nephrotoxicity. Caspases often used as a marker to study apoptosis, are from the family of endoproteases that provide critical links in cell regulatory networks controlling inflammation and cell death.²⁷ Sahu et al.²⁸ has shown that Gentamicin results in apoptosis in glomeruli and tubules. While, this toxicity was ameliorated by the co-administration of propolis. Renoprotective effect of Brazilian red propolis has also been demonstrated by Teles et al.²⁹ Other biomarkers to study nephrotoxicity include Kidney injury molecule 1. Prozialeck et al.³⁰ has suggested the use of KIM-1 as a nephrotoxicity biomarker in preclinical studies of drug candidates. Furthermore, Food and Drug




Fig. 9 – Immunohistochemical staining of, caspase 3 in control (a1), Gm (a2) and GmP (a3), Kim-1 control (b1), Gm (b2) and GmP (b3) and Malondialdehyde in control (c1), Gm (c2) and GmP (c3) group of mice.

Table 4 – Histochemical and immunohistochemical analysis in control, gentamicin (Gm), gentamicin treated with propolis (GmP) groups: –, means negative; +, little; ++, mild; +++, extensive.			
Parameters	Control	Gentamicin (Gm)	Gentamicin + propolis (GmP)
Collagenous fibres	–	+++	–
Reticular fibres	–	++	–
Caspase3 gene (glomeruli)	–	+++	–
Kim-1 gene (glomeruli)	–	+++	+
Malondialdehyde gene (glomeruli)	–	+++	–
Caspase3 gene (tubules)	–	++	+
Kim-1 gene (tubules)	–	+++	+
Malondialdehyde (tubules)	–	+++	+

Administration (USA) has also recently recognized KIM-1 as an appropriate biomarker for renal injury in preclinical studies of pharmacological agents. Besides being a sensitive diagnostic marker of nephrotoxicity, KIM-1 also has predictive value for AKI in patients undergoing cardiac surgery.³¹ Results obtained in our study confirmed that gentamicin administration produced severe kidney injury as evident from intense immunoreactions of kim-1 gene in glomeruli and tubules. These findings are in agreement with the reports of Chen et al.,³² Mcduffie et al.,³³ and Qiu et al.³⁴ As in these studies also an intense immunoreaction of Kim-1 was observed following exposure to gentamicin. Interestingly, a decrease in kim-1 immunoreaction was observed in this study when Gentamicin was co-administered with propolis; a trend which was also observed in caspase-3 immunoreactions.

Another mode through which gentamicin exert its nephrotoxicity, is through the generation of Reactive oxygen species (ROS) or oxidative stress.³⁵ These ROS target a number of biomolecules including lipids. Malondialdehyde (MDA) is the principal and most studied product of polyunsaturated fatty acid peroxidation. And hence is considered as an important marker of lipid peroxidation.³⁶ In agreement with previous studies,³⁷ gentamicin administration produced intense immunoreaction of (MDA) gene as an oxidative stress marker in glomeruli and tubules confirming the gentamicin mediated oxidative stress in kidney tissue. However, oral administration of propolis resulted in a decrease of MDA immunoprecipitation suggesting a decrease in oxidative stress. However, the pathway through which propolis result in this change is not known.

Based on the results presented in this study, it can be concluded that propolis is a good renoprotective agent and can effectively ameliorate the renotoxicity of gentamicin.

Q2 Conflicts of interest

The authors declare no conflicts of interest.

Q3 Uncited reference

22.

Acknowledgement

Thanks and sincere appreciation to the Deanship of Scientific Research at King Saud University for its funding this research group No. (RG-1435-030) and its perfect support for this project.

REFERENCES

1. Ali M, Goetz M. A meta-analysis of the relative efficacy and toxicity of single daily dosing versus multiple daily dosing of aminoglycosides. *Clin Infect Dis*. 1997;24:796–809.
2. Qadir M, Tahir M, Lone K, Munir B, Sam W. Protective role of ginseng against gentamicin induced changes in kidney of albino mice. *J Ayub Med Coll Abbottabad*. 2011;23:53–7.
3. Alarifi S, Al-Doaiss A, Alkahtani S, Al-Farraj S, Al-Eissa M, Al-Dahmash B, et al. Blood chemical changes and renal histological alterations induced by gentamicin in rats. *Saudi J Biol Sci*. 2012;19:103–10.
4. Rybak L, Whitworth C. Ototoxicity: therapeutic opportunities. *Drug Dis Today*. 2005;10:1313–21.
5. Aubrecht J, Goad M, Simpson E. Expression of hygR in transgenic mice causes resistance to toxic effects of hygromycin B in vivo. *J Pharmacol Exp Ther*. 1997;281:992–7.
6. Sanchez-Gonzalez PD, Lopez-Hernandez FJ, Perez-Barriocanal F, Morales AI, Lopez-Novoa JM. Quercetin reduces cisplatin nephrotoxicity in rats without compromising its anti-tumour activity. *Nephrol Dial Transplant*. 2011;26:3484–95.
7. Basile DP, Anderson MD, Sutton TA. Pathophysiology of acute kidney injury. *Comp Physiol*. 2012;2:1303–53.
8. Endre Z, Pickering J, Walker R, Devarajan P, Edelstein C, Bonventre J, et al. Improved performance of urinary biomarkers of acute kidney injury in the critically ill by stratification for injury duration and baseline renal function. *Kidney Int*. 2012;79:1119–30.
9. Havasi A, Borkan S. Apoptosis and acute kidney injury. *Kidney Int*. 2011;80:29–40.
10. Ichimura T, Hung C, Yang S, Stevens J, Bonventre J. Kidney injury molecule-1 (Kim-1): a tissue and urinary biomarker for nephrotoxicant-induced renal injury. *Am J Physiol Renal Physiol*. 2003;286:552–63.
11. Ali SS, Rizvi SZ, Muzaffar S, Ahmad A, Ali A, Hassan SH. Renal cortical necrosis: a case series of nine patients & review of literature. *J Ayub Med Coll Abbottabad*. 2003;15:41–4.
12. Cekmen M, Otunctemur A, Ozbek E, Cakir S, Dursun M, Polat E, et al. Pomegranate extract attenuates gentamicin-induced nephrotoxicity in rats by reducing oxidative stress. *Ren Fail*. 2013;2013:268–74.
13. Nagai J, Takano M. Molecular aspects of renal handling of aminoglycosides and strategies for preventing the nephrotoxicity. *Drug Metab Pharmacokin*. 2004;19:159–70.
14. Burdock GA. Review of the biological properties and toxicity of bee propolis (propolis). *Food Chem Toxicol*. 1998;36:347–63.
15. Chen C, Weng M, Wu C, Lin J. Comparison of radical scavenging activity, cytotoxic effects and apoptosis induction in human melanoma cells by Taiwanese propolis from different sources. *Evid Based Complement Alternat Med*. 2004;1:175–85.
16. Nassar S, Mohamed A, Soufy H, Nasr S, Mahran K. Immunostimulant effect of Egyptian propolis in rabbits. *Sci World J*. 2012;901516.
17. Nassar S, Mohamed A, Soufy H, Nasr S. Protective effect of Egyptian propolis against rabbit pasteurellosis. *Biomed Res Int*. 2013;2013:24.
18. Perron N, Brumaghim J. A review of the antioxidant mechanisms of polyphenol compounds related to iron binding. *Cell Biochem Biophys*. 2009;53:75–100.
19. El-Kott A, Owayss A. Protective effects of propolis against the amitraz hepatotoxicity in mice. *J Pharmacol Toxicol*. 2008;3:402–8.
20. Sun F, Hayami S, Haruna S, Ogiri Y, Tanaka K, Yamada Y, et al. In vivo antioxidative activity of propolis evaluated by the interaction with vitamins C and E and the level of lipid hydroperoxides in rats. *J Agric Food Chem*. 2000;48:1462.
21. Biswas M, Kar B, Karan TK, Bhattacharya S, Kumar SRB, Ghosh AK, et al. Acute and subchronic toxicity study of *Dregea volubilis* fruit in mice. *J Phytol*. 2010;2:6–10.
22. Noorani A, Gupta K, Bhadada K, Kale M. Protective effect of methanolic leaf extract of *Caesalpinia bonduc* (L.) on gentamicin-induced hepatotoxicity and nephrotoxicity in rats. *Iran J Pharmacol Ther*. 2011;10:21–5.
23. Nale L, More P, More B, Ghumare B, Shendre S, Mote C. Protective effect of *Carica papaya* L. seed extract in gentamicin induced hepatotoxicity and nephrotoxicity in rats. *Int J Pharm Biol Sci*. 2012;3:508–15.
24. Ullah N, Khan M, Khan T, Ahmad W. Protective effect of *Cinnamomum tamala* extract on gentamicin-induced nephrotic damage in rabbits. *Trop J Pharm Res*. 2013;12:215–9.
25. Azab A, Fetouh F, Albasha M. Nephro-protective effects of curcumin, rosemary and propolis against gentamicin induced toxicity in guinea pigs: morphological and biochemical study. *Am J Clin Exp Med*. 2014;2:28–35.
26. Han S, Chang E, Choi H, Kwak C, Park S, Kim H. Apoptosis by cyclosporine in mesangial cells. *Transplant Proc*. 2006;38:2244–6.
27. McIlwain D, Berger T, Mak T. Caspase functions in cell death and disease. *Cold Spring Harb Perspect Biol*. 2013;5:1–28.
28. Sahu BD, Tatireddy S, Koneru M, Borkar RM, Kumar JM, Kuncha M, et al. Naringin ameliorates gentamicin-induced nephrotoxicity and associated mitochondrial dysfunction, apoptosis and inflammation in rats: possible mechanism of nephroprotection. *Toxicol Appl Pharmacol*. 2014;277:8–20.
29. Teles F, da Silva TM, da Cruz Júnior FP, Honorato VH, de Oliveira Costa H, Barbosa AP, et al. Brazilian red propolis attenuates hypertension and renal damage in 5/6 renal ablation model. *PLOS ONE*. 2015;21:e0116535, <http://dx.doi.org/10.1371/journal.pone.0116535>.
30. Prozialeck WC, Vaidya VS, Liu J, Waalkes MP, Edwards JR, Lamar PC, et al. Kidney injury molecule-1 is an early biomarker of cadmium nephrotoxicity. *Kidney Int*. 2007;72:985–93.
31. Vaidya V, Ramirez V, Ichimura T, Bobadilla N, Bonventre J. Urinary kidney injury molecule-1: a sensitive quantitative biomarker for early detection of kidney tubular injury. *Am J Physiol Renal Physiol*. 2006;290:517–29.
32. Chen F, Smith R, Gu YZ, Collins ND, Nioi P. Toxicogenetic alteration of the kidney injury molecule 1 gene in gentamicin-exposed rat kidney. *Toxicol Sci*. 2010;117:375–80.

33. McDuffie J, Gao J, Ma J, La D, Bittner A, Sonee M, et al. Novel genomic biomarkers for acute gentamicin nephrotoxicity in dog. *J Mol Integr Physiol*. 2013;3:125–33.
34. Qiu Y, Hong M, Li H, Tang N, Ma J, Hsu C, et al. Time-series pattern of gene expression profile in gentamycin-induced nephrotoxicity. *Toxicol Mech Methods*. 2015;24:142–51.
35. Pedraza C, Maldonado P, Medina C. Garlic ameliorates gentamicin nephrotoxicity: relation antioxidant enzymes. *Free Radic Biol Med*. 2000;29:602–11.
36. Rio D, Stewart A, Pellegrini N. A review of recent studies on malondialdehyde as toxic molecule and biological marker of oxidative stress. *NMCD*. 2005;15:316–28.
37. Alqasoumi S. Protective effect of *Ipomea aquatica* forsk. On gentamicin-induced oxidative stress and nephropathy in rats. *Topclass J Herb Med*. 2013;2:13–9.