

# Introduction to Stochastic Processes

Lothar Breuer

# Contents

<b>1</b>	<b>Some general definitions</b>	<b>1</b>
<b>2</b>	<b>Markov Chains and Queues in Discrete Time</b>	<b>3</b>
2.1	Definition . . . . .	3
2.2	Classification of States . . . . .	8
2.3	Stationary Distributions . . . . .	13
2.4	Restricted Markov Chains . . . . .	20
2.5	Conditions for Positive Recurrence . . . . .	22
2.6	The M/M/1 queue in discrete time . . . . .	24
<b>3</b>	<b>Markov Processes on Discrete State Spaces</b>	<b>33</b>
3.1	Definition . . . . .	33
3.2	Stationary Distribution . . . . .	40
3.3	Skip-Free Markov Processes and Markovian Queues . . . . .	44
3.3.1	The M/M/1 Queue . . . . .	44
3.3.2	Skip-Free Markov Processes . . . . .	46
3.3.3	The M/M/ $\infty$ Queue . . . . .	48
3.3.4	The M/M/k Queue . . . . .	49
3.3.5	The M/M/k/k Queue . . . . .	51
3.3.6	The M/M/k/k+c/N Queue . . . . .	52
<b>4</b>	<b>Renewal Theory</b>	<b>57</b>
4.1	Renewal Processes . . . . .	57
4.2	Renewal Function and Renewal Equations . . . . .	60
4.3	Renewal Theorems . . . . .	62
4.4	Stationary Renewal Processes . . . . .	65

<b>5</b>	<b>Appendix</b>	<b>71</b>
5.1	Conditional Expectations and Probabilities . . . . .	71
5.2	Extension Theorems . . . . .	74
5.2.1	Stochastic chains . . . . .	74
5.2.2	Stochastic processes . . . . .	75
5.3	Transforms . . . . .	76
5.3.1	$z$ -transforms . . . . .	76
5.3.2	Laplace–Stieltjes transforms . . . . .	78
5.4	Gershgorin’s circle theorem . . . . .	79



# **Chapter 1**

## **Some general definitions**

see notes under

<http://www.kent.ac.uk/IMS/personal/lb209/files/notes1.pdf>



## Chapter 2

# Markov Chains and Queues in Discrete Time

### 2.1 Definition

Let  $X_n$  with  $n \in \mathbb{N}_0$  denote random variables on a discrete space  $E$ . The sequence  $\mathcal{X} = (X_n : n \in \mathbb{N}_0)$  is called a **stochastic chain**. If  $\mathbb{P}$  is a probability measure  $\mathcal{X}$  such that

$$\mathbb{P}(X_{n+1} = j | X_0 = i_0, \dots, X_n = i_n) = \mathbb{P}(X_{n+1} = j | X_n = i_n) \quad (2.1)$$

for all  $i_0, \dots, i_n, j \in E$  and  $n \in \mathbb{N}_0$ , then the sequence  $\mathcal{X}$  shall be called a **Markov chain** on  $E$ . The probability measure  $\mathbb{P}$  is called the distribution of  $\mathcal{X}$ , and  $E$  is called the **state space** of  $\mathcal{X}$ .

If the conditional probabilities  $\mathbb{P}(X_{n+1} = j | X_n = i_n)$  are independent of the time index  $n \in \mathbb{N}_0$ , then we call the Markov chain  $\mathcal{X}$  **homogeneous** and denote

$$p_{ij} := \mathbb{P}(X_{n+1} = j | X_n = i)$$

for all  $i, j \in E$ . The probability  $p_{ij}$  is called **transition probability** from state  $i$  to state  $j$ . The matrix  $P := (p_{ij})_{i,j \in E}$  shall be called **transition matrix** of the chain  $\mathcal{X}$ . Condition (2.1) is referred to as the **Markov property**.

**Example 2.1** If  $(X_n : n \in \mathbb{N}_0)$  are random variables on a discrete space  $E$ , which are stochastically independent and identically distributed (shortly: iid), then the chain  $\mathcal{X} = (X_n : n \in \mathbb{N}_0)$  is a homogeneous Markov chain.

**Example 2.2** Discrete Random Walk

Set  $E := \mathbb{Z}$  and let  $(S_n : n \in \mathbb{N})$  be a sequence of iid random variables with values in  $\mathbb{Z}$  and distribution  $\pi$ . Define  $X_0 := 0$  and  $X_n := \sum_{k=1}^n S_k$  for all  $n \in \mathbb{N}$ . Then the chain  $\mathcal{X} = (X_n : n \in \mathbb{N}_0)$  is a homogeneous Markov chain with transition probabilities  $p_{ij} = \pi_{j-i}$ . This chain is called **discrete random walk**.

**Example 2.3** Bernoulli process

Set  $E := \mathbb{N}_0$  and choose any parameter  $0 < p < 1$ . The definitions  $X_0 := 0$  as well as

$$p_{ij} := \begin{cases} p, & j = i + 1 \\ 1 - p, & j = i \end{cases}$$

for  $i \in \mathbb{N}_0$  determine a homogeneous Markov chain  $\mathcal{X} = (X_n : n \in \mathbb{N}_0)$ . It is called **Bernoulli process** with parameter  $p$ .

So far, all examples have been chosen as to be homogeneous. The following theorem shows that there is a good reason for this:

**Theorem 2.4** *Be  $\mathcal{X} = (X_n : n \in \mathbb{N}_0)$  a Markov chain on a discrete state space  $E$ . Then there is a homogeneous Markov chain  $\mathcal{X}' = (X'_n : n \in \mathbb{N}_0)$  on the state space  $E \times \mathbb{N}_0$  such that  $X_n = pr_1(X'_n)$  for all  $n \in \mathbb{N}_0$ , with  $pr_1$  denoting the projection to the first dimension.*

**Proof:** Let  $\mathcal{X}$  be a Markov chain with transition probabilities

$$p_{n;ij} := \mathbb{P}(X_{n+1} = j | X_n = i)$$

which may depend on the time instant  $n$ . Define the two-dimensional random variables  $X'_n := (X_n, n)$  for all  $n \in \mathbb{N}_0$  and denote the resulting distribution of the chain  $\mathcal{X}' = (X'_n : n \in \mathbb{N}_0)$  by  $\mathbb{P}'$ . By definition we obtain  $X_n = pr_1(X'_n)$  for all  $n \in \mathbb{N}_0$ .

Further  $\mathbb{P}'(X'_0 = (i, k)) = \delta_{k0} \cdot \mathbb{P}(X_0 = i)$  holds for all  $i \in E$ , and all transition probabilities

$$p'_{(i,k),(j,l)} = \mathbb{P}'(X'_{k+1} = (j, l) | X'_k = (i, k)) = \delta_{l,k+1} \cdot p_{k;ij}$$

can be expressed without a time index. Hence the Markov chain  $\mathcal{X}'$  is homogeneous.

□

Because of this result, we will from now on treat only homogeneous Markov chains and omit the adjective "homogeneous".



Let  $P$  denote the transition matrix of a Markov chain on  $E$ . Then as an immediate consequence of its definition we obtain  $p_{ij} \in [0, 1]$  for all  $i, j \in E$  and  $\sum_{j \in E} p_{ij} = 1$  for all  $i \in E$ . A matrix  $P$  with these properties is called a **stochastic matrix** on  $E$ . In the following we shall demonstrate that, given an initial distribution, a Markov chain is uniquely determined by its transition matrix. Thus any stochastic matrix defines a family of Markov chains.

**Theorem 2.5** *Let  $\mathcal{X}$  denote a homogeneous Markov chain on  $E$  with transition matrix  $P$ . Then the relation*

$$\mathbb{P}(X_{n+1} = j_1, \dots, X_{n+m} = j_m | X_n = i) = p_{i,j_1} \cdot \dots \cdot p_{j_{m-1},j_m}$$

*holds for all  $n \in \mathbb{N}_0$ ,  $m \in \mathbb{N}$ , and  $i, j_1, \dots, j_m \in E$ .*

**Proof:** This is easily shown by induction on  $m$ . For  $m = 1$  the statement holds by definition of  $P$ . For  $m > 1$  we can write

$$\begin{aligned} & \mathbb{P}(X_{n+1} = j_1, \dots, X_{n+m} = j_m | X_n = i) \\ &= \frac{\mathbb{P}(X_{n+1} = j_1, \dots, X_{n+m} = j_m, X_n = i)}{\mathbb{P}(X_n = i)} \\ &= \frac{\mathbb{P}(X_{n+1} = j_1, \dots, X_{n+m-1} = j_{m-1}, X_n = i)}{\mathbb{P}(X_{n+1} = j_1, \dots, X_{n+m-1} = j_{m-1}, X_n = i)} \\ & \quad \times \frac{\mathbb{P}(X_{n+1} = j_1, \dots, X_{n+m-1} = j_{m-1}, X_n = i)}{\mathbb{P}(X_n = i)} \\ &= \mathbb{P}(X_{n+m} = j_m | X_n = i, X_{n+1} = j_1, \dots, X_{n+m-1} = j_{m-1}) \\ & \quad \times p_{i,j_1} \cdot \dots \cdot p_{j_{m-2},j_{m-1}} \\ &= p_{j_{m-1},j_m} \cdot p_{i,j_1} \cdot \dots \cdot p_{j_{m-2},j_{m-1}} \end{aligned}$$

because of the induction hypothesis and the Markov property.

□

Let  $\pi$  be a probability distribution on  $E$  with  $\mathbb{P}(X_0 = i) = \pi_i$  for all  $i \in E$ . Then theorem 2.5 immediately yields

$$\mathbb{P}(X_0 = j_0, X_1 = j_1, \dots, X_m = j_m) = \pi_{j_0} \cdot p_{j_0,j_1} \cdot \dots \cdot p_{j_{m-1},j_m} \quad (2.2)$$

for all  $m \in \mathbb{N}$  and  $j_0, \dots, j_m \in E$ . The chain with this distribution  $\mathbb{P}$  is denoted by  $\mathcal{X}^\pi$  and called the  $\pi$ -**version** of  $\mathcal{X}$ . The probability measure  $\pi$  is called **initial distribution** for  $\mathcal{X}$ .

Theorem 2.5 and the extension theorem by Tulcea (see appendix 5.2) show that a Markov chain is uniquely determined by its transition matrix and its initial distribution. Whenever the initial distribution  $\pi$  is not important or understood from the context, we will simply write  $\mathcal{X}$  instead of  $\mathcal{X}^\pi$ . However, in an exact manner the notation  $\mathcal{X}$  denotes the family of all the versions  $\mathcal{X}^\pi$  of  $\mathcal{X}$ , indexed by their initial distribution  $\pi$ .

**Theorem 2.6** *Let  $\mathcal{X}$  denote a homogeneous Markov chain with transition matrix  $P$ . Then the relation*

$$\mathbb{P}(X_{n+m} = j | X_n = i) = P^m(i, j)$$

*holds for all  $m, n \in \mathbb{N}_0$  and  $i, j \in E$ , with  $P^m(i, j)$  denoting the  $(i, j)$ th entry of the  $m$ th power of the matrix  $P$ . In particular,  $P^0$  equals the identity matrix.*

**Proof:** This follows by induction on  $m$ . For  $m = 1$  the statement holds by definition of  $P$ . For  $m > 1$  we can write

$$\begin{aligned} \mathbb{P}(X_{n+m} = j | X_n = i) &= \frac{\mathbb{P}(X_{n+m} = j, X_n = i)}{\mathbb{P}(X_n = i)} \\ &= \sum_{k \in E} \frac{\mathbb{P}(X_{n+m} = j, X_{n+m-1} = k, X_n = i)}{\mathbb{P}(X_{n+m-1} = k, X_n = i)} \\ &\quad \times \frac{\mathbb{P}(X_{n+m-1} = k, X_n = i)}{\mathbb{P}(X_n = i)} \\ &= \sum_{k \in E} \mathbb{P}(X_{n+m} = j | X_{n+m-1} = k, X_n = i) \cdot P^{m-1}(i, k) \\ &= \sum_{k \in E} p_{kj} \cdot P^{m-1}(i, k) = P^m(i, j) \end{aligned}$$

because of the induction hypothesis and the Markov property.

□

Thus the probabilities for transitions in  $m$  steps are given by the  $m$ th power of the transition matrix  $P$ . The rule  $P^{m+n} = P^m P^n$  for the multiplication of matrices and theorem 2.6 lead to the decompositions

$$\mathbb{P}(X_{m+n} = j | X_0 = i) = \sum_{k \in E} \mathbb{P}(X_m = k | X_0 = i) \cdot \mathbb{P}(X_n = j | X_0 = k)$$

which are known as the **Chapman–Kolmogorov equations**.

For later purposes we will need a relation closely related to the Markov property, which is called the **strong Markov property**. Let  $\tau$  denote a random variable with values in  $\mathbb{N}_0 \cup \{\infty\}$ , such that the condition

$$\mathbb{P}(\tau \leq n | \mathcal{X}) = \mathbb{P}(\tau \leq n | X_0, \dots, X_n) \quad (2.3)$$

holds for all  $n \in \mathbb{N}_0$ . Such a random variable is called a (discrete) **stopping time** for  $\mathcal{X}$ . The defining condition means that the probability for the event  $\{\tau \leq n\}$  depends only on the evolution of the chain until time  $n$ . In other words, the determination of a stopping time does not require any knowledge of the future. Now the strong Markov property is stated in

**Theorem 2.7** *Let  $\mathcal{X}$  denote a Markov chain and  $\tau$  a stopping time for  $\mathcal{X}$  with  $\mathbb{P}(\tau < \infty) = 1$ . Then the relation*

$$\mathbb{P}(X_{\tau+m} = j | X_0 = i_0, \dots, X_\tau = i_\tau) = \mathbb{P}(X_m = j | X_0 = i_\tau)$$

*holds for all  $m \in \mathbb{N}$  and  $i_0, \dots, i_\tau, j \in E$ .*

**Proof:** The fact that the stopping time  $\tau$  is finite and may assume only countably many values can be exploited in the transformation

$$\begin{aligned} \mathbb{P}(X_{\tau+m} = j | X_0 = i_0, \dots, X_\tau = i_\tau) &= \sum_{n=0}^{\infty} \mathbb{P}(\tau = n, X_{\tau+m} = j | X_0 = i_0, \dots, X_\tau = i_\tau) \\ &= \sum_{n=0}^{\infty} \mathbb{P}(X_{\tau+m} = j | \tau = n, X_0 = i_0, \dots, X_\tau = i_\tau) \\ &\quad \times \mathbb{P}(\tau = n | X_0 = i_0, \dots, X_\tau = i_\tau) \\ &= \sum_{n=0}^{\infty} \mathbb{P}(X_{n+m} = j | X_n = i_\tau) \cdot \mathbb{P}(\tau = n | \mathcal{X}) \\ &= \sum_{n=0}^{\infty} \mathbb{P}(\tau = n | \mathcal{X}) \cdot \mathbb{P}(X_m = j | X_0 = i_\tau) \end{aligned}$$

which yields the statement, as  $\tau$  is finite with probability one.

□

## 2.2 Classification of States

Let  $\mathcal{X}$  denote a Markov chain with state space  $E$  and transition matrix  $P$ . We call a state  $j \in E$  **accessible** from a state  $i \in E$  if there is a number  $m \in \mathbb{N}_0$  with  $P(X_m = j | X_0 = i) > 0$ . This relation shall be denoted by  $i \rightarrow j$ . If for two states  $i, j \in E$ , the relations  $i \rightarrow j$  and  $j \rightarrow i$  hold, then  $i$  and  $j$  are said to **communicate**, in notation  $i \leftrightarrow j$ .

**Theorem 2.8** *The relation  $\leftrightarrow$  of communication between states is an equivalence relation.*

**Proof:** Because of  $P^0 = I$ , communication is reflexive. Symmetry holds by definition. Thus it remains to show transitivity. For this, assume  $i \leftrightarrow j$  and  $j \leftrightarrow k$  for three states  $i, j, k \in E$ . This means that there are numbers  $m, n \in \mathbb{N}_0$  with  $P^m(i, j) > 0$  and  $P^n(j, k) > 0$ . Hence, by the Chapman–Kolmogorov equation, we obtain

$$\begin{aligned} \mathbb{P}(X_{m+n} = k | X_0 = i) &= \sum_{h \in E} \mathbb{P}(X_m = h | X_0 = i) \cdot \mathbb{P}(X_n = k | X_0 = h) \\ &\geq \mathbb{P}(X_m = j | X_0 = i) \cdot \mathbb{P}(X_n = k | X_0 = j) > 0 \end{aligned}$$

which proves  $i \rightarrow k$ . The remaining proof of  $k \rightarrow i$  is completely analogous.

□

Because of this result and the countability, we can divide the state space  $E$  of a Markov chain into a partition of countably many equivalence classes with respect to the communication of states. Any such equivalence class shall be called **communication class**. A communication class  $C \subset E$  that does not allow access to states outside itself, i.e. for which the implication

$$i \rightarrow j, \quad i \in C \quad \Rightarrow \quad j \in C$$

holds, is called **closed**. If a closed equivalence class consists only of one state, then this state shall be called **absorbing**. If a Markov chain has only one communication class, i.e. if all states are communicating, then it is called **irreducible**. Otherwise it is called **reducible**.

**Example 2.9** Let  $\mathcal{X}$  denote a discrete random walk (see example 2.2) with the specification  $\pi_1 = p$  and  $\pi_{-1} = 1 - p$  for some parameter  $0 < p < 1$ . Then  $\mathcal{X}$  is irreducible.

**Example 2.10** The Bernoulli process (see example 2.3) with non-trivial parameter  $0 < p < 1$  is to the highest degree reducible. Every state  $x \in \mathbb{N}_0$  forms an own communication class. None of these is closed, thus there are no absorbing states.

**Theorem 2.11** *Be  $\mathcal{X}$  a Markov chain with state space  $E$  and transition matrix  $P$ . Let  $C = \{c_n : n \in I\} \subset E$  with  $I \subset \mathbb{N}$  be a closed communication class. Define the matrix  $P'$  by its entries  $p'_{ij} := p_{c_i, c_j}$  for all  $i, j \in I$ . Then  $P'$  is stochastic.*

**Proof:** By definition,  $p'_{ij} \in [0, 1]$  for all  $i, j \in I$ . Since  $C$  is closed,  $p_{c_i, k} = 0$  for all  $i \in I$  and  $k \notin C$ . This implies

$$\sum_{j \in I} p'_{ij} = \sum_{j \in I} p_{c_i, c_j} = 1 - \sum_{k \notin C} p_{c_i, k} = 1$$

for all  $i \in I$ , as  $P$  is stochastic.

□

Thus the restriction of a Markov chain  $\mathcal{X}$  with state space  $E$  to the states of one of its closed communication classes  $C$  defines a new Markov chain with state space  $C$ . If the states are relabeled according to their affiliation to a communication class, the transition matrix of  $\mathcal{X}$  can be displayed in a block matrix form as

$$P = \begin{bmatrix} Q & Q_1 & Q_2 & Q_3 & Q_4 & \dots \\ 0 & P_1 & 0 & 0 & 0 & \dots \\ 0 & 0 & P_2 & 0 & 0 & \dots \\ 0 & 0 & 0 & P_3 & 0 & \dots \\ \vdots & \vdots & \ddots & \ddots & \ddots & \ddots \end{bmatrix} \quad (2.4)$$

with  $P_n$  being stochastic matrices on the closed communication classes  $C_n$ . The first row contains the transition probabilities starting from communication classes that are not closed.

Let  $\mathcal{X}$  denote a Markov chain with state space  $E$ . In the rest of this section we shall investigate distribution and expectation of the following random variables: Define  $\tau_j$  as the stopping time of the **first visit** to the state  $j \in E$ , i.e.

$$\tau_j := \min\{n \in \mathbb{N} : X_n = j\}$$

Denote the distribution of  $\tau_j$  by

$$F_k(i, j) := \mathbb{P}(\tau_j = k | X_0 = i)$$

for all  $i, j \in E$  and  $k \in \mathbb{N}$ .

**Lemma 2.12** *The conditional distribution of the first visit to the state  $j \in E$ , given an initial state  $X_0 = i$ , can be determined iteratively by*

$$F_k(i, j) = \begin{cases} p_{ij}, & k = 1 \\ \sum_{h \neq j} p_{ih} F_{k-1}(h, j), & k \geq 2 \end{cases}$$

for all  $i, j \in E$ .

**Proof:** For  $k = 1$ , the definition yields

$$F_1(i, j) = \mathbb{P}(\tau_j = 1 | X_0 = i) = \mathbb{P}(X_1 = j | X_0 = i) = p_{ij}$$

for all  $i, j \in E$ . For  $k \geq 2$ , conditioning upon  $X_1$  yields

$$\begin{aligned} F_k(i, j) &= \mathbb{P}(X_1 \neq j, \dots, X_{k-1} \neq j, X_k = j | X_0 = i) \\ &= \sum_{h \neq j} \mathbb{P}(X_1 = h | X_0 = i) \\ &\quad \times \mathbb{P}(X_2 \neq j, \dots, X_{k-1} \neq j, X_k = j | X_0 = i, X_1 = h) \\ &= \sum_{h \neq j} p_{ih} \cdot \mathbb{P}(X_1 \neq j, \dots, X_{k-2} \neq j, X_{k-1} = j | X_0 = h) \end{aligned}$$

due to the Markov property.

□

Now define

$$f_{ij} := \mathbb{P}(\tau_j < \infty | X_0 = i) = \sum_{k=1}^{\infty} F_k(i, j) \quad (2.5)$$

for all  $i, j \in E$ , which represents the probability of ever visiting state  $j$  after beginning in state  $i$ . Summing up over all  $k \in \mathbb{N}$  in the formula of Lemma 2.12 leads to

$$f_{ij} = p_{ij} + \sum_{h \neq j} p_{ih} f_{hj} \quad (2.6)$$

for all  $i, j \in E$ . The proof is left as an exercise.

Define  $N_j$  as the random variable of the **total number of visits** to the state  $j \in E$ . Expression (2.6) is useful for computing the distribution of  $N_j$ :

**Theorem 2.13** *Let  $\mathcal{X}$  denote a Markov chain with state space  $E$ . The total number of visits to a state  $j \in E$  under the condition that the chain starts in state  $i$  is given by*

$$\mathbb{P}(N_j = m | X_0 = j) = f_{jj}^{m-1}(1 - f_{jj})$$

and for  $i \neq j$

$$\mathbb{P}(N_j = m | X_0 = i) = \begin{cases} 1 - f_{ij}, & m = 0 \\ f_{ij}f_{jj}^{m-1}(1 - f_{jj}), & m \geq 1 \end{cases}$$

Thus the distribution of  $N_j$  is modified geometric.

**Proof:** Define  $\tau_j^{(1)} := \tau_j$  and  $\tau_j^{(k+1)} := \min\{n > \tau_j^{(k)} : X_n = j\}$  for all  $k \in \mathbb{N}$ , with the convention that  $\min \emptyset = \infty$ . Note that  $\tau_j^{(k)} = \infty$  implies  $\tau_j^{(l)} = \infty$  for all  $l > k$ .

Then the sequence  $(\tau_j^{(k)} : k \in \mathbb{N})$  is a sequence of stopping times. The event  $\{N_j = m\}$  is the same as the intersection of the events  $\{\tau_j^{(k)} < \infty\}$  for  $k = 1, \dots, M$  and  $\{\tau_j^{(M+1)} = \infty\}$ , with  $M = m$  if  $i \neq j$  and  $M = m-1$  if  $i = j$ . Now this event can be further described by the intersection of the events  $\{\tau_j^{(k+1)} - \tau_j^{(k)} < \infty\}$  for  $k = 0, \dots, M-1$  and  $\{\tau_j^{(M+1)} - \tau_j^{(M)} = \infty\}$ , with  $M$  as above and the convention  $\tau_j^{(0)} := 0$ .

The subevent  $\{\tau_j^{(k+1)} - \tau_j^{(k)} < \infty\}$  has probability  $f_{ij}$  for  $k = 0$  and because of the strong Markov property (see theorem 2.7) probability  $f_{jj}$  for  $k > 0$ . The probability for  $\{\tau_j^{(M+1)} - \tau_j^{(M)} = \infty\}$  is  $1 - f_{ij}$  for  $M = 0$  and  $1 - f_{jj}$  for  $M > 0$ . Once more the strong Markov property is the reason for independence of the subevents. Now multiplication of the probabilities leads to the formulae in the statement.

□

Summing over all  $m$  in the above theorem leads to

**Corollary 2.14** *For all  $j \in E$ , the zero-one law*

$$\mathbb{P}(N_j < \infty | X_0 = j) = \begin{cases} 1, & f_{jj} < 1 \\ 0, & f_{jj} = 1 \end{cases}$$

holds, i.e. depending on  $f_{jj}$  there are almost certainly infinitely many visits to a state  $j \in E$ .

This result gives rise to the following definitions: A state  $j \in E$  is called **re-current** if  $f_{jj} = 1$  and **transient** otherwise. Let us further define the **potential matrix**  $R = (r_{ij})_{i,j \in E}$  of the Markov chain by its entries

$$r_{ij} := \mathbb{E}(N_j | X_0 = i)$$

for all  $i, j \in E$ . Thus an entry  $r_{ij}$  gives the expected number of visits to the state  $j \in E$  under the condition that the chain starts in state  $i \in E$ . As such,  $r_{ij}$  can be computed by

$$r_{ij} = \sum_{n=0}^{\infty} P^n(i, j) \quad (2.7)$$

for all  $i, j \in E$ . The results in theorem 2.13 and corollary 2.14 yield

**Corollary 2.15** *For all  $i, j \in E$  the relations*

$$r_{jj} = (1 - f_{jj})^{-1} \quad \text{and} \quad r_{ij} = f_{ij} r_{jj}$$

*hold, with the conventions  $0^{-1} := \infty$  and  $0 \cdot \infty := 0$  included. In particular, the expected number  $r_{jj}$  of visits to the state  $j \in E$  is finite if  $j$  is transient and infinite if  $j$  is recurrent.*

**Theorem 2.16** *Recurrence and transience of states are class properties with respect to the relation  $\leftrightarrow$ . Furthermore, a recurrent communication class is always closed.*

**Proof:** Assume that  $i \in E$  is transient and  $i \leftrightarrow j$ . Then there are numbers  $m, n \in \mathbb{N}$  with  $0 < P^m(i, j) \leq 1$  and  $0 < P^n(j, i) \leq 1$ . The inequalities

$$\sum_{k=0}^{\infty} P^k(i, i) \geq \sum_{h=0}^{\infty} P^{m+h+n}(i, i) \geq P^m(i, j) P^n(j, i) \sum_{k=0}^{\infty} P^k(j, j)$$

now imply  $r_{jj} < \infty$  because of representation (2.7). According to corollary 2.15 this means that  $j$  is transient, too.

If  $j$  is recurrent, then the same inequalities lead to

$$r_{ii} \geq P^m(i, j) P^n(j, i) r_{jj} = \infty$$

which signifies that  $i$  is recurrent, too. Since the above arguments are symmetric in  $i$  and  $j$ , the proof of the first statement is complete.



For the second statement assume that  $i \in E$  belongs to a communication class  $C \subset E$  and  $p_{ij} > 0$  for some state  $j \in E \setminus C$ . Then

$$f_{ii} = p_{ii} + \sum_{h \neq i} p_{ih} f_{hi} \leq 1 - p_{ij} < 1$$

according to formula (2.6), since  $f_{ji} = 0$  (otherwise  $i \leftrightarrow j$ ). Thus  $i$  is transient, which proves the second statement.

□

**Theorem 2.17** *If the state  $j \in E$  is transient, then  $\lim_{n \rightarrow \infty} P^n(i, j) = 0$ , regardless of the initial state  $i \in E$ .*

**Proof:** If the state  $j$  is transient, then the first equation in corollary 2.15 yields  $r_{jj} < \infty$ . The second equation in the same corollary now implies  $r_{ij} < \infty$ , which by the representation (2.7) completes the proof.

□

## 2.3 Stationary Distributions

Let  $\mathcal{X}$  denote a Markov chain with state space  $E$  and  $\pi$  a measure on  $E$ . If  $\mathbb{P}(X_n = i) = \mathbb{P}(X_0 = i) = \pi_i$  for all  $n \in \mathbb{N}$  and  $i \in E$ , then  $\mathcal{X}^\pi$  is called **stationary**, and  $\pi$  is called a **stationary measure** for  $\mathcal{X}$ . If furthermore  $\pi$  is a probability measure, then it is called **stationary distribution** for  $\mathcal{X}$ .

**Theorem 2.18** *Let  $\mathcal{X}$  denote a Markov chain with state space  $E$  and transition matrix  $P$ . Further, let  $\pi$  denote a probability distribution on  $E$  with  $\pi P = \pi$ , i.e.*

$$\pi_i = \sum_{j \in E} \pi_j p_{ji} \quad \text{and} \quad \sum_{j \in E} \pi_j = 1$$

*for all  $i \in E$ . Then  $\pi$  is a stationary distribution for  $\mathcal{X}$ . If  $\pi$  is a stationary distribution for  $\mathcal{X}$ , then  $\pi P = \pi$  holds.*

**Proof:** Let  $\mathbb{P}(X_0 = i) = \pi_i$  for all  $i \in E$ . Then  $\mathbb{P}(X_n = i) = \mathbb{P}(X_0 = i)$  for all  $n \in \mathbb{N}$  and  $i \in E$  follows by induction on  $n$ . The case  $n = 1$  holds by assumption, and the induction step follows by induction hypothesis and the Markov property. The last statement is obvious.

□

The following examples show some features of stationary distributions:

**Example 2.19** Let the transition matrix of a Markov chain  $\mathcal{X}$  be given by

$$P = \begin{pmatrix} 0.8 & 0.2 & 0 & 0 \\ 0.2 & 0.8 & 0 & 0 \\ 0 & 0 & 0.4 & 0.6 \\ 0 & 0 & 0.6 & 0.4 \end{pmatrix}$$

Then  $\pi = (0.5, 0.5, 0, 0)$ ,  $\pi' = (0, 0, 0.5, 0.5)$  as well as any linear combination of them are stationary distributions for  $\mathcal{X}$ . This shows that a stationary distribution does not need to be unique.

**Example 2.20** Bernoulli process (see example 2.1)

The transition matrix of a Bernoulli process has the structure

$$P = \begin{pmatrix} 1-p & p & 0 & 0 & \dots \\ 0 & 1-p & p & 0 & \ddots \\ 0 & 0 & 1-p & p & \ddots \\ \vdots & \ddots & \ddots & \ddots & \ddots \end{pmatrix}$$

Hence  $\pi P = \pi$  implies first

$$\pi_0 \cdot (1-p) = \pi_0 \quad \Rightarrow \quad \pi_0 = 0$$

since  $0 < p < 1$ . Assume that  $\pi_n = 0$  for any  $n \in \mathbb{N}_0$ . This and the condition  $\pi P = \pi$  further imply for  $\pi_{n+1}$

$$\pi_n \cdot p + \pi_{n+1} \cdot (1-p) = \pi_{n+1} \quad \Rightarrow \quad \pi_{n+1} = 0$$

which completes an induction argument proving  $\pi_n = 0$  for all  $n \in \mathbb{N}_0$ . Hence the Bernoulli process does not have a stationary distribution.

**Example 2.21** The solution of  $\pi P = \pi$  and  $\sum_{j \in E} \pi_j = 1$  is unique for

$$P = \begin{pmatrix} 1-p & p \\ p & 1-p \end{pmatrix}$$

with  $0 < p < 1$ . Thus there are transition matrices which have exactly one stationary distribution.

The question of existence and uniqueness of a stationary distribution is one of the most important problems in the theory of Markov chains. A simple answer can be given in the transient case (cf. example 2.20):

**Theorem 2.22** *A transient Markov chain (i.e. a Markov chain with transient states only) has no stationary distribution.*

**Proof:** Assume that  $\pi P = \pi$  holds for some distribution  $\pi$  and take any enumeration  $E = (s_n : n \in \mathbb{N})$  of the state space  $E$ . Choose any index  $m \in \mathbb{N}$  with  $\pi_{s_m} > 0$ . Since  $\sum_{n=1}^{\infty} \pi_{s_n} = 1$  is bounded, there is an index  $M > m$  such that  $\sum_{n=M}^{\infty} \pi_{s_n} < \pi_{s_m}$ . Set  $\varepsilon := \pi_{s_m} - \sum_{n=M}^{\infty} \pi_{s_n}$ . According to theorem 2.17, there is an index  $N \in \mathbb{N}$  such that  $P^n(s_i, s_m) < \varepsilon$  for all  $i \leq M$  and  $n \geq N$ . Then the stationarity of  $\pi$  implies

$$\begin{aligned} \pi_{s_m} &= \sum_{i=1}^{\infty} \pi_{s_i} P^N(s_i, s_m) = \sum_{i=1}^{M-1} \pi_{s_i} P^N(s_i, s_m) + \sum_{i=M}^{\infty} \pi_{s_i} P^N(s_i, s_m) \\ &< \varepsilon + \sum_{i=M}^{\infty} \pi_{s_i} = \pi_{s_m} \end{aligned}$$

which is a contradiction.

□

For the recurrent case, a finer distinction will be necessary. While the expected total number  $r_{jj}$  of visits to a recurrent state  $j \in E$  is always infinite (see corollary 2.15), there are differences in the rate of visits to a recurrent state. In order to describe these, define  $N_i(n)$  as the number of visits to state  $i$  until time  $n$ . Further define for a recurrent state  $i \in E$  the mean time

$$m_i := \mathbb{E}(\tau_i | X_0 = i)$$

until the first visit to  $i$  (after time zero) under the condition that the chain starts in  $i$ . By definition  $m_i > 0$  for all  $i \in E$ . The elementary renewal theorem (which will be proven later as theorem 4.12) states that

$$\lim_{n \rightarrow \infty} \frac{\mathbb{E}(N_i(n) | X_0 = j)}{n} = \frac{1}{m_i} \quad (2.8)$$

for all recurrent  $i \in E$  and independently of  $j \in E$  provided  $j \leftrightarrow i$ , with the convention of  $1/\infty := 0$ . Thus the asymptotic rate of visits to a recurrent state is determined by the mean recurrence time of this state. This gives reason to the

following definition: A recurrent state  $i \in E$  with  $m_i = \mathbb{E}(\tau_i | X_0 = i) < \infty$  will be called **positive recurrent**, otherwise  $i$  is called **null recurrent**. The distinction between positive and null recurrence is supported by the equivalence relation  $\leftrightarrow$ , as shown in

**Theorem 2.23** *Positive recurrence and null recurrence are class properties with respect to the relation of communication between states.*

**Proof:** Assume that  $i \leftrightarrow j$  for two states  $i, j \in E$  and  $i$  is null recurrent. Thus there are numbers  $m, n \in \mathbb{N}$  with  $P^n(i, j) > 0$  and  $P^m(j, i) > 0$ . Because of the representation  $\mathbb{E}(N_i(k) | X_0 = i) = \sum_{l=0}^k P^l(i, i)$ , we obtain

$$\begin{aligned} 0 &= \lim_{k \rightarrow \infty} \frac{\sum_{l=0}^k P^l(i, i)}{k} \\ &\geq \lim_{k \rightarrow \infty} \frac{\sum_{l=0}^{k-m-n} P^l(j, j)}{k} \cdot P^n(i, j) P^m(j, i) \\ &= \lim_{k \rightarrow \infty} \frac{k-m-n}{k} \cdot \frac{\sum_{l=0}^{k-m-n} P^l(j, j)}{k-m-n} \cdot P^n(i, j) P^m(j, i) \\ &= \lim_{k \rightarrow \infty} \frac{\sum_{l=0}^k P^l(j, j)}{k} \cdot P^n(i, j) P^m(j, i) \\ &= \frac{P^n(i, j) P^m(j, i)}{m_j} \end{aligned}$$

and thus  $m_j = \infty$ , which signifies the null recurrence of  $j$ .

□

Thus we can call a communication class positive recurrent or null recurrent. In the former case, a construction of a stationary distribution is given in

**Theorem 2.24** *Let  $i \in E$  be positive recurrent and define the mean first visit time  $m_i := \mathbb{E}(\tau_i | X_0 = i)$ . Then a stationary distribution  $\pi$  is given by*

$$\pi_j := m_i^{-1} \cdot \sum_{n=0}^{\infty} \mathbb{P}(X_n = j, \tau_i > n | X_0 = i)$$

for all  $j \in E$ . In particular,  $\pi_i = m_i^{-1}$  and  $\pi_k = 0$  for all states  $k$  outside of the communication class belonging to  $i$ .

**Proof:** First of all,  $\pi$  is a probability measure since

$$\begin{aligned} \sum_{j \in E} \sum_{n=0}^{\infty} \mathbb{P}(X_n = j, \tau_i > n | X_0 = i) &= \sum_{n=0}^{\infty} \sum_{j \in E} \mathbb{P}(X_n = j, \tau_i > n | X_0 = i) \\ &= \sum_{n=0}^{\infty} \mathbb{P}(\tau_i > n | X_0 = i) = m_i \end{aligned}$$

The particular statements in the theorem are obvious from theorem 2.16 and the definition of  $\pi$ . The stationarity of  $\pi$  is shown as follows. First we obtain

$$\begin{aligned} \pi_j &= m_i^{-1} \cdot \sum_{n=0}^{\infty} \mathbb{P}(X_n = j, \tau_i > n | X_0 = i) \\ &= m_i^{-1} \cdot \sum_{n=1}^{\infty} \mathbb{P}(X_n = j, \tau_i \geq n | X_0 = i) \\ &= m_i^{-1} \cdot \sum_{n=1}^{\infty} \mathbb{P}(X_n = j, \tau_i > n-1 | X_0 = i) \end{aligned}$$

since  $X_0 = X_{\tau_i} = i$  in the conditioning set  $\{X_0 = i\}$ . Because of

$$\begin{aligned} \mathbb{P}(X_n = j, \tau_i > n-1 | X_0 = i) &= \frac{\mathbb{P}(X_n = j, \tau_i > n-1, X_0 = i)}{\mathbb{P}(X_0 = i)} \\ &= \sum_{k \in E} \frac{\mathbb{P}(X_n = j, X_{n-1} = k, \tau_i > n-1, X_0 = i)}{\mathbb{P}(X_0 = i)} \\ &= \sum_{k \in E} \frac{\mathbb{P}(X_n = j, X_{n-1} = k, \tau_i > n-1, X_0 = i)}{\mathbb{P}(X_{n-1} = k, \tau_i > n-1, X_0 = i)} \\ &\quad \times \frac{\mathbb{P}(X_{n-1} = k, \tau_i > n-1, X_0 = i)}{\mathbb{P}(X_0 = i)} \\ &= \sum_{k \in E} p_{kj} \mathbb{P}(X_{n-1} = k, \tau_i > n-1 | X_0 = i) \end{aligned}$$

we can transform further

$$\pi_j = m_i^{-1} \cdot \sum_{n=1}^{\infty} \sum_{k \in E} p_{kj} \mathbb{P}(X_{n-1} = k, \tau_i > n-1 | X_0 = i)$$

$$= \sum_{k \in E} p_{kj} \cdot m_i^{-1} \sum_{n=0}^{\infty} \mathbb{P}(X_n = k, \tau_i > n | X_0 = i) = \sum_{k \in E} \pi_k p_{kj}$$

which completes the proof.

□

**Theorem 2.25** *Let  $\mathcal{X}$  denote an irreducible, positive recurrent Markov chain. Then  $\mathcal{X}$  has a unique stationary distribution.*

**Proof:** Existence has been shown in theorem 2.24. Uniqueness of the stationary distribution can be seen as follows. Let  $\pi$  denote the stationary distribution as constructed in theorem 2.24 and  $i$  the positive recurrent state that served as recurrence point for  $\pi$ . Further, let  $\nu$  denote any stationary distribution for  $\mathcal{X}$ . Then there is a state  $j \in E$  with  $\nu_j > 0$  and a number  $m \in \mathbb{N}$  with  $P^m(j, i) > 0$ , since  $\mathcal{X}$  is irreducible. Consequently we obtain

$$\nu_i = \sum_{k \in E} \nu_k P^m(k, i) \geq \nu_j P^m(j, i) > 0$$

Hence we can multiply  $\nu$  by a skalar factor  $c$  such that  $c \cdot \nu_i = \pi_i = 1/m_i$ . Denote  $\tilde{\nu} := c \cdot \nu$ .

Let  $\tilde{P}$  denote the transition matrix  $P$  without the  $i$ th column, i.e. we define the  $(j, k)$ th entry of  $\tilde{P}$  by  $\tilde{p}_{jk} = p_{jk}$  if  $k \neq i$  and zero otherwise. Denote further the Dirac measure on  $i$  by  $\delta^i$ , i.e.  $\delta_j^i = 1$  if  $i = j$  and zero otherwise. Then the stationary distribution  $\pi$  can be represented by  $\pi = m_i^{-1} \cdot \delta^i \sum_{n=0}^{\infty} \tilde{P}^n$ .

We first claim that  $m_i \tilde{\nu} = \delta^i + m_i \tilde{\nu} \tilde{P}$ . This is clear for the entry  $\tilde{\nu}_i$  and easily seen for  $\tilde{\nu}_j$  with  $j \neq i$  because in this case  $(\tilde{\nu} \tilde{P})_j = c \cdot (\nu P)_j = \tilde{\nu}_j$ . Now we can proceed with the same argument to see that

$$\begin{aligned} m_i \tilde{\nu} &= \delta^i + (\delta^i + m_i \tilde{\nu} \tilde{P}) \tilde{P} = \delta^i + \delta^i \tilde{P} + m_i \tilde{\nu} \tilde{P}^2 = \dots \\ &= \delta^i \sum_{n=0}^{\infty} \tilde{P}^n = m_i \pi \end{aligned}$$

Hence  $\tilde{\nu}$  already is a probability measure and the skalar factor must be  $c = 1$ . This yields  $\nu = \tilde{\nu} = \pi$  and thus the statement.

□

**Remark 2.26** At a closer look the assumption of irreducibility may be relaxed to some extend. For example, if there is exactly one closed positive recurrent

communication class and a set of transient and inaccessible states (i.e. states  $j$  for which there is no state  $i$  with  $i \rightarrow j$ ), then the above statement still holds although  $\mathcal{X}$  is not irreducible.

A first consequence of the uniqueness is the following simpler representation of the stationary distribution:

**Theorem 2.27** *Let  $\mathcal{X}$  denote an irreducible, positive recurrent Markov chain. Then the stationary distribution  $\pi$  of  $\mathcal{X}$  is given by*

$$\pi_j = m_j^{-1} = \frac{1}{\mathbb{E}(\tau_j | X_0 = j)}$$

for all  $j \in E$ .

**Proof:** Since all states in  $E$  are positive recurrent, the construction in theorem 2.24 can be pursued for any initial state  $j$ . This yields  $\pi_j = m_j^{-1}$  for all  $j \in E$ . The statement now follows from the uniqueness of the stationary distribution.  $\square$

**Corollary 2.28** *For an irreducible, positive recurrent Markov chain, the stationary probability  $\pi_j$  of a state  $j$  coincides with its asymptotic rate of recurrence, i.e.*

$$\lim_{n \rightarrow \infty} \frac{\mathbb{E}(N_j(n) | X_0 = i)}{n} = \pi_j$$

for all  $j \in E$  and independently of  $i \in E$ . Further, if an asymptotic distribution  $p = \lim_{n \rightarrow \infty} \mathbb{P}(X_n = \cdot)$  does exist, then it coincides with the stationary distribution. In particular, it is independent of the initial distribution of  $\mathcal{X}$ .

**Proof:** The first statement immediately follows from equation (2.8). For the second statement, it suffices to employ  $\mathbb{E}(N_j(n) | X_0 = i) = \sum_{l=0}^n P^l(i, j)$ . If an asymptotic distribution  $p$  does exist, then for any initial distribution  $\nu$  we obtain

$$\begin{aligned} p_j &= \lim_{n \rightarrow \infty} (\nu P^n)_j = \sum_{i \in E} \nu_i \lim_{n \rightarrow \infty} P^n(i, j) \\ &= \sum_{i \in E} \nu_i \lim_{n \rightarrow \infty} \frac{\sum_{l=0}^n P^l(i, j)}{n} = \sum_{i \in E} \nu_i \pi_j \\ &= \pi_j \end{aligned}$$

independently of  $\nu$ .  $\square$

## 2.4 Restricted Markov Chains

Now let  $F \subset E$  denote any subset of the state space  $E$ . Define  $\tau_F(k)$  to be the stopping time of the  $k$ th visit of  $\mathcal{X}$  to the set  $F$ , i.e.

$$\tau_F(k+1) := \min\{n > \tau_F(k) : X_n \in F\}$$

with  $\tau_F(0) := 0$ . If  $\mathcal{X}$  is recurrent, then the strong Markov property (theorem 2.7) ensures that the chain  $\mathcal{X}^F = (X_n^F : n \in \mathbb{N})$  with  $X_n^F := X_{\tau_F(n)}$  is a recurrent Markov chain, too. It is called the Markov chain restricted to  $F$ . In case of positive recurrence, we can obtain the stationary distribution of  $\mathcal{X}^F$  from the stationary distribution of  $\mathcal{X}$  in a simple manner:

**Theorem 2.29** *If the Markov chain  $\mathcal{X}$  is positive recurrent, then the stationary distribution of  $\mathcal{X}^F$  is given by*

$$\pi_j^F = \frac{\pi_j}{\sum_{k \in F} \pi_k}$$

for all  $j \in F$ .

**Proof:** Choose any state  $i \in F$  and recall from theorem 2.24 the expression

$$\pi_j := m_i^{-1} \cdot \sum_{n=0}^{\infty} \mathbb{P}(X_n = j, \tau_i > n | X_0 = i)$$

which holds for all  $j \in F$ . For  $\pi_j^F$  we can perform the same construction with respect to the chain  $\mathcal{X}^F$ . By the definition of  $\mathcal{X}^F$  it is clear that the number of visits to the state  $j$  between two consecutive visits to  $i$  is the same for the chains  $\mathcal{X}$  and  $\mathcal{X}^F$ . Hence the sum expression for  $\pi_j^F$ , which is the expectation of that number of visits, remains the same as for  $\pi_j$ . The other factor  $m_i^{-1}$  in the formula above is independent of  $j$  and serves only as a normalization constant, i.e. in order to secure that  $\sum_{j \in E} \pi_j = 1$ . Hence for a construction of  $\pi_j^F$  with respect to  $\mathcal{X}^F$  this needs to be replaced by  $(m_i \cdot \sum_{k \in F} \pi_k)^{-1}$ , which then yields the statement.  $\square$

**Theorem 2.30** *Let  $\mathcal{X} = (X_n : n \in \mathbb{N}_0)$  denote an irreducible and positive recurrent Markov chain with discrete state space  $E$ . Further let  $F \subset E$  denote any subset of  $E$ , and  $\mathcal{X}^F$  the Markov chain restricted to  $F$ . Denote*

$$\tau_F := \min\{n \in \mathbb{N} : X_n \in F\}$$



Then a measure  $\nu$  on  $E$  is stationary for  $\mathcal{X}$  if and only if  $\nu' = (\nu_i : i \in F)$  is stationary for  $\mathcal{X}^F$  and

$$\nu_j = \sum_{k \in F} \nu_k \sum_{n=0}^{\infty} \mathbb{P}(X_n = j, \tau_F > n | X_0 = k) \quad (2.9)$$

for all  $j \in E \setminus F$ .

**Proof:** Due to theorem 2.29 it suffices to prove equation (2.9) for  $j \in E \setminus F$ . Choose any state  $i \in F$  and define

$$\tau_i := \min\{n \in \mathbb{N} : X_n = i\}$$

According to theorem 2.24 the stationary measure  $\nu$  for  $\mathcal{X}$  is given by

$$\nu_j = \nu_i \cdot \sum_{n=0}^{\infty} \mathbb{P}(X_n = j, \tau_i > n | X_0 = i) = \nu_i \cdot \mathbb{E}_i \left( \sum_{n=0}^{\tau_i-1} \mathbf{1}_{X_n=j} \right)$$

for  $j \in E \setminus F$ , where  $\mathbb{E}_i$  denotes the conditional expectation given  $X_0 = i$ . Define further

$$\tau_i^F := \min\{n \in \mathbb{N} : X_n^F = i\}$$

Because of the strong Markov property we can proceed as

$$\begin{aligned} \nu_j &= \nu_i \cdot \mathbb{E}_i \left( \sum_{n=0}^{\tau_i^F-1} \mathbb{E}_{X_n^F} \sum_{m=0}^{\tau_F-1} \mathbf{1}_{X_m=j} \right) \\ &= \nu_i \cdot \sum_{k \in F} \mathbb{E}_i \left( \sum_{n=0}^{\tau_i^F-1} \mathbf{1}_{X_n^F=k} \right) \cdot \mathbb{E}_k \left( \sum_{m=0}^{\tau_F-1} \mathbf{1}_{X_m=j} \right) \end{aligned}$$

Regarding the restricted Markov chain  $\mathcal{X}^F$ , theorem 2.24 states that

$$\mathbb{E}_i \left( \sum_{n=0}^{\tau_i^F-1} \mathbf{1}_{X_n^F=k} \right) = \sum_{n=0}^{\infty} \mathbb{P}(X_n^F = k, \tau_i^F > n | X_0^F = i) = \frac{\nu_k}{\nu_i}$$

for all  $k \in F$ . Hence we obtain

$$\nu_j = \sum_{k \in F} \nu_k \sum_{n=0}^{\infty} \mathbb{P}(X_n = j, \tau_F > n | X_0 = k)$$

which was to be proven.

□

## 2.5 Conditions for Positive Recurrence

In the third part of this course we will need some results on the behaviour of a Markov chain on a finite subset of its state space. As a first fundamental result we state

**Theorem 2.31** *An irreducible Markov chain with finite state space  $F$  is positive recurrent.*

**Proof:** For all  $n \in \mathbb{N}$  and  $i \in F$  we have  $\sum_{j \in E} P^n(i, j) = 1$ . Hence it is not possible that  $\lim_{n \rightarrow \infty} P^n(i, j) = 0$  for all  $j \in F$ . Thus there is one state  $h \in F$  such that  $r_{hh} = \sum_{n=0}^{\infty} P^n(h, h) = \infty$ , which means by corollary 2.15 that  $h$  is recurrent and by irreducibility that the chain is recurrent.

If the chain were null recurrent, then according to the relation in (2.8)

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n P^k(i, j) = 0$$

would hold for all  $j \in F$ , independently of  $i$  because of irreducibility. But this would imply that  $\lim_{n \rightarrow \infty} P^n(i, j) = 0$  for all  $j \in F$ , which contradicts our first observation in this proof. Hence the chain must be positive recurrent.

□

For irreducible Markov chains the condition  $\mathbb{E}(\tau_i | X_0 = i) < \infty$  implies positive recurrence of state  $i$  and hence positive recurrence of the whole chain. Writing  $\tau_F$  for the time of the first visit to the set  $F$ , we now can state the following generalization of this condition:

**Theorem 2.32** *Let  $\mathcal{X}$  denote an irreducible Markov chain with state space  $E$  and be  $F \subset E$  a finite subset of  $E$ . The chain  $\mathcal{X}$  is positive recurrent if and only if  $\mathbb{E}(\tau_F | X_0 = i) < \infty$  for all  $i \in F$ .*

**Proof:** If  $\mathcal{X}$  is positive recurrent, then  $\mathbb{E}(\tau_F | X_0 = i) \leq \mathbb{E}(\tau_i | X_0 = i) < \infty$  for all  $i \in F$ , by the definition of positive recurrence.

Now assume that  $\mathbb{E}(\tau_F | X_0 = i) < \infty$  for all  $i \in F$ . Define the stopping times  $\sigma(i) := \min\{k \in \mathbb{N} : X_k^F = i\}$  and random variables  $Y_k := \tau_F(k) - \tau_F(k-1)$ . Since  $F$  is finite,  $m := \max_{j \in F} \mathbb{E}(\tau_F | X_0 = j) < \infty$ . We shall denote the

conditional expectation given  $X_0 = i$  by  $\mathbb{E}_i$ . For  $i \in F$  we now obtain

$$\begin{aligned} \mathbb{E}(\tau_i | X_0 = i) &= \mathbb{E}_i \left( \sum_{k=1}^{\sigma(i)} Y_k \right) = \sum_{k=1}^{\infty} \mathbb{E}_i \left( \mathbb{E}(Y_k | X_{\tau_F(k-1)}) \cdot 1_{k \leq \sigma(i)} \right) \\ &\leq m \cdot \sum_{k=1}^{\infty} \mathbb{P}(\sigma(i) \geq k | X_0 = i) = m \cdot \mathbb{E}(\sigma(i) | X_0 = i) \end{aligned}$$

Since  $F$  is finite,  $\mathcal{X}^F$  is positive recurrent by theorem 2.31. Hence we know that  $\mathbb{E}(\sigma(i) | X_0 = i) < \infty$ , and thus  $\mathbb{E}(\tau_i | X_0 = i) < \infty$  which shows that  $\mathcal{X}$  is positive recurrent.

□

An often difficult problem is to determine whether a given Markov chain is positive recurrent or not. Concerning this, we now introduce one of the most important criteria for the existence of stationary distributions of Markov chains occurring in queueing theory. It is known as **Foster's criterion**.

**Theorem 2.33** *Let  $\mathcal{X}$  denote an irreducible Markov chain with countable state space  $E$  and transition matrix  $P$ . Further let  $F$  denote a finite subset of  $E$ . If there is a function  $h : E \rightarrow \mathbb{R}$  with  $\inf\{h(i) : i \in E\} > -\infty$ , such that the conditions*

$$\sum_{k \in E} p_{ik} h(k) < \infty \quad \text{and} \quad \sum_{k \in E} p_{jk} h(k) \leq h(j) - \varepsilon$$

*hold for some  $\varepsilon > 0$  and all  $i \in F$  and  $j \in E \setminus F$ , then  $\mathcal{X}$  is positive recurrent.*

**Proof:** Without loss of generality we can assume  $h(i) \geq 0$  for all  $i \in E$ , since otherwise we only need to increase  $h$  by a suitable constant. Define the stopping time  $\tau_F := \min\{n \in \mathbb{N}_0 : X_n \in F\}$ . First we observe that

$$\begin{aligned} \mathbb{E}(h(X_{n+1}) \cdot 1_{\tau_F > n+1} | X_0, \dots, X_n) &\leq \mathbb{E}(h(X_{n+1}) \cdot 1_{\tau_F > n} | X_0, \dots, X_n) \\ &= 1_{\tau_F > n} \cdot \sum_{k \in E} p_{X_n, k} h(k) \\ &\leq 1_{\tau_F > n} \cdot (h(X_n) - \varepsilon) \\ &= h(X_n) \cdot 1_{\tau_F > n} - \varepsilon \cdot 1_{\tau_F > n} \end{aligned}$$

holds for all  $n \in \mathbb{N}_0$ , where the first equality is due to (5.3). We now proceed with

$$\begin{aligned}
0 &\leq \mathbb{E}(h(X_{n+1}) \cdot 1_{\tau_F > n+1} | X_0 = i) \\
&= \mathbb{E}(\mathbb{E}(h(X_{n+1}) \cdot 1_{\tau_F > n+1} | X_0, \dots, X_n) | X_0 = i) \\
&\leq \mathbb{E}(h(X_n) \cdot 1_{\tau_F > n} | X_0 = i) - \varepsilon \mathbb{P}(\tau_F > n | X_0 = i) \\
&\leq \dots \\
&\leq \mathbb{E}(h(X_0) \cdot 1_{\tau_F > 0} | X_0 = i) - \varepsilon \sum_{k=0}^n \mathbb{P}(\tau_F > k | X_0 = i)
\end{aligned}$$

which holds for all  $i \in E \setminus F$  and  $n \in \mathbb{N}_0$ . For  $n \rightarrow \infty$  this implies

$$\mathbb{E}(\tau_F | X_0 = i) = \sum_{k=0}^{\infty} \mathbb{P}(\tau_F > k | X_0 = i) \leq h(i)/\varepsilon < \infty$$

for  $i \in E \setminus F$ . Now the mean return time to the state set  $F$  is bounded by

$$\begin{aligned}
\mathbb{E}(\tau_F | X_0 = i) &= \sum_{j \in F} p_{ij} + \sum_{j \in E \setminus F} p_{ij} \mathbb{E}(\tau_F + 1 | X_0 = j) \\
&\leq 1 + \varepsilon^{-1} \sum_{j \in E} p_{ij} h(j) < \infty
\end{aligned}$$

for all  $i \in F$ , which completes the proof.

□

## 2.6 The M/M/1 queue in discrete time

Choose any parameters  $0 < p, q < 1$ . Let the arrival process be distributed as a Bernoulli process with parameter  $p$  and the service times  $(S_n : n \in \mathbb{N}_0)$  be iid according to the geometric distribution with parameter  $q$ .

The geometric service time distribution and the Bernoulli arrival process have been chosen because this simplifies the formulation of the system process in terms of a Markov model due to the following **memoryless property**:

**Theorem 2.34** *Let  $S$  be distributed geometrically with parameter  $q$ , i.e. let  $\mathbb{P}(S = k) = (1 - q)^{k-1}q$  for all  $k \in \mathbb{N}$ . Then  $\mathbb{P}(S = k | S > k - 1) = q$  holds for the conditional distribution, independently of  $k$ . Likewise, if  $Z_n$  is the  $n$ th inter-arrival time of a Bernoulli process with parameter  $p$ , then the relation  $\mathbb{P}(Z_n = k | Z_n > k - 1) = p$  holds, independently of  $k$  and  $n$ .*

**Proof:** First the proof for the geometric distribution: For all  $k \in \mathbb{N}$ , the argument

$$\begin{aligned}\mathbb{P}(S = k | S > k - 1) &= \frac{\mathbb{P}(S = k, S > k - 1)}{\mathbb{P}(S > k - 1)} = \frac{\mathbb{P}(S = k)}{\mathbb{P}(S > k - 1)} \\ &= \frac{(1 - q)^{k-1} q}{(1 - q)^{k-1}} = q\end{aligned}$$

holds, which shows the first statement. For a Bernoulli process, the  $n$ th inter-arrival time  $Z_n = T_n - T_{n-1}$  is distributed geometrically with parameter  $p$ , due to the strong Markov property. This completes the proof for the second statement.

□

Thus the memoryless property states that no matter how long a service time or an inter-arrival time has already passed, the probability of a service completion or an arrival at the next time instant is always the same. Hence the system process  $\mathcal{Q} = (Q_n : n \in \mathbb{N}_0)$  of the M/M/1 queue in discrete time with arrival process  $\mathcal{T}$  and service times  $S_n$  can be formulated easily as a homogeneous Markov chain. It has state space  $E = \mathbb{N}_0$  and transition probabilities  $p_{01} := p$ ,  $p_{00} := 1 - p$ , and

$$p_{ij} := \begin{cases} p(1 - q), & j = i + 1 \\ pq + (1 - p)(1 - q), & j = i \\ q(1 - p), & j = i - 1 \end{cases}$$

for  $i \geq 1$ . Because of the simple state space, the transition matrix can be displayed in the form of a triangular matrix

$$P = \begin{pmatrix} 1 - p & p & 0 & \dots \\ q(1 - p) & pq + (1 - p)(1 - q) & p(1 - q) & \ddots \\ 0 & q(1 - p) & pq + (1 - p)(1 - q) & \ddots \\ \vdots & \ddots & \ddots & \ddots \end{pmatrix}$$

Since  $p, q > 0$ , the chain  $\mathcal{Q}$  is irreducible. If  $p < q$ , then  $h(n) := n$  defines a function which satisfies the conditions for Foster's criterion, as

$$\begin{aligned}\sum_{k=0}^{\infty} p_{ik} h(k) &= q(1 - p) \cdot (i - 1) + (qp + (1 - q)(1 - p)) \cdot i \\ &\quad + p(1 - q) \cdot (i + 1) \\ &= i - q(1 - p) + p(1 - q) = i - q + p \leq i - \varepsilon\end{aligned}$$

for all  $i \in \mathbb{N}$ , with  $\varepsilon = q - p > 0$ , and  $\sum_{k=0}^{\infty} p_{0k} \cdot h(k) = p < \infty$  show. The ratio  $p/q$  is called the **load** of the queue. Thus the system process  $\mathcal{Q}$  is positive recurrent if the queue load is less than one.

In order to derive a stationary distribution for  $\mathcal{Q}$ , we first introduce notations  $p' := p(1 - q)$  and  $q' := q(1 - p)$ . Then we translate the condition  $\pi P = \pi$  into the equations

$$\pi_0 = \pi_0(1 - p) + \pi_1 q' \quad (2.10)$$

$$\pi_1 = \pi_0 p + \pi_1(1 - p - q') + \pi_2 q' \quad (2.11)$$

$$\pi_n = \pi_{n-1} p' + \pi_n(1 - (p' + q')) + \pi_{n+1} q' \quad (2.12)$$

for all  $n \geq 2$ . For the solution, we guess the geometric form

$$\pi_{n+1} = \pi_n \cdot r$$

for all  $n \geq 1$ , with  $r > 0$ . Thus equation (2.12) becomes

$$0 = \pi_n p' - \pi_n r(p' + q') + \pi_n r^2 q' = \pi_n (p' - r(p' + q') + r^2 q')$$

for all  $n \geq 1$ , which leads for non-trivial  $\pi \neq 0$  to the roots  $r = 1$  and  $r = p'/q'$  of the quadratic term.

In the first case  $r = 1$ , we obtain  $\pi_{n+1} = \pi_n$  for all  $n \geq 1$ . This implies  $\sum_{j \in E} \pi_j = \infty$  and thus cannot lead to a stationary distribution. Hence in the case  $r = 1$  the geometric approach is not successful.

The second root  $r = p'/q'$  allows solutions for the other equations (2.10) and (2.11) too. This can be checked as follows: First, the relation

$$\pi_1 = \pi_0 \frac{p}{q'} = \pi_0 \frac{\rho}{1 - p}$$

is a requirement from equation (2.10). Then the second equation (2.11) yields

$$\begin{aligned} \pi_2 &= \frac{1}{q'} (\pi_1(p' + q') - \pi_0 p) = \frac{1}{q'} \left( \frac{p}{q'} (p' + q') - p \right) \pi_0 \\ &= \pi_0 \frac{p}{q'} \left( \frac{p' + q'}{q'} - 1 \right) = \pi_1 \frac{p'}{q'} \end{aligned}$$

in accordance with our geometric approach. Now normalization of  $\pi$  leads to

$$1 = \sum_{n=0}^{\infty} \pi_n = \pi_0 \left( 1 + \frac{p}{q'} \sum_{n=1}^{\infty} \left( \frac{p'}{q'} \right)^{n-1} \right)$$

from which we obtain

$$\begin{aligned}\pi_0 &= \left(1 + \frac{p}{q'} \sum_{n=1}^{\infty} \left(\frac{p'}{q'}\right)^{n-1}\right)^{-1} = \left(1 + \frac{p}{q'(1 - p'/q')}\right)^{-1} \\ &= \left(1 + \frac{p}{q' - p'}\right)^{-1} = (q' - p')(q' - p' + p)^{-1} = \frac{q - p}{q} \\ &= 1 - \rho\end{aligned}$$

with  $\rho := p/q$ , because of  $q' - p' = q - p$ . Hence the approach  $\pi_{n+1} = \pi_n \cdot r$  with  $r = p'/q'$  leads to a solution of  $\pi P = \pi$ .

Note that  $r < 1$  if and only if  $p < q$ . Further, the mean inter-arrival time is  $\mathbb{E}(T_1) = 1/p$  and the mean service time is  $\mathbb{E}(S_1) = 1/q$ . Thus the geometric approach is successful if the so-called **stability condition**

$$\rho = \frac{p}{q} = \frac{\mathbb{E}(S_1)}{\mathbb{E}(T_1)} < 1$$

holds. This condition simply postulates that the mean service time be shorter than the mean inter-arrival time. In this case, the stationary distribution  $\pi$  of  $Q$  has the form

$$\pi_0 = 1 - \rho \quad \text{and} \quad \pi_n = (1 - \rho) \frac{\rho}{1 - p} r^{n-1}$$

for all  $n \geq 1$ . It thus is a modified geometric distribution with parameter  $r = p'/q' < 1$ .

## Notes

Markov chains originate from a series of papers written by A. Markov at the beginning of the 20th century. His first application is given here as exercise 2.37. However, methods and terminology at that time were very different from today's presentations.

The literature on Markov chains is perhaps the most extensive in the field of stochastic processes. This is not surprising, as Markov chains form a simple and useful starting point for the introduction of other processes.

Textbook presentations are given in Feller [8], Breiman [1], Karlin and Taylor [14], or Çinlar [4], to name but a few. The treatment in Ross [18] contains the useful concept of time-reversible Markov chains. An exhaustive introduction to Markov chains on general state spaces and conditions for their positive recurrence is given in Meyn and Tweedie [16].

**Exercise 2.35** Let  $(X_n : n \in \mathbb{N}_0)$  be a family of iid random variables with discrete state space. Show that  $\mathcal{X} = (X_n : n \in \mathbb{N}_0)$  is a homogeneous Markov chain.

**Exercise 2.36** Let  $(X_n : n \in \mathbb{N}_0)$  be iid random variables on  $\mathbb{N}_0$  with probabilities  $a_i := \mathbb{P}(X_n = i)$  for all  $n, i \in \mathbb{N}_0$ . The event  $X_n > \max(X_0, \dots, X_{n-1})$  for  $n \geq 1$  is called a record at time  $n$ . Define  $T_i$  as the time of the  $i$ th record, i.e.  $T_0 := 0$  and  $T_{i+1} := \min\{n \in \mathbb{N} : X_n > X_{T_i}\}$  for all  $i \in \mathbb{N}_0$ . Denote the  $i$ th record value by  $R_i := X_{T_i}$ . Show that  $(R_i : i \in \mathbb{N}_0)$  and  $((R_i, T_i) : i \in \mathbb{N}_0)$  are Markov chains by determining their transition probabilities.

**Exercise 2.37** Diffusion model by Bernoulli and Laplace

The following is a stochastic model for the flow of two incompressible fluids between two containers: Two boxes contain  $m$  balls each. Of these  $2m$  balls,  $b$  are black and the others are white. The system is said to be in state  $i$  if the first box contains  $i$  black balls. A state transition is performed by choosing one ball out of each box at random (meaning here that each ball is chosen with equal probability) and then interchanging the two. Derive a Markov chain model for the system and determine the transition probabilities.

**Exercise 2.38** Let  $\mathcal{X}$  denote a Markov chain with  $m < \infty$  states. Show that if state  $j$  is accessible from state  $i$ , then it is accessible in at most  $m - 1$  transitions.

**Exercise 2.39** Let  $p = (p_n : n \in \mathbb{N}_0)$  be a discrete probability distribution and define

$$P = \begin{pmatrix} p_0 & p_1 & p_2 & \dots \\ & p_0 & p_1 & \ddots \\ & & p_0 & \ddots \\ & & & \ddots \end{pmatrix}$$

with all non-specified entries being zero. Let  $\mathcal{X}$  denote a Markov chain with state space  $\mathbb{N}_0$  and transition matrix  $P$ . Derive an expression (in terms of discrete convolutions) for the transition probabilities  $\mathbb{P}(X_{n+m} = j | X_n = i)$  with  $n, m \in \mathbb{N}_0$  and  $i, j \in \mathbb{N}_0$ . Apply the result to the special case of a Bernoulli process (see example 2.3).

**Exercise 2.40** Prove equation (2.6).




**Exercise 2.41** Prove the equation  $P^n(i, j) = \sum_{k=1}^n F_k(i, j) P^{n-k}(j, j)$  for all  $n \in \mathbb{N}$  and  $i, j \in E$ .

**Exercise 2.42** Let  $\mathcal{X}$  denote a Markov chain with state space  $E = \{1, \dots, 10\}$  and transition matrix

$$P = \begin{pmatrix} 1/2 & 0 & 1/2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1/3 & 0 & 0 & 0 & 0 & 2/3 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1/3 & 1/3 & 0 & 0 & 0 & 1/3 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1/4 & 0 & 3/4 & 0 \\ 0 & 0 & 1/4 & 1/4 & 0 & 0 & 0 & 1/4 & 0 & 1/4 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1/3 & 0 & 0 & 1/3 & 0 & 0 & 0 & 0 & 1/3 \end{pmatrix}$$

Reorder the states according to their communication classes and determine the resulting form of the transition matrix as in representation (2.4). Determine further a transition graph, in which



means that  $f_{ij} > 0$ .

**Exercise 2.43** Prove equation (2.7).

Hint: Derive a representation of  $N_j$  in terms of the random variables

$$A_n := \begin{cases} 1, & X_n = j \\ 0, & X_n \neq j \end{cases}$$

**Exercise 2.44** Prove corollary 2.15.

**Exercise 2.45** Prove remark 2.26.

**Exercise 2.46** A server's up time is  $k$  time units with probability  $p_k = 2^{-k}$ ,  $k \in \mathbb{N}$ . After failure the server is immediately replaced by an identical new one. The up time of the new server is of course independent of the behaviour of all preceding servers.

Let  $X_n$  denote the remaining up time of the server at time  $n \in \mathbb{N}_0$ . Determine the transition probabilities for the Markov chain  $\mathcal{X} = (X_n : n \in \mathbb{N}_0)$  and determine the stationary distribution of  $\mathcal{X}$ .

**Exercise 2.47** Let  $P$  denote the transition matrix of an irreducible Markov chain  $\mathcal{X}$  with discrete state space  $E = F \cup F^c$ , where  $F^c = E \setminus F$ . Write  $P$  in block notation as

$$P = \begin{pmatrix} P_{FF} & P_{FF^c} \\ P_{F^cF} & P_{F^cF^c} \end{pmatrix}$$

Show that the Markov chain  $\mathcal{X}^F$  restricted to the state space  $F$  has transition matrix

$$P^F = P_{FF} + P_{FF^c}(I - P_{F^cF^c})^{-1}P_{F^cF}$$

with  $I$  denoting the identity matrix on  $F^c$ .

**Exercise 2.48** Let  $\mathcal{X}$  denote a Markov chain with state space  $E = \{0, \dots, m\}$  and transition matrix

$$P = \begin{pmatrix} p_{00} & p_{01} & & & \\ p_{10} & p_{11} & p_{12} & & \\ & p_{21} & p_{22} & p_{23} & \\ & & \ddots & \ddots & \ddots \\ & & & p_{m,m-1} & p_{mm} \end{pmatrix}$$

where  $p_{ij} > 0$  for  $|i - j| = 1$ . Show that the stationary distribution  $\pi$  of  $\mathcal{X}$  is uniquely determined by

$$\pi_n = \pi_0 \cdot \prod_{i=1}^n \frac{p_{i-1,i}}{p_{i,i-1}} \quad \text{and} \quad \pi_0 = \left( \sum_{j=0}^m \prod_{i=1}^j \frac{p_{i-1,i}}{p_{i,i-1}} \right)^{-1}$$

for all  $n = 1, \dots, m$ .

Use this result to determine the stationary distribution of the Bernoulli–Laplace diffusion model with  $b = m$  (see exercise 2.37).

**Exercise 2.49** Show that the second condition in theorem 2.33 can be substituted by the condition

$$\sum_{j \in E} p_{ij} h(j) \leq h(i) - 1 \quad \text{for all } i \in E \setminus F.$$

**Exercise 2.50** Show the following complement to theorem 2.33: Let  $P$  denote the transition matrix of a positive recurrent Markov chain with discrete state space  $E$ . Then there is a function  $h : E \rightarrow \mathbb{R}$  and a finite subset  $F \subset E$  such that

$$\begin{aligned} \sum_{j \in E} p_{ij} h(j) &< \infty && \text{for all } i \in F, \text{ and} \\ \sum_{j \in E} p_{ij} h(j) &\leq h(i) - 1 && \text{for all } i \in E \setminus F. \end{aligned}$$

Hint: Consider the conditional expectation of the remaining time until returning to a fixed set  $F$  of states.

**Exercise 2.51** For the discrete, non-negative random walk with transition matrix

$$P = \begin{pmatrix} p_{00} & p_{01} & & & \\ p_{10} & 0 & p_{12} & & \\ & p_{10} & 0 & p_{12} & \\ & & \ddots & \ddots & \ddots \end{pmatrix}$$

determine the criterion of positive recurrence according to theorem 2.33.



## Chapter 3

# Homogeneous Markov Processes on Discrete State Spaces

In the present chapter we will transfer the discrete time results of the previous chapter to Markov processes in continuous time.

### 3.1 Definition

Define  $T_0 := 0$  and let  $(T_n : n \in \mathbb{N})$  denote a sequence of positive real-valued random variables with  $T_{n+1} > T_n$  for all  $n \in \mathbb{N}_0$  and  $T_n \rightarrow \infty$  as  $n \rightarrow \infty$ . Further, let  $E$  denote a countable state space and  $(X_n : n \in \mathbb{N}_0)$  a sequence of  $E$ -valued random variables. A process  $\mathcal{Y} = (Y_t : t \in \mathbb{R}_0^+)$  in continuous time with

$$Y_t := X_n \quad \text{for} \quad T_n \leq t < T_{n+1}$$

is called a **pure jump process**. The variable  $H_n := T_{n+1} - T_n$  (resp.  $X_n$ ) is called the  $n$ th **holding time** (resp. the  $n$ th **state**) of the process  $\mathcal{Y}$ . If further  $\mathcal{X} = (X_n : n \in \mathbb{N}_0)$  is a Markov chain with transition matrix  $P = (p_{ij})_{i,j \in E}$  and the variables  $H_n$  are independent and distributed exponentially with parameter  $\lambda_{X_n}$  only depending on the state  $X_n$ , then  $\mathcal{Y}$  is called homogeneous **Markov process** with discrete **state space**  $E$ . The chain  $\mathcal{X}$  is called the **embedded Markov chain** of  $\mathcal{Y}$ . As a technical assumption we always agree upon the condition  $\hat{\lambda} := \sup\{\lambda_i : i \in E\} < \infty$ , i.e. the parameters for the exponential holding times shall be bounded.

An immediate consequence of the definition is that the paths of a Markov process are step functions. The lengths of the holding times are almost certainly strictly

positive, since exponential distributions are zero with probability zero.

**Example 3.1 Poisson process**

Define  $X_n := n$  deterministically. Then  $\mathcal{X} = (X_n : n \in \mathbb{N}_0)$  is a Markov chain with state space  $E = \mathbb{N}_0$  and transition probabilities  $p_{n,n+1} = 1$  for all  $n \in \mathbb{N}_0$ . Let the holding times  $H_n$  be distributed exponentially with identical parameter  $\lambda > 0$ . Then the resulting process  $\mathcal{Y}$  as defined in the above definition is a Markov process with state space  $\mathbb{N}_0$ . It is called **Poisson process** with **intensity** (also: rate or parameter)  $\lambda$ .

Next we want to prove a property similar to the Markov property for Markov chains in discrete time. To this aim, we need to show the **memoryless property** for the exponential distribution, which is the analogue to the memoryless property for geometric distributions in discrete time.

**Lemma 3.2** *Let  $H$  denote a random variable having an exponential distribution with parameter  $\lambda$ . Then the memoryless property*

$$\mathbb{P}(H > t + s | H > s) = \mathbb{P}(H > t)$$

*holds for all time durations  $s, t > 0$ .*

**Proof:** We immediately check

$$\begin{aligned} \mathbb{P}(H > t + s | H > s) &= \frac{\mathbb{P}(H > t + s, H > s)}{\mathbb{P}(H > s)} = \frac{\mathbb{P}(H > t + s)}{\mathbb{P}(H > s)} \\ &= \frac{e^{-\lambda \cdot (t+s)}}{e^{-\lambda \cdot s}} = e^{-\lambda \cdot t} = \mathbb{P}(H > t) \end{aligned}$$

which holds for all  $s, t > 0$ .

□

**Theorem 3.3** *Let  $\mathcal{Y}$  denote a Markov process with discrete state space  $E$ . Then the Markov property*

$$\mathbb{P}(Y_t = j | Y_u : u \leq s) = \mathbb{P}(Y_t = j | Y_s)$$

*holds for all times  $s < t$  and states  $j \in E$ .*

**Proof:** Denote the state at time  $s$  by  $Y_s = i$ . Because of the memoryless property of the exponential holding times, the remaining time in state  $i$  is distributed exponentially with parameter  $\lambda_i$ , no matter how long the preceding holding time has

been. After the holding time in the present state elapses, the process changes to another state  $j$  according to the homogeneous Markov chain  $\mathcal{X}$ . Hence the probability for the next state being  $j$  is given by  $p_{ij}$ , independently of any state of the process before time  $s$ . Now another exponential holding time begins, and thus the past before time  $s$  will not have any influence on the future of the process  $\mathcal{Y}$ .

□

Analogous to the discrete time case, for any two time instances  $s < t$  the conditional probabilities  $\mathbb{P}(Y_t = j | Y_s = i)$  shall be called the **transition probabilities** from time  $s$  to time  $t$ . We will now derive a recursion formula for the transition probabilities of a Markov process by conditioning on the number of jumps between time  $s$  and time  $t$ :

**Theorem 3.4** *The transition probabilities of a Markov process  $\mathcal{Y}$  are given by*

$$\mathbb{P}(Y_t = j | Y_s = i) = \sum_{n=0}^{\infty} P_{ij}^{(n)}(s, t)$$

for all times  $s < t$  and states  $i, j \in E$ , with

$$P_{ij}^{(0)}(s, t) = \delta_{ij} \cdot e^{-\lambda_i \cdot (t-s)}$$

and recursively

$$P_{ij}^{(n+1)}(s, t) = \int_s^t e^{-\lambda_i \cdot u} \lambda_i \sum_{k \in E} p_{ik} P_{kj}^{(n)}(u, t) du$$

for all  $n \in \mathbb{N}_0$ .

**Proof:** The above representation follows immediately by conditioning on the number of jumps in  $]s, t]$ . The expressions  $P_{ij}^{(n)}(s, t)$  represent the conditional probabilities that  $Y_t = j$  and there are  $n$  jumps in  $]s, t]$  given that  $Y_s = i$ . In the recursion formula the integral comprises all times  $u$  of a possible first jump along with the Lebesgue density  $e^{-\lambda_i \cdot u} \lambda_i$  of this event, after which the probability of  $n$  remaining jumps reaching state  $j$  at time  $t$  is given by  $\sum_{k \in E} p_{ik} P_{kj}^{(n)}(u, t)$ .

□

For every two time instances  $s < t$ , define the **transition probability matrix**  $P(s, t)$  from time  $s$  to time  $t$  by its entries

$$P_{ij}(s, t) := \mathbb{P}(Y_t = j | Y_s = i)$$

Using the recursion formula, it is shown by induction on  $n$  that the conditional probabilities  $P_{ij}^{(n)}(s, t)$  are homogeneous in time, i.e. they satisfy

$$P_{ij}^{(n)}(s, t) = P_{ij}^{(n)}(0, t - s)$$

for all  $s < t$ . Thus we can from now on restrict the analysis to the transition probability matrices

$$P(t) := P(0, t)$$

with  $t \geq 0$ . With this notation the Markov property yields the **Chapman–Kolmogorov equations**

$$P(s + t) = P(s)P(t)$$

for all time durations  $s, t \geq 0$ . Thus the family  $\{P(t) : t \geq 0\}$  of transition probability matrices forms a semi-group under the composition of matrix multiplication. In particular, we obtain for the neutral element of this semi-group  $P(0) = I_E := (\delta_{ij})_{i,j \in E}$  with  $\delta_{ij} = 1$  for  $i = j$  and zero otherwise.

In order to derive a simpler expression for the transition probability matrices, we need to introduce another concept, which will be called the **generator matrix**. This is defined as the matrix  $G = (g_{ij})_{i,j \in E}$  on  $E$  with entries


$$g_{ij} := \begin{cases} -\lambda_i \cdot (1 - p_{ii}), & i = j \\ \lambda_i \cdot p_{ij}, & i \neq j \end{cases}$$

for all states  $i, j \in E$ . In particular, the relation

$$g_{ii} = - \sum_{j \neq i} g_{ij} \tag{3.1}$$

holds for all  $i \in E$ .

The  $(i, j)$ th entry of the generator  $G$  is called the **infinitesimal transition rate** from state  $i$  to state  $j$ . Using these, we can illustrate the dynamics of a Markov process in a directed graph where the nodes represent the states and an edge



means that  $g_{ij} = r > 0$ . Such a graph is called a **state transition graph** of the Markov process. With the convention  $p_{ii} = 0$  the state transition graph uniquely determines the Markov process.






Figure 3.1: Poisson process

**Example 3.5** The state transition graph of the Poisson process with intensity  $\lambda$  (see example 3.1) is given by

**Theorem 3.6** *The transition probabilities  $P_{ij}(t)$  of a Markov process satisfy the systems*

$$\frac{dP_{ij}(t)}{dt} = \sum_{k \in E} P_{ik}(t)g_{kj} = \sum_{k \in E} g_{ik}P_{kj}(t)$$

*of differential equations. These are called the **Kolmogorov forward and backward equations**.*

**Proof:** From the representation in theorem 3.4, it follows by induction on the number of jumps that all restricted probabilities  $P^{(n)}(t)$  are Lebesgue integrable with respect to  $t$  over finite intervals. Since the sum of all  $P_{ij}^{(n)}(t)$  is a probability and thus bounded, we conclude by majorized convergence that also  $P(t)$  is Lebesgue integrable with respect to  $t$  over finite intervals.

Now we can state the recursion

$$P_{ij}(t) = e^{-\lambda_i \cdot t} \cdot \delta_{ij} + \int_0^t e^{-\lambda_i \cdot s} \lambda_i \sum_{k \in E} p_{ik} P_{kj}(t-s) ds$$

which results from conditioning on the time  $s$  of the first jump from state  $i$ . We obtain further

$$P_{ij}(t) = e^{-\lambda_i \cdot t} \cdot \left( \delta_{ij} + \int_0^t e^{+\lambda_i \cdot u} \lambda_i \sum_{k \in E} p_{ik} P_{kj}(u) du \right)$$

by substituting  $u = t - s$  in the integral. Since  $\sum_{k \in E} p_{ik} = 1$  is bounded, we conclude that  $P(t)$  is continuous in  $t$ . Further, we can differentiate  $P(t)$  as given in the recursion and obtain

$$\frac{dP_{ij}(t)}{dt} = -\lambda_i e^{-\lambda_i \cdot t} \cdot \left( \delta_{ij} + \int_0^t f(u) du \right) + e^{-\lambda_i \cdot t} \cdot f(t)$$

with  $f$  denoting the integrand function. This means nothing else than

$$\begin{aligned}\frac{dP_{ij}(t)}{dt} &= -\lambda_i P_{ij}(t) + \lambda_i \sum_{k \in E} p_{ik} P_{kj}(t) \\ &= -\lambda_i(1 - p_{ii}) \cdot P_{ii}(t) + \sum_{k \neq i} g_{ik} P_{kj}(t)\end{aligned}$$

and thus proves the backward equations. For the forward equations, one only needs to use the Chapman–Kolmogorov equations and apply the backward equations in

$$\begin{aligned}\frac{dP_{ij}(t)}{dt} &= \lim_{h \rightarrow 0} \frac{P_{ij}(t+h) - P_{ij}(t)}{h} = \lim_{h \rightarrow 0} \sum_{k \in E} P_{ik}(t) \frac{P_{kj}(h) - \delta_{kj}}{h} \\ &= \sum_{k \in E} P_{ik}(t) \lim_{h \rightarrow 0} \frac{P_{kj}(h) - P_{kj}(0)}{h} = \sum_{k \in E} P_{ik}(t) g_{kj}\end{aligned}$$

which holds for all  $i, j \in E$ .

□

**Theorem 3.7** *The transition probability matrices can be expressed in terms of the generator by*

$$P(t) = e^{G \cdot t} := \sum_{n=0}^{\infty} \frac{t^n}{n!} G^n$$

for all  $t \geq 0$ , with  $G^n$  denoting the  $n$ th power of the matrix  $G$ .

**Proof:** First we validate the solution by

$$\frac{d}{dt} e^{G \cdot t} = \frac{d}{dt} \sum_{n=0}^{\infty} \frac{t^n}{n!} G^n = \sum_{n=1}^{\infty} G^n \frac{d}{dt} \frac{t^n}{n!} = \sum_{n=1}^{\infty} G^n \frac{t^{n-1}}{(n-1)!} = G e^{G \cdot t}$$

which holds for all  $t \geq 0$ . Furthermore, it is obvious that

$$G e^{G \cdot t} = G \sum_{n=0}^{\infty} \frac{t^n}{n!} G^n = \left( \sum_{n=0}^{\infty} \frac{t^n}{n!} G^n \right) G = e^{G \cdot t} G$$

and thus  $P(t) = e^{G \cdot t}$  is a solution of Kolmogorov's forward and backward equations.

Now we show uniqueness of the solution. Let  $\tilde{P}(t)$  denote another solution of the forward equations. The differential equations with initial condition translate into the integral equations

$$P(t) = I_E + \int_0^t P(u)G \, du \quad \text{and} \quad \tilde{P}(t) = I_E + \int_0^t \tilde{P}(u)G \, du$$

Define a norm for matrices  $M = (m_{ij})_{i,j \in E}$  on  $E$  by

$$\|M\| := \sup \left\{ \sum_{j \in E} |m_{ij}| : i \in E \right\}$$

Then  $\|G\| \leq 2 \cdot \hat{\lambda}$  and  $\|AB\| \leq \|A\| \cdot \|B\|$  for any two matrices  $A$  and  $B$  on  $E$ . Further we obtain

$$\begin{aligned} \|P(t) - \tilde{P}(t)\| &= \left\| \int_0^t P(u) - \tilde{P}(u) \, du \, G \right\| \\ &\leq \int_0^t \|P(u) - \tilde{P}(u)\| \, du \cdot \|G\| \end{aligned} \quad (3.2)$$

$$\leq \Delta_t \cdot t \cdot \|G\| \quad (3.3)$$

with  $\Delta_t := \sup\{\|P(u) - \tilde{P}(u)\| : u \leq t\}$ , which is finite, since for all  $u \geq 0$  we know that  $\|P(u)\| = \|\tilde{P}(u)\| = 1$ . Plugging the result (3.3) into the right hand of the bound (3.2) again (with time  $u$  instead of  $t$ ), we obtain

$$\|P(t) - \tilde{P}(t)\| \leq \int_0^t \Delta_t \cdot u \cdot \|G\| \, du \cdot \|G\| = \Delta_t \cdot \frac{t^2}{2} \cdot \|G\|^2$$

Likewise,  $n$ -fold repetition of this step achieves the bound

$$\|P(t) - \tilde{P}(t)\| \leq \Delta_t \cdot \frac{t^n}{n!} \cdot \|G\|^n \leq \Delta_t \cdot \frac{(2\hat{\lambda} \cdot t)^n}{n!}$$

which in the limit  $n \rightarrow \infty$  yields  $0 \leq \|P(t) - \tilde{P}(t)\| \leq 0$  and consequently  $P(t) = \tilde{P}(t)$ . As  $t$  has been chosen arbitrarily, the statement is proven.

□

Hence the generator of a Markov process uniquely determines all its transition matrices. This can also be seen from the definition, if we agree (without loss of

generality) upon the convention  $p_{ii} = 0$  for all  $i \in E$ . Then the parameters for the definition of the Markov process can be recovered by

$$\lambda_i = -g_{ii} \quad \text{and} \quad p_{ij} = \frac{g_{ij}}{-g_{ii}}$$

for all  $i \neq j \in E$ .

However, as in the discrete time case of Markov chains, Markov processes are not completely determined by their transition probability matrices only. The missing link to a complete characterization again is given by the **initial distribution**  $\pi$  with  $\pi_i = \mathbb{P}(Y_0 = X_0 = i)$  for all  $i \in E$ . Then we can express all **finite-dimensional marginal distributions** as in

**Theorem 3.8** *For a Markov process  $\mathcal{Y}$  with initial distribution  $\pi$  and time instances  $0 < t_1 < \dots < t_n$ ,  $n \in \mathbb{N}$ , the equation*

$$\begin{aligned} \mathbb{P}(Y_{t_1} = j_1, \dots, Y_{t_n} = j_n) \\ = \sum_{i \in E} \pi_i P_{i,j_1}(t_1) P_{j_1,j_2}(t_2 - t_1) \dots P_{j_{n-1},j_n}(t_n - t_{n-1}) \end{aligned}$$

holds for all  $j_1, \dots, j_n \in E$ .

The proof is left as an exercise. Thus a Markov process  $\mathcal{Y}$  with transition probability matrices  $(P(t) : t \geq 0)$  admits a variety of versions depending on the initial distribution  $\pi$ . Any such **version** shall be denoted by  $\mathcal{Y}^\pi$ .

## 3.2 Stationary Distribution

From now on we shall convene on the technical assumption

$$\check{\lambda} := \inf\{\lambda_i : i \in E\} > 0$$

which holds for all queueing systems that we will examine. Then a Markov process is called **irreducible, transient, recurrent** or **positive recurrent** if the defining Markov chain is.

An initial distribution  $\pi$  is called **stationary** if the process  $\mathcal{Y}^\pi$  is stationary, i.e. if

$$\mathbb{P}(Y_{t_1}^\pi = j_1, \dots, Y_{t_n}^\pi = j_n) = \mathbb{P}(Y_{t_1+s}^\pi = j_1, \dots, Y_{t_n+s}^\pi = j_n)$$

for all  $n \in \mathbb{N}$ ,  $0 \leq t_1 < \dots < t_n$ , and states  $j_1, \dots, j_n \in E$ , and  $s \geq 0$ .

**Theorem 3.9** *A distribution  $\pi$  on  $E$  is stationary if and only if  $\pi G = 0$  holds.*

**Proof:** First we obtain

$$\pi P(t) = \pi e^{G \cdot t} = \sum_{n=0}^{\infty} \frac{t^n}{n!} \pi G^n = \pi I_E + \sum_{n=1}^{\infty} \frac{t^n}{n!} \pi G^n = \pi + \mathbf{0} = \pi$$

for all  $t \geq 0$ , with  $\mathbf{0}$  denoting the zero measure on  $E$ . With this, theorem 3.8 yields

$$\begin{aligned} \mathbb{P}(Y_{t_1}^\pi = j_1, \dots, Y_{t_n}^\pi = j_n) \\ &= \sum_{i \in E} \pi_i P_{i,j_1}(t_1) P_{j_1,j_2}(t_2 - t_1) \dots P_{j_{n-1},j_n}(t_n - t_{n-1}) \\ &= \pi_{j_1} P_{j_1,j_2}(t_2 - t_1) \dots P_{j_{n-1},j_n}(t_n - t_{n-1}) \\ &= \sum_{i \in E} \pi_i P_{i,j_1}(t_1 + s) P_{j_1,j_2}(t_2 - t_1) \dots P_{j_{n-1},j_n}(t_n - t_{n-1}) \\ &= \mathbb{P}(Y_{t_1+s}^\pi = j_1, \dots, Y_{t_n+s}^\pi = j_n) \end{aligned}$$

for all times  $t_1 < \dots < t_n$  with  $n \in \mathbb{N}$ , and states  $j_1, \dots, j_n \in E$ . Hence the process  $\mathcal{Y}^\pi$  is stationary.

On the other hand, if  $\pi$  is a stationary distribution, then we necessarily obtain  $\pi P(t) = \pi e^{G \cdot t} = \pi$  for all  $t \geq 0$ . As above, this means  $\sum_{n=1}^{\infty} \frac{t^n}{n!} \pi G^n = \mathbf{0}$  for all  $t \geq 0$ , which yields  $\pi G = \mathbf{0}$  because of the uniqueness of the zero power series.

□

By definition of the generator  $G$  and equation (3.1), the equation  $\pi G = 0$  is equivalent to an equation system

$$\sum_{i \neq j} \pi_i g_{ij} = -\pi_j g_{jj} \iff \sum_{i \neq j} \pi_i g_{ij} = \pi_j \sum_{i \neq j} g_{ji} \quad (3.4)$$

for all  $j \in E$ . This system can be interpreted as follows. We call the value  $\pi_i g_{ij}$  **stochastic flow** from state  $i$  to state  $j$  in equilibrium. Then the above equations mean that the accrued stochastic flow into any state  $j$  equals the flow out of this state. Equations (3.4) are called the (global) **balance equations**.

**Example 3.10** The generator of the Poisson process with parameter  $\lambda$  (see exam-

ple 3.1) is given by

$$G = \begin{pmatrix} -\lambda & \lambda & 0 & 0 & \dots \\ 0 & -\lambda & \lambda & 0 & \ddots \\ 0 & 0 & -\lambda & \lambda & \ddots \\ \vdots & \ddots & \ddots & \ddots & \ddots \end{pmatrix}$$

This process has no stationary distribution, which can be seen as follows. The balance equations for the Poisson process are given by

$$\pi_0 \lambda = 0 \quad \text{and} \quad \pi_i \lambda = \pi_{i-1} \lambda$$

for all  $i \geq 1$ . It is immediately evident that these are solvable only by  $\pi_i = 0$  for all  $i \in E$ , which means that there is no stationary distribution  $\pi$ .

The question of existence and uniqueness of a stationary distribution for  $\mathcal{Y}$  can be reduced to the same question for  $\mathcal{X}$ , which we have examined in the preceding chapter:

**Theorem 3.11** *Let the underlying Markov chain  $\mathcal{X}$  in the definition of the Markov process  $\mathcal{Y}$  be irreducible and positive recurrent. Further assume that  $\check{\lambda} := \inf\{\lambda_i : i \in E\} > 0$ . Then there is a unique stationary distribution for  $\mathcal{Y}$ .*

**Proof:** According to theorems 2.25 and 2.18, the transition matrix  $P$  of  $\mathcal{X}$  admits a unique stationary distribution  $\nu$  with  $\nu P = \nu$ . The generator  $G$  is defined by  $G = \Lambda(P - I_E)$ , with  $\Lambda = \text{diag}(\lambda_i : i \in E)$ . Hence the measure  $\mu := \nu \Lambda^{-1}$  is stationary for  $\mathcal{Y}$ . Since  $\check{\lambda} > 0$ , the measure  $\mu$  is finite, with total mass bounded by  $\check{\lambda}^{-1} < \infty$ . Now the normalization

$$\pi_j := \frac{\mu_j}{\sum_{i \in E} \mu_i} = \frac{\nu_j / \lambda_j}{\sum_{i \in E} \nu_i / \lambda_i} \quad (3.5)$$

for all  $j \in E$  yields a stationary distribution for  $\mathcal{Y}$ . This is unique because  $\nu$  is unique and the construction of  $\pi$  from  $\nu$  is reversible.

□

Finally we give two important results for the asymptotic behaviour of a Markov process. We call a Markov process **regular** if it satisfies the conditions given in the preceding theorem. If  $\mathcal{Y}$  is a regular Markov process, then the limit

$$\lim_{t \rightarrow \infty} \mathbb{P}(Y_t = j) = \pi_j \quad (3.6)$$

of the marginal distribution at time  $t$  tends to the stationary distribution as  $t$  tends to infinity. Further the limit

$$\lim_{t \rightarrow \infty} P_{ij}(t) = \pi_j \quad (3.7)$$

holds for all  $i, j \in E$  and is independent of  $i$ .

## Notes

An early text book on Markov processes with discrete state space is Chung [5]. Other classical text book presentation are Karlin and Taylor [14], Breiman [1], or Çinlar [4]. An exposition on non-homogeneous Markov processes on discrete state spaces can be found under the name Markov jump processes in Gikhman and Skorokhod [12, 11].

**Exercise 3.12** Consider a population of male and female species. There is an infinitesimal rate  $\lambda > 0$  that any male and female produce a single offspring, which will be female with probability  $p$ . Determine a Markov process which models the numbers  $F_t$  and  $M_t$  of female and male species at any time  $t$ .

**Exercise 3.13** Let  $X$  and  $Y$  denote two independent random variables which are distributed exponentially with parameters  $\lambda$  and  $\mu$ , respectively. Prove the following properties:

- (a)  $X \neq Y$  almost certainly.
- (b) The random variable  $Z := \min\{X, Y\}$  is distributed exponentially with parameter  $\lambda + \mu$ .
- (c)  $\mathbb{P}(X < Y) = \lambda/(\lambda + \mu)$

**Exercise 3.14** Let  $\mathcal{Y}^{(1)}$  and  $\mathcal{Y}^{(2)}$  denote independent Poisson processes with intensities  $\lambda_1$  and  $\lambda_2$ , respectively. Show that the process  $\mathcal{Y} = (Y_t : t \in \mathbb{R}_0^+)$  defined by  $Y_t = Y_t^{(1)} + Y_t^{(2)}$  for all  $t \geq 0$  is a Poisson process with intensity  $\lambda = \lambda_1 + \lambda_2$ . The process  $\mathcal{Y}$  is called the **superposition** of  $\mathcal{Y}^{(1)}$  and  $\mathcal{Y}^{(2)}$ .

**Exercise 3.15** Prove theorem 3.8.

**Exercise 3.16** Determine the finite-dimensional marginal distributions for a Poisson process with parameter  $\lambda$ .

**Exercise 3.17** Let  $\mathcal{Y}$  denote a Poisson process with parameter  $\lambda$ . Given that there is exactly one arrival in the interval  $[0, t]$ , show that the exact time of the arrival within  $[0, t]$  is uniformly distributed.

**Exercise 3.18** Verify the Chapman–Kolmogorov equations for a Poisson process.

### 3.3 Skip–Free Markov Processes and Markovian Queues

The methods of analyzing Markov processes are already sufficient for the treatment of quite a variety of queueing systems. These are commonly known as elementary or **Markovian queues**. The most classical of them shall be examined in this chapter.

#### 3.3.1 The M/M/1 Queue

The M/M/1 queue in continuous time is defined by the following characteristics: The arrival process is a Poisson process with some rate  $\lambda > 0$ . The service times are iid and distributed exponentially with service rate  $\mu > 0$ . There is one server and the service discipline is first come first served.




Figure 3.2: M/M/1 queue

For the Poisson process, the inter–arrival times are distributed exponentially with parameter  $\lambda$ . Since the exponential distribution is memoryless, the system process  $\mathcal{Q} = (Q_t : t \in \mathbb{R}_0^+)$  can be modelled by a Markov process with state space  $E = \mathbb{N}_0$  and generator

$$G = \begin{pmatrix} -\lambda & \lambda & 0 & 0 & \dots \\ \mu & -\lambda - \mu & \lambda & 0 & \ddots \\ 0 & \mu & -\lambda - \mu & \lambda & \ddots \\ \vdots & \ddots & \ddots & \ddots & \ddots \end{pmatrix}$$

Here, the first line represents the possible transitions if the system is empty. In this case there can only occur single arrivals according to the Poisson process with rate  $\lambda$ . If the system is not empty, there are two possibilities: Either an arrival occurs (with rate  $\lambda$ ) or a service is completed (with rate  $\mu$ ). Contrary to the M/M/1 queue



in discrete time, arrivals and service completions cannot occur at the same time. This follows from the memoryless property of the exponential distribution and exercise 3.13. The parameter of the holding time for the states of a non-empty system is explained by exercise 3.13.

Clearly, the structure of the matrix  $G$  shows that the process  $Q$  is irreducible and hence there is at most one stationary distribution  $\pi$  for  $Q$ . According to theorem 3.9, this must satisfy  $\pi G = 0$ , which translates into the system

$$\pi_0 \lambda = \pi_1 \mu \quad (3.8)$$

$$\pi_n (\lambda + \mu) = \pi_{n-1} \lambda + \pi_{n+1} \mu \quad \text{for all } n \geq 1 \quad (3.9)$$

$$\sum_{n=0}^{\infty} \pi_n = 1 \quad (3.10)$$

of equations, where the latter is simply the normalization of the distribution  $\pi$ . The first two equations are the global **balance equations** and can be illustrated by the following scheme:




Figure 3.3: Transition rates for the M/M/1 queue

This gives the rates of jumps between the states of the system. If we encircle any one state, then the sum of the rates belonging to the arcs reaching into this state must equal the sum of the rates which belong to the arcs that go out of this state. If this is the case, then we say that the system is in balance. The conditions for this are given in equations (3.8) and (3.9).

The solution of the above system of equations can be obtained by the following considerations: The first equation yields

$$\pi_1 = \pi_0 \frac{\lambda}{\mu} =: \pi_0 \rho$$

with  $\rho := \lambda/\mu$ . By induction on  $n$  we obtain from the second equation

$$\begin{aligned} \pi_{n+1} &= \frac{1}{\mu} (\pi_n (\lambda + \mu) - \pi_{n-1} \lambda) = \pi_n \frac{\lambda}{\mu} + \pi_n - \pi_{n-1} \frac{\lambda}{\mu} \\ &= \pi_n \rho \end{aligned}$$

for all  $n \in \mathbb{N}$ , where the last equality holds by induction hypothesis. Thus the geometric approach  $\pi_n = \pi_0 \rho^n$  for all  $n \in \mathbb{N}_0$  solves the first two equations. The last equation now yields

$$1 = \sum_{n=0}^{\infty} \pi_n = \pi_0 \sum_{n=0}^{\infty} \rho^n = \frac{1}{1-\rho} \pi_0$$

if and only if  $\rho < 1$ , which means  $\lambda < \mu$ . Hence there is a stationary distribution of the system, given by

$$\pi_n = (1-\rho)\rho^n$$

for all  $n \in \mathbb{N}_0$ , if and only if the so-called queue **load**  $\rho = \lambda/\mu$  remains smaller than one.

In this case several performance measures of the queueing system can be derived immediately. All of them are computed by means of the stationary distribution. Thus they hold only for the system being in equilibrium, which is attained asymptotically.

For instance, the probability that the system is empty is given by  $\pi_0 = 1 - \rho$ . The mean and the variance of the number  $N$  of users in the system are given as

$$\mathbb{E}(N) = \sum_{n=1}^{\infty} n\pi_n = (1-\rho) \sum_{n=1}^{\infty} n\rho^n = \frac{\rho}{1-\rho}$$

and  $\text{Var}(N) = \rho/(1-\rho)^2$ . The probability  $R_K$  that there are at least  $K$  users in the system is

$$R_K = \sum_{n=K}^{\infty} \pi_n = (1-\rho) \sum_{n=K}^{\infty} \rho^n = \rho^K$$

As expected, these equations show that with increasing load  $\rho \rightarrow 1$  the mean number of users in the system grows and the probability of an idle system decreases.

### 3.3.2 Skip-Free Markov Processes

There are many variations of the M/M/1 queue which can be analyzed by the same method. In order to show this we first put the analysis presented in the preceding section in a more general context. This will be applicable to a large variety of queueing models.

The Markov process which models the M/M/1 queue has the decisive property that transitions are allowed to neighbouring states only, i.e.  $g_{ij} = 0$  for states

$i, j \in \mathbb{N}_0$  with  $|i - j| > 1$ . The result is a very simple state transition graph of a linear form and correspondingly a set of balance equations, given by (3.8) and (3.9), which can be solved easily. We can retain the same method of analysis if we relax the special assumption that  $g_{i,i+1}$  and  $g_{i,i-1}$  be independent of  $i$ .

Thus we define a **skip-free Markov process** by the property that its generator  $G = (g_{ij})_{i,j \in E}$  satisfies  $g_{ij} = 0$  for all states  $i, j \in E \subset \mathbb{N}_0$  with  $|i - j| > 1$ . For queueing systems this means that there are only single arrivals or departures. Thus every Markovian queueing system with single arrivals and departures can be modelled by a skip-free Markov process.

Denote the remaining infinitesimal transition rates by

$$\lambda_i := g_{i,i+1} \quad \text{and} \quad \mu_i := g_{i,i-1}$$

for all possible values of  $i$ . The rates  $\lambda_i$  and  $\mu_i$  are called **arrival rates** and **departure rates**, respectively. The state transition graph of such a process assumes the form




Figure 3.4: A skip-free Markov process

Its balance equations are given by  $\lambda_0\pi_0 = \mu_1\pi_1$  and

$$(\lambda_i + \mu_i)\pi_i = \lambda_{i-1}\pi_{i-1} + \mu_{i+1}\pi_{i+1}$$

for all  $i \in \mathbb{N}$ . By induction on  $i$  it is easily shown that these are equivalent to the equation system

$$\lambda_{i-1}\pi_{i-1} = \mu_i\pi_i \quad (3.11)$$

for all  $i \in \mathbb{N}$ . This system is solved by successive elimination with a solution of the form

$$\pi_i = \pi_0 \prod_{j=0}^{i-1} \frac{\lambda_j}{\mu_{j+1}} = \pi_0 \frac{\lambda_0 \lambda_1 \cdots \lambda_{i-1}}{\mu_1 \mu_2 \cdots \mu_i} \quad (3.12)$$

for all  $i \geq 1$ . The solution  $\pi$  is a probability distribution if and only if it can be normalized, i.e. if  $\sum_{n \in E} \pi_n = 1$ . This condition implies

$$1 = \sum_{n \in E} \pi_0 \prod_{j=0}^{n-1} \frac{\lambda_j}{\mu_{j+1}} = \pi_0 \sum_{n \in E} \prod_{j=0}^{n-1} \frac{\lambda_j}{\mu_{j+1}}$$

with the empty product being defined as one. This means that

$$\pi_0 = \left( \sum_{n \in E} \prod_{j=0}^{n-1} \frac{\lambda_j}{\mu_{j+1}} \right)^{-1} \quad (3.13)$$

and thus  $\pi$  is a probability distribution if and only if the series in the brackets converges. In this case, the stationary distribution of a skip-free Markov process is given by (3.13) and (3.12).

### 3.3.3 The M/M/ $\infty$ Queue

The first application of the analysis of the last section to a queueing system shall be the M/M/ $\infty$  queue. This is a queue without queueing: There are infinitely many servers such that every incoming user finds an idle server immediately. Arrivals are governed by a Poisson process with intensity  $\lambda > 0$ , and the service times are exponentially distributed with rate  $\mu > 0$ , equal for each server. Due to lemma 3.13, the system process is Markovian. Furthermore, there are only single arrivals and departures. Hence the M/M/ $\infty$  queue can be modelled by a skip-free Markov process.

Since the arrival process is independent of the rest of the queue, the arrival rates of the respective skip-free Markov process are constant. In the notation of section 3.3.2 we can thus specify  $\lambda_n = \lambda$  for all  $n \in \mathbb{N}_0$ . Departures occur upon service completions. According to lemma 3.13 and due to the memoryless property of the exponential distribution (see lemma 3.2), the departure rates are given by  $\mu_n = n \cdot \mu$  for all  $n \in \mathbb{N}$ .

Define  $\rho := \lambda/\mu$ . Then the series in (3.13) assumes the value

$$\sum_{n=0}^{\infty} \prod_{j=0}^{n-1} \frac{\lambda_j}{\mu_{j+1}} = \sum_{n=0}^{\infty} \frac{\rho^n}{n!} = e^\rho$$

and thus converges regardless of the value of  $\rho$ . This means that the M/M/ $\infty$  queue always has a stationary distribution, which is not surprising as infinitely many servers cannot be exhausted, whatever the arrival intensity amounts to.

Due to formulae (3.13) and (3.12), we obtain the stationary distribution  $\pi$  as given by  $\pi_0 = e^{-\rho}$  and

$$\pi_n = e^{-\rho} \cdot \frac{\rho^n}{n!}$$

for all  $n \in \mathbb{N}$ , which is a Poisson distribution with parameter  $\rho$ . Hence the mean and the variance of the number  $N$  of users in the stationary system are given by  $\mathbb{E}(N) = \mathbb{V}ar(N) = \rho$ .

Since there is no queueing in the M/M/ $\infty$  system, all waiting times are zero and the mean sojourn time in the system equals  $1/\mu$ . This means that all users passing through such a system are independently kept there for an exponentially distributed time. In the context of queueing networks, the M/M/ $\infty$  queue is therefore often called an (independent) **delay system**.

### 3.3.4 The M/M/k Queue

The M/M/k queue is provided with  $k$  identical servers which can serve users in parallel. Users arrive according to a Poisson process with intensity  $\lambda > 0$ , and the service time distribution is exponential with parameter  $\mu > 0$  at all servers. Whenever a user arrives and finds all servers busy (i.e. at least  $k$  users in the system) he queues up in the waiting room. From there the next waiting user is served in the order of a FIFO discipline as soon as one of the servers becomes idle. An arriving user finding less than  $k$  users already in the system (i.e. there are idle servers at the time of arrival) chooses any server and starts service immediately.

For this type of queue the dynamics is a mixture between the M/M/ $\infty$  queue and the M/M/1 queue. Up to the value of  $k$  users in the system, the service (and thus the departure) rate increases like  $\mu_n = n \cdot \mu$  for  $1 \leq n \leq k$ . Starting from  $k$  users in the system there are no servers anymore to keep up with newly arriving users, and the departure rate remains  $\mu_n = k \cdot \mu$  for all  $n \geq k + 1$ . The independence of the arrival process yields constant arrival rates  $\lambda_n = \lambda$  for all  $n \in \mathbb{N}_0$ .

Again we define  $\rho := \lambda/\mu$ . The series in (3.13) specifies to

$$\sum_{n=0}^{\infty} \prod_{j=0}^{n-1} \frac{\lambda_j}{\mu_{j+1}} = \sum_{n=0}^{k-1} \frac{\rho^n}{n!} + \frac{\rho^k}{k!} \sum_{n=0}^{\infty} \left(\frac{\rho}{k}\right)^n$$

which is finite if and only if  $\rho < k$ . In this case the stationary distribution  $\pi$  is




Figure 3.5: M/M/k queue

given by formulae (3.13) and (3.12) as

$$\pi_0 = \left( \sum_{n=0}^{k-1} \frac{\rho^n}{n!} + \frac{\rho^k}{(k-1)! \cdot (k-\rho)} \right)^{-1}$$

and

$$\begin{aligned} \pi_n &= \pi_0 \cdot \frac{\rho^n}{n!}, & 1 \leq n \leq k \\ \pi_n &= \pi_k \cdot \left( \frac{\rho}{k} \right)^{n-k}, & n > k \end{aligned}$$

Here we see the M/M/ $\infty$  form for  $n \leq k$  and the M/M/1 form beginning with  $n \geq k$ , where  $\pi_k$  substitutes the base value that is played by  $\pi_0$  for the pure M/M/1 queue.

The fact that the M/M/k queue behaves for more than  $k$  users in the system like an M/M/1 queue with load  $\rho/k$  is further illustrated by the following observation. Let  $N$  denote the number of users in the system that is in equilibrium. Consider the conditional probability  $p_n := \mathbb{P}(N = n | N \geq k)$  for  $n \geq k$ . This is computed as

$$p_n = \frac{\pi_n}{\sum_{i=k}^{\infty} \pi_i} = \pi_k \left( \frac{\rho}{k} \right)^{n-k} \bigg/ \pi_k \sum_{i=k}^{\infty} \left( \frac{\rho}{k} \right)^{i-k} = \left( \frac{\rho}{k} \right)^{n-k} \left( 1 - \frac{\rho}{k} \right)$$

Since  $n - k$  is the number  $N_q$  of users waiting in the queue, the conditional distribution of  $N_q$  given that all servers are busy has exactly the same (geometric) form as the stationary distribution for the M/M/1 system process.

The probability  $\mathbb{P}\{N \geq k\}$  of the conditioning event that in equilibrium all servers are busy is given by

$$\sum_{n=k}^{\infty} \pi_n = \left( 1 + (k-1)! \cdot (k-\rho) \cdot \sum_{n=0}^{k-1} \frac{\rho^{n-k}}{n!} \right)^{-1} \quad (3.14)$$

This is the probability that a newly arriving user must wait before he is served. The above formula for it is called **Erlang's delay formula**.

### 3.3.5 The M/M/k/k Queue

In stochastic modelling there always is a trade-off between the adaptation of the model to reality and its simplicity, i.e. its analytic tractability. We have seen that the nicest solutions could be derived for the M/M/1 queue (a geometric distribution) and the M/M/ $\infty$  queue (a Poisson distribution). The solution for the M/M/k queue, which is more realistic for most practical applications, is also more involved. For all these models we kept the often unrealistic assumption of an infinite waiting room. The models in this and the following sections stem from more realistic specifications. Historically, they belong to the first applications which founded the field of queueing theory.

In the times of A.K. Erlang, at the beginning of the 20th century, telephone calls had to be connected by an operator. The telephone companies installed call centers where a number  $k$  of operators served call requests which arrived from a large number of subscribers. Whenever all operators are busy with serving call requests and a new subscriber calls to get a line, this subscriber will be rejected.

If we model the arriving requests by a Poisson process and the duration of the operators' services by an exponential distribution, then we get an M/M/k/k queue as a model for this application. The subscribers with their call requests are the users and the operators are the servers. There are  $k$  servers and as many places in the system, i.e. there is no additional waiting room.

Let the intensity of the Poisson arrival process and the rate of the exponential service times be denoted by  $\lambda > 0$  and  $\mu > 0$ , respectively. Again we can use a skip-free Markov process to analyze this system. In this notation, we obtain  $\lambda_n = \lambda$  for all  $n = 0, \dots, k-1$  and  $\mu_n = n \cdot \mu$  for  $n = 1, \dots, k$ . The values of

$\lambda_n$  and  $\mu_n$  are zero for all other indices  $n$ . Define  $\rho := \lambda/\mu$ . The series in (3.13) is in this case

$$\sum_{n \in E} \prod_{j=0}^{n-1} \frac{\lambda_j}{\mu_{j+1}} = \sum_{n=0}^k \frac{\rho^n}{n!}$$

which is finite, regardless of the value for  $\rho$ . Hence a stationary distribution  $\pi$  always exists and is given by

$$\pi_0 = \left( \sum_{n=0}^k \frac{\rho^n}{n!} \right)^{-1} \quad \text{and} \quad \pi_n = \pi_0 \cdot \frac{\rho^n}{n!}$$

for all  $n = 1, \dots, k$ . The main performance measure for this application is the probability that all operators are busy and the company is unable to accept new call requests. This is given by

$$\pi_k = \frac{\rho^k}{k!} \left( \sum_{n=0}^k \frac{\rho^n}{n!} \right)^{-1}$$

which of course is valid only under the stationary regime, i.e. in equilibrium. This expression is known as **Erlang's loss formula**.

Note that the expression of  $\pi_0$  for the M/M/ $\infty$  queue is the limit of the respective expression for the M/M/k/k model as  $k$  tends to infinity. Even further, the stationary distribution for the M/M/k/k queue converges to the stationary distribution of the M/M/ $\infty$  for increasing  $k$ .

### 3.3.6 The M/M/k/k+c/N Queue

A simplifying assumption in the previous model has been the constant arrival rates  $\lambda_n = \lambda$ . This implies that even for a high number of users in the queue the intensity of new arrivals does not diminish. While this is a reasonable assumption for an application to call centers, where the number of operators (and thus the maximal number of users in the system) is only marginal compared to the number of all subscribers, there are other applications for which such an assumption would not be realistic.

Consider a closed computer network with  $k$  servers and  $N$  terminals. Every terminal sends a job to one of the servers after some exponentially distributed think time. If a server is available, i.e. idle, then this job is served, demanding an exponential service time. A terminal that has a job in a server may not send another job






Figure 3.6: A closed computer network

request during the service time. Whenever a terminal sends a job request and all servers are busy at that time, then the job is put into a queue. This queue has maximal capacity  $c$ , i.e. if a terminal sends a job request and the queue is already filled with  $c$  jobs, then this new job request is rejected and the terminal starts another think time.

This application can be modelled by an M/M/k/k+c/N queue if we interpret the users in the system as the job requests that are in service or waiting. Denote the parameters of the exponential think time and service time distributions by  $\lambda > 0$  and  $\mu > 0$ , respectively. Without loss of generality we may assume that  $k+c \leq N$ . Then the queue in consideration is a skip-free Markov process with arrival rates  $\lambda_n = (N-n) \cdot \lambda$  for  $n = 0, \dots, k+c-1$  and departure rates  $\mu_n = \min(n, k) \cdot \mu$  for  $n = 1, \dots, k+c$ . As always, define  $\rho := \lambda/\mu$ . The series in (3.13) amounts to

$$\sum_{n \in E} \prod_{j=0}^{n-1} \frac{\lambda_j}{\mu_{j+1}} = \sum_{n=0}^k \binom{N}{n} \cdot \rho^n + \sum_{n=k+1}^{k+c} \frac{N! \cdot \rho^n}{(N-n)! \cdot k! \cdot k^{n-k}} \quad (3.15)$$

and thus is finite for every value of  $\rho$ . The stationary distribution  $\pi$  is given by

$$\pi_0 = \left( \sum_{n=0}^k \binom{N}{n} \cdot \rho^n + \sum_{n=k+1}^{k+c} \frac{N! \cdot \rho^n}{(N-n)! \cdot k! \cdot k^{n-k}} \right)^{-1}$$

and

$$\pi_n = \pi_0 \cdot \binom{N}{n} \cdot \rho^n, \quad 1 \leq n \leq k$$

$$\pi_n = \pi_0 \cdot \frac{N! \cdot \rho^n}{(N-n)! \cdot k! \cdot k^{n-k}}, \quad k+1 \leq n \leq k+c$$

There are several interesting special cases. For  $c = 0$  there is no room for a queue of waiting jobs. Then the stationary distribution simplifies to a binomial distribution with parameters  $(N, p)$ , where  $p = \rho/(1 + \rho)$ , which is truncated to the states  $n = 0, \dots, k$ . Such a distribution is called an **Engset distribution**.

For  $c = N - k$  the queue has an important application in reliability theory. This is known as the **machine repair problem**. In a production site there are  $N$  machines which are prone to failure. Each of them breaks down after a working time which is exponentially distributed with parameter  $\lambda$ . There are  $k$  repairmen that take care of the broken machines sequentially. The repair times are exponential with parameter  $\mu$ . Then the system process of the  $M/M/k/N/N$  queue yields the number of broken machines.

## Notes

The models presented in this chapter are the oldest within queueing theory. Applications to telephone networks date back to the beginning of the 20th century, notably Erlang [7] and Engset [6].

Skip-free Markov processes have been extensively used for populations models. Therefore the name birth-and-death processes is very popular for them, with  $\lambda_i$  and  $\mu_i$  denoting the transition rates for a birth and a death, respectively, if the population has  $i$  members. However, the authors think that such a name is inappropriate for queueing models and thus prefer the more technical term skip-free. For more Markovian queueing models see Kleinrock [15]. An analysis of non-homogeneous (namely periodic) Markovian queues is given in Breuer [2, 3].

**Exercise 3.19** Verify the formula  $\text{Var}(N) = \rho/(1 - \rho)^2$  for the stationary variance of the number of users in the M/M/1 queue.

**Exercise 3.20** Show that the equation system (3.11) is equivalent to the balance equations for a skip-free Markov process. Prove the form (3.12) of its solution.

**Exercise 3.21** Prove Erlang's delay formula (3.14).

**Exercise 3.22** Compare the stationary mean number of users in the system for the following three queueing systems: (a) an M/M/1 queue with arrival intensity  $\lambda$  and service rate  $\mu$ , (b) an M/M/2 system with arrival intensity  $\lambda$  and service rate  $\mu/2$ , and (c) two independent M/M/1 queues with arrival intensity  $\lambda/2$  to each of them and equal service rate  $\mu$ . Explain the differences.

**Exercise 3.23** Explain equation (3.15).

**Exercise 3.24** Show that the stationary distribution for an  $M/M/k/k/N$  queue is an Engset distribution.

**Exercise 3.25** Analyze the M/M/1/c queue with arrival intensity  $\lambda$  and service rate  $\mu$ . This always has a stationary distribution  $\pi$ . Show that in the limit  $c \rightarrow \infty$ , there are two possibilities: Either  $\rho < 1$  and  $\pi$  converges to the stationary distribution of the M/M/1 queue, or  $\rho \geq 1$  and  $\pi$  converges to the zero measure.

**Exercise 3.26** Examine the M/M/1 queue with users who are discouraged by long queue lengths. This can be modelled by arrival rates  $\lambda_n = \lambda/(n+1)$  for all  $n \in \mathbb{N}_0$ . Show that the stationary distribution is Poisson.



# Chapter 4

## Renewal Theory

### 4.1 Renewal Processes

Be  $(X_n : n \in \mathbb{N}_0)$  a sequence of independent positive random variables, and assume that  $(X_n : n \in \mathbb{N})$  are identically distributed. Define the sequence  $\mathcal{S} = (S_n : n \in \mathbb{N})$  by  $S_1 := X_0$  and  $S_{n+1} := S_n + X_n$  for all  $n \in \mathbb{N}$ . The random variable  $S_n$ , with  $n \in \mathbb{N}$ , is called the  $n$ th **renewal time**, while the time duration  $X_n$  is called the  $n$ th **renewal interval**. Further define the random variable of the number of renewals until time  $t$  by

$$N_t := \max\{n \in \mathbb{N} : S_n \leq t\}$$

for all  $t \geq 0$  with the convention  $\max \emptyset = 0$ . Then the continuous time process  $\mathcal{N} = (N_t : t \in \mathbb{R}_0^+)$  is called a **renewal process**. The random variable  $X_0$  is called the **delay** of  $\mathcal{N}$ . If  $X_0$  and  $X_1$  have the same distribution, then  $\mathcal{N}$  is called an **ordinary renewal process**.

We will always assume that  $\mathbb{P}(X_1 = 0) = 0$  and  $m := \mathbb{E}(X_1) < \infty$  is finite. The strong law of large numbers implies that  $S_n/n \rightarrow m$  with probability one as  $n \rightarrow \infty$ . Hence  $S_n < t$  cannot hold for infinitely many  $n$  and thus  $N_t$  is finite with probability one. By standard notation we will write

$$G(x) := \mathbb{P}(X_0 \leq x) \quad \text{and} \quad F(x) := \mathbb{P}(X_1 \leq x)$$

for all  $x \in \mathbb{R}_0^+$ .

**Example 4.1** A light bulb has been installed at time zero. After a duration  $X_0$ , it will go out of order. We assume that it will be immediately replaced by a new light

bulb at time  $S_1 = X_0$ . Assume that the new light bulb is of a type identical to the old one. Then the duration  $X_1$  until it goes out of order is distributed identically to  $X_0$ . Of course, the life times of the light bulbs are independent from one another. Keeping up this rechange policy over time, the number  $N_t$  of used light bulbs until time  $t$  forms an ordinary renewal process.

**Remark 4.2** A Poisson process with intensity  $\lambda$  (see example 3.1) is an ordinary renewal process with  $F(x) = G(x) = 1 - e^{-\lambda x}$ , i.e. the renewal intervals have an exponential distribution. Thus a renewal process can be seen as a generalization of the Poisson process with respect to the distribution of the renewal intervals.

In order to derive an expression for the distribution and the expectation of  $N_t$  at any time  $t$ , we need to introduce the concept of **convolutions** of a non-negative function and a distribution function. Let  $F$  denote a distribution function on  $\mathbb{R}_0^+$  and  $g : \mathbb{R}_0^+ \rightarrow \mathbb{R}_0^+$  a Lebesgue-measurable function which is bounded on all finite intervals  $[0, t]$  with  $t \geq 0$ . Then the function defined by

$$F * g(t) := \int_0^t g(t-u) dF(u)$$

for all  $t \in \mathbb{R}$  is called the convolution of  $F$  and  $g$ . In particular, the definition of a convolution applies if  $g$  is a distribution function. As an exercise the reader can prove

**Theorem 4.3** For any distribution functions  $F$  and  $G$  as well as non-negative Lebesgue-measurable functions  $(g_n : n \in \mathbb{N})$  on  $\mathbb{R}_0^+$ , the following properties hold:

- (1) The convolution  $F * G$  is a distribution function on  $\mathbb{R}_0^+$ .
- (2)  $F * G = G * F$
- (3)  $F * \sum_{n=1}^{\infty} g_n = \sum_{n=1}^{\infty} F * g_n$
- (4) The Dirac measure  $\delta_0$  on 0 with distribution function  $I_0$ , which is defined by  $I_0(t) := 1$  for all  $t \geq 0$  and  $I_0(t) := 0$  otherwise, is neutral in regard to convolutions, i.e.  $I_0 * G = G$  for all distribution functions  $G$ .
- (5) If the random variables  $X$  and  $Y$  are independent and distributed according to  $F$  and  $G$ , respectively, then  $\mathbb{P}(X + Y \leq t) = F * G(t)$  for all  $t \geq 0$ .
- (6)  $F * (G * g) = (F * G) * g$

Let  $F$  denote any distribution function for a real-valued random variable. Define the **convolutional powers** by  $F^{*1} := F$  and recursively  $F^{*n+1} := F^{*n} * F$  for all

$n \in \mathbb{N}$ . Because of property (4) in the above theorem, we define  $F^{*0} := I_0$  for every distribution function  $F$ .

Now denote the distribution function of the random variable  $X_1$  (and hence of all  $X_n$  with  $n \geq 1$ ) and  $X_0$  by  $F$  and  $G$ , respectively. Since the random variables  $(X_n : n \in \mathbb{N})$  are iid, part (5) of the above theorem yields for all  $n \in \mathbb{N}_0$  the relation  $\mathbb{P}(N_t \geq n) = \mathbb{P}(S_n \leq t) = G * F^{*n-1}(t)$  and thus we obtain  $\mathbb{P}(N_t = 0) = 1 - G(t)$  and

$$\mathbb{P}(N_t = n) = \mathbb{P}(S_n \leq t) - \mathbb{P}(S_{n+1} \leq t) = G * F^{*n-1}(t) - G * F^{*n}(t)$$

for  $n \geq 1$ . The expectation of  $N_t$  is given by

$$\mathbb{E}(N_t) = \sum_{n=1}^{\infty} \mathbb{P}(N_t \geq n) = \sum_{n=1}^{\infty} \mathbb{P}(S_n \leq t) = G * \sum_{n=0}^{\infty} F^{*n}(t) \quad (4.1)$$

for all  $t \geq 0$  (for the first equality see Exercise 4.19). The rate of growth of a renewal process is described by

**Theorem 4.4** *Let  $\mathcal{N} = (N_t : t \geq 0)$  denote a renewal process with renewal intervals having mean length  $m < \infty$ . Then*

$$\lim_{t \rightarrow \infty} \frac{N_t}{t} = \frac{1}{m}$$

*holds with probability one.*

**Proof:** By definition of  $N_t$ , the inequalities  $S_{N_t} \leq t \leq S_{N_t+1}$  hold with probability one for all times  $t$ . Dividing these by  $N_t$  and using the strong law of large numbers, we obtain

$$\begin{aligned} m &= \lim_{n \rightarrow \infty} \frac{S_n}{n} = \lim_{t \rightarrow \infty} \frac{S_{N_t}}{N_t} \\ &\leq \lim_{t \rightarrow \infty} \frac{t}{N_t} \\ &\leq \lim_{t \rightarrow \infty} \left( \frac{S_{N_t+1}}{N_t+1} \cdot \frac{N_t+1}{N_t} \right) = \lim_{n \rightarrow \infty} \frac{S_{n+1}}{n+1} \cdot \lim_{n \rightarrow \infty} \frac{n+1}{n} = m \cdot 1 \end{aligned}$$

which proves the statement.

□

Because of this theorem, the inverse  $1/m$  of the mean length of a renewal interval is called the **rate** of the renewal process. It describes the asymptotic rate at which renewals occur.

**Example 4.5** Regarding a Poisson process  $\mathcal{N} = (N_t : t \geq 0)$  with intensity  $\lambda > 0$ , it can be shown that

$$\mathbb{P}(N_t = n) = \frac{(\lambda t)^n}{n!} e^{-\lambda t} \quad (4.2)$$

for all  $t \geq 0$  and  $n \in \mathbb{N}_0$ . The expectation of  $N_t$  is given by  $\mathbb{E}(N_t) = \lambda \cdot t$ .

Thus a Poisson process with intensity  $\lambda$  has at time  $t$  a Poisson distribution with parameter  $\lambda \cdot t$ . Moreover, the intensity  $\lambda$  is also the rate of the Poisson process, since a mean renewal interval has length  $1/\lambda$ .

Given an observed stream of events (e.g. job requests at a server) over some time interval of length  $t$ , we can count the number  $N(t)$  of events that have occurred in this interval. If we want to model such event streams by a Poisson process, then we need to find a statistical estimator for the intensity  $\lambda$ . Now theorem 4.4 states that the fraction  $N(t)/t$  comes close to  $\lambda$  for large interval lengths  $t$ . Thus a consistent statistical estimator for the intensity  $\lambda$  is given by  $\hat{\lambda} = N(t)/t$ .

**Example 4.6** There is a discrete-time analogue of the Poisson process, which is called **Bernoulli process**. This is an ordinary renewal process with renewal intervals that have a geometric distribution. Given a parameter  $p \in ]0, 1[$ , the length of the renewal intervals is distributed as  $\mathbb{P}(X_1 = n) = p \cdot (1 - p)^{n-1}$  for  $n \in \mathbb{N}$ .

## 4.2 Renewal Function and Renewal Equations

The function defined by  $R(t) := \sum_{n=1}^{\infty} F^{*n}(t)$  for all  $t \geq 0$  is called the **renewal function** of the process  $\mathcal{N}$ . The renewal function will play a central role in renewal theory. First we need to show that it remains finite:

**Theorem 4.7** *If  $F(0) < 1$ , then  $R(t) = \sum_{n=1}^{\infty} F^{*n}(t) < \infty$  for all  $t \geq 0$ .*

**Proof:** Since  $F(0) < 1$  and  $F$  is continuous to the right, there is a number  $\alpha > 0$  such that  $F(\alpha) < 1$ . Fix any  $t \geq 0$  and choose  $k \in \mathbb{N}$  such that  $k \cdot \alpha > t$ . Then  $F^{*k}(t) \leq 1 - (1 - F(\alpha))^k =: 1 - \beta$  with  $0 < \beta < 1$ . Thence we obtain the bound  $F^{*mk}(t) \leq (1 - \beta)^m$  for any  $m \in \mathbb{N}$ . Since  $F(0-) = 0$ , we can use  $F^{*n}(t) \geq F^{*h}(t)$  for all  $n < h \in \mathbb{N}$ . Putting these bounds together, we obtain

$$R(t) = \sum_{n=1}^{\infty} F^{*n}(t) \leq k \cdot \sum_{m=0}^{\infty} F^{*mk}(t) \leq k \cdot \sum_{m=0}^{\infty} (1 - \beta)^m = \frac{k}{\beta} < \infty$$



since  $\beta > 0$ .

□

**Theorem 4.8** *An ordinary renewal process is uniquely determined by its renewal function.*

**Proof:** First we take the Laplace–Stieltjes transform (LST, see appendix 5.3) on both sides of the equation  $R(t) = \sum_{n=1}^{\infty} F^{*n}(t)$ . This yields

$$\tilde{R}(s) = \sum_{n=1}^{\infty} \widetilde{F^{*n}}(s) = \tilde{F}(s) \cdot \sum_{n=0}^{\infty} (\tilde{F}(s))^n = \frac{\tilde{F}(s)}{1 - \tilde{F}(s)} \quad (4.3)$$

for  $s > 0$ , or

$$\tilde{F}(s) = \frac{\tilde{R}(s)}{1 + \tilde{R}(s)}$$

and thus determines the LST  $\tilde{F}(s)$  of  $F$  uniquely in terms of  $\tilde{R}(s)$ . Now uniqueness of the LST yields the statement.

□

For an ordinary renewal process we can derive an implicit integral equation for the renewal function, which is known as a renewal equation. Note that for an ordinary renewal process  $\mathbb{E}(N_t) = R(t)$  for all times  $t$  (see (4.1) with  $G = F$ ). Hence the function  $R$  is increasing. If we condition upon the length  $x$  of the first renewal interval  $X_0$ , we obtain

$$\mathbb{E}(N_t) = \int_0^{\infty} \mathbb{E}(N_t | X_0 = x) dF(x)$$

Since  $\mathbb{E}(N_t | X_0 = x) = 1 + R(t - x)$  for  $t \geq x$  and  $\mathbb{E}(N_t | X_0 = x) = 0$  for  $t < x$ , we can simplify this equation to

$$R(t) = \int_0^t (1 + R(t - x)) dF(x) = F(t) + \int_0^t R(t - x) dF(x)$$

for all  $t \geq 0$ . A **renewal equation** is the generalized form

$$g(t) = h(t) + \int_0^t g(t - x) dF(x), \quad t \geq 0 \quad (4.4)$$

where a function  $h$  on  $[0, \infty[$  and a distribution function  $F$  on  $[0, \infty[$  are given and the function  $g$  on  $[0, \infty[$  is unknown. The solution is given in

**Theorem 4.9** *The unique solution  $g$  to equation (4.4) is given by*

$$g(t) = \int_0^t h(t-x) dR(t) + h(t)$$

where  $R(t) = \sum_{n=1}^{\infty} F^{*n}(t)$  denotes the renewal function for  $F$ .

**Proof:** Equation (4.4) can be written as  $g = h + g * F$ . Because of the definition  $R = \sum_{n=1}^{\infty} F^{*n}$  we obtain

$$F * (R * h + h) = F * h + \sum_{n=1}^{\infty} F^{*n+1} * h = \sum_{n=1}^{\infty} F^{*n} * h = R * h$$

which shows that  $g = R * h + h$  is indeed a solution of (4.4).

Let  $g'$  denote another solution and define the function

$$\delta := g' - R * h - h$$

Then (4.4) implies  $\delta = F * \delta$  and thus  $\delta = F^{*n} * \delta$  for all  $n \in \mathbb{N}$ . Since  $R(t) < \infty$  for any fixed  $t \geq 0$ , we infer that  $F^{*n} \rightarrow 0$  as  $n \rightarrow \infty$ . Hence  $\delta(t) = 0$  for all  $t \geq 0$ , which completes the proof.

□

### 4.3 Renewal Theorems

In order to present the most powerful results of renewal theory, it will be useful to introduce stopping times and Wald's lemma. Recall from (2.3) that a random variable  $S$  with values in  $\mathbb{N}_0 \cup \{\infty\}$  is called a stopping time for the sequence  $\mathcal{X} = (X_n : n \in \mathbb{N}_0)$  if

$$\mathbb{P}(S \leq n | \mathcal{X}) = \mathbb{P}(S \leq n | X_0, \dots, X_n) \quad (4.5)$$

holds for all  $n \in \mathbb{N}_0$ .

**Lemma 4.10** *For a renewal process  $\mathcal{N}$  with delay  $X_0$  and renewal intervals  $(X_n : n \in \mathbb{N})$ , the random variable  $N_t$  is a stopping time for the sequence  $(X_n : n \in \mathbb{N}_0)$ .*

**Proof:** This follows from the observation that  $N_t = k$  is equivalent to

$$\sum_{n=0}^{k-1} X_n \leq t < \sum_{n=0}^k X_n$$

which implies that the event  $N_t \leq k$  depends only on  $X_0, \dots, X_k$ .

□

**Lemma 4.11 Wald's Lemma**

Be  $\mathcal{X} = (X_n : n \in \mathbb{N}_0)$  a sequence of stochastically independent positive random variables with the same expectation  $\mathbb{E}(X_n) = m$  for all  $n \in \mathbb{N}$ . The expectations  $\mathbb{E}(X_0)$  and  $\mathbb{E}(X_1)$  shall be finite. Further be  $S$  a stopping time of the sequence  $\mathcal{X}$  with  $\mathbb{E}(S) < \infty$ . Then

$$\mathbb{E}\left(\sum_{n=0}^S X_n\right) = \mathbb{E}(X_0) + \mathbb{E}(S) \cdot m$$

**Proof:** For all  $n \in \mathbb{N}_0$  define the random variables  $I_n := 1$  on the set  $\{S \geq n\}$  and  $I_n := 0$  else. Then  $\sum_{n=0}^S X_n = \sum_{n=0}^{\infty} I_n X_n$  and hence

$$\mathbb{E}\left(\sum_{n=0}^S X_n\right) = \mathbb{E}\left(\sum_{n=0}^{\infty} I_n X_n\right) = \sum_{n=0}^{\infty} \mathbb{E}(I_n X_n)$$

by monotone convergence, as  $I_n$  and  $X_n$  are non-negative.  $S$  being a stopping time for  $\mathcal{X}$ , we obtain by definition  $\mathbb{P}(S \geq 0) = 1$ , and further

$$\mathbb{P}(S \geq n | \mathcal{X}) = 1 - \mathbb{P}(S \leq n-1 | \mathcal{X}) = 1 - \mathbb{P}(S \leq n-1 | X_0, \dots, X_{n-1})$$

for all  $n \in \mathbb{N}$ . Since the  $X_n$  are independent,  $I_n$  and  $X_n$  are independent, too, which implies  $\mathbb{E}(I_0 X_0) = \mathbb{E}(X_0)$  and

$$\mathbb{E}(I_n X_n) = \mathbb{E}(I_n) \cdot \mathbb{E}(X_n) = \mathbb{P}(S \geq n) \cdot m$$

for all  $n \in \mathbb{N}$ . Now the relation  $\sum_{n=1}^{\infty} \mathbb{P}(S \geq n) = \mathbb{E}(S)$  yields

$$\begin{aligned} \mathbb{E}\left(\sum_{n=0}^S X_n\right) &= \sum_{n=0}^{\infty} \mathbb{E}(I_n X_n) = \mathbb{E}(X_0) + \sum_{n=1}^{\infty} \mathbb{P}(S \geq n) \cdot m \\ &= \mathbb{E}(X_0) + \mathbb{E}(S) \cdot m \end{aligned}$$

□

**Theorem 4.12 Elementary Renewal Theorem**

Be  $\mathcal{N}$  a renewal process with renewal intervals  $(X_n : n \in \mathbb{N})$  and mean renewal time  $\mathbb{E}(X_1) = m > 0$ . Assume further that the mean delay is finite, i.e.  $\mathbb{E}(X_0) < \infty$ . Then for the counting function  $N_t$  the limit

$$\lim_{t \rightarrow \infty} \frac{\mathbb{E}(N_t)}{t} = \frac{1}{m}$$

holds, with the convention  $1/\infty := 0$ .

**Proof:** For every  $t \geq 0$ , the bound  $t < \sum_{n=0}^{N_t} X_n$  holds almost surely. By Wald's lemma, this implies

$$t < \mathbb{E} \left( \sum_{n=0}^{N_t} X_n \right) = \mathbb{E}(X_0) + \mathbb{E}(N_t) \cdot m$$

and thence for  $m < \infty$

$$\frac{1}{m} - \frac{\mathbb{E}(X_0)}{m \cdot t} < \frac{\mathbb{E}(N_t)}{t}$$

for all  $t \geq 0$ . For  $\mathbb{E}(X_0) < \infty$  and  $t \rightarrow \infty$ , this yields the bound

$$\liminf_{t \rightarrow \infty} \frac{\mathbb{E}(N_t)}{t} \geq \frac{1}{m}$$

which trivially holds for the case  $m = \infty$ .

Now it remains to show that  $\limsup_{t \rightarrow \infty} \mathbb{E}(N_t)/t \leq 1/m$ . To this aim we consider the truncated renewal process, denoted by  $\tilde{\mathcal{N}}$ , with the same delay  $\tilde{X}_0 = X_0$  but renewal intervals  $\tilde{X}_n = \min(X_n, M)$  for all  $n \in \mathbb{N}$ , with  $M$  being a fixed constant. Denote further  $\tilde{m} = \mathbb{E}(\tilde{X}_1)$ .

Because of  $\tilde{X}_n \leq M$  the bound  $\sum_{n=0}^{\tilde{N}_t} \tilde{X}_n \leq t + M$  holds almost certainly for all  $t \geq 0$ . Taking expectations and applying Wald's lemma, we obtain

$$\mathbb{E}(X_0) + \mathbb{E}(\tilde{N}_t) \cdot \tilde{m} = \mathbb{E} \left( \sum_{n=0}^{\tilde{N}_t} \tilde{X}_n \right) \leq t + M$$

For  $\mathbb{E}(X_0) < \infty$  and  $t \rightarrow \infty$ , this yields

$$\limsup_{t \rightarrow \infty} \frac{\mathbb{E}(\tilde{N}_t)}{t} \leq \frac{1}{\tilde{m}}$$

Since  $\tilde{X}_n \leq X_n$  for all  $n \in \mathbb{N}$ , we know that  $\tilde{N}_t \geq N_t$  for all  $t \geq 0$ . Thus we obtain further

$$\limsup_{t \rightarrow \infty} \frac{\mathbb{E}(N_t)}{t} \leq \frac{1}{\tilde{m}}$$

for any constant  $M$ . Now the result follows for  $M \rightarrow \infty$ .

□

**Remark 4.13** In view of theorem 4.4 one might be tempted to think that this trivially implied the statement of the above theorem 4.12. However, the following example shows that a limit with probability one in general does not imply a limit in expectation.

Let  $U$  denote a random variable which is uniformly distributed on the interval  $]0, 1[$ . Further define the random variables  $(V_n : n \in \mathbb{N})$  by

$$V_n := \begin{cases} 0, & U > 1/n \\ n, & U \leq 1/n \end{cases}$$

Since  $U > 0$  with probability one, we obtain the limit

$$V_n \rightarrow 0, \quad n \rightarrow \infty$$

with probability one. On the other hand, the expectation for  $V_n$  is given by

$$\mathbb{E}(V_n) = n \cdot \mathbb{P}(U \leq 1/n) = n \cdot \frac{1}{n} = 1$$

for all  $n \in \mathbb{N}$  and thus  $\mathbb{E}(V_n) \rightarrow 1$  as  $n \rightarrow \infty$ .

## 4.4 Residual Life Times and Stationary Renewal Processes

Choose any time  $t \geq 0$ . Denote the duration from  $t$  until the next arrival by  $B_t := S_{N_t+1} - t$  and call it the **residual life time** (or the **excess life**) at  $t$ . Further we define  $A_t := t - S_{N_t}$  and call  $A_t$  the **age** at  $t$ . These definitions imply the equality  $\{A_t > x\} = \{B_{t-x} > x\}$  for all  $x < t$ .

For a **stationary renewal process** we would postulate that the distribution of the counts in an interval  $[s, s+t]$  be independent of  $s$  and thus equal the distribution of  $N_t$ . If this holds for a process  $\mathcal{N}$ , then we also say that  $\mathcal{N}$  has **stationary**

**increments.** This implies in particular that the distribution of the residual life time must be independent of  $t$ , i.e. it coincides with the distribution of  $B_0$  and hence of  $X_0$ . We first guess that it satisfies

$$\mathbb{P}(X_0 \leq x) = \frac{1}{m} \int_0^x (1 - F(y)) dy \quad (4.6)$$

for all  $x \geq 0$ , where  $F$  denotes the distribution function of  $X_1$  and further  $m = \mathbb{E}(X_1) < \infty$ . Indeed we can show

**Theorem 4.14** *For a renewal process  $\mathcal{N}$  defined by (4.6) the following properties hold:*

- (1)  $\mathbb{E}(N_t) = t/m$  for all  $t \geq 0$
- (2)  $\mathbb{P}(B_t \leq x) = m^{-1} \int_0^x (1 - F(y)) dy$  for all  $t \geq 0$
- (3)  $\mathcal{N}$  has stationary increments.

**Proof:** (1) The distribution  $G$  of  $X_0$  has a density  $g(t) = \frac{1}{m}(1 - F(t))$ . Hence the Laplace–Stieltjes transform (LST) of  $G$  is

$$\begin{aligned} \tilde{G}(s) &= \int_0^\infty e^{-st} \frac{1}{m} (1 - F(t)) dt = \frac{1}{m} \left( \int_0^\infty e^{-st} dt - \int_0^\infty e^{-st} F(t) dt \right) \\ &= \frac{1}{m} \left( \frac{1}{s} - \frac{1}{s} \int_0^\infty e^{-st} dF(t) \right) = \frac{1 - \tilde{F}(s)}{sm} \end{aligned}$$

with  $\tilde{F}(s)$  denoting the LST of  $F$ . According to (4.1) we have the representation  $\mathbb{E}(N_t) = G * \sum_{n=0}^\infty F^{*n}(t)$  for all  $t \geq 0$ . Hence the LST of  $M(t) := \mathbb{E}(N_t)$  is given by

$$\tilde{M}(s) = \frac{\tilde{G}(s)}{1 - \tilde{F}(s)} = \frac{1}{sm}$$

for all  $s > 0$ , and thus coincides with the LST of the measure  $dx/m$ . Since the LST uniquely determines a function on  $[0, \infty[$ , this proves the first statement.

(2) The joint distributions

$$\begin{aligned} \mathbb{P}(B_t > x, N_t = 0) &= 1 - G(t + x) \\ \mathbb{P}(B_t > x, N_t = n) &= \int_0^\infty \mathbb{P}(B_t > x, N_t = n | S_n = y) dG * F^{*n-1}(y) \\ &= \int_0^t (1 - F(t + x - y)) dG * F^{*n-1}(y) \end{aligned}$$

for  $n \geq 1$  are immediate from the definition. Abbreviating  $F^c(x) := 1 - F(x)$ ,  $G^c(x) := 1 - G(x)$ , and denoting  $M(t) := \mathbb{E}(N_t)$ , we can write

$$\begin{aligned}
 \mathbb{P}(B_t > x) &= \sum_{n=0}^{\infty} \mathbb{P}(B_t > x, N_t = n) \\
 &= G^c(t+x) + \sum_{n=1}^{\infty} \int_0^t F^c(t+x-y) dG * F^{*n-1}(y) \\
 &= G^c(t+x) + \int_0^t F^c(t+x-y) d\left(\sum_{n=1}^{\infty} G * F^{*n-1}\right)(y) \\
 &= G^c(t+x) + \int_0^t F^c(t+x-y) dM(y)
 \end{aligned}$$

Using statement (1) and the definition of  $G$ , we obtain

$$\begin{aligned}
 \mathbb{P}(B_t > x) &= 1 - \frac{1}{m} \int_0^{t+x} (1 - F(y)) dy + \frac{1}{m} \int_0^t (1 - F(t+x-y)) dy \\
 &= 1 - \frac{1}{m} \int_0^x (1 - F(y)) dy
 \end{aligned}$$

which proves the second statement.

(3) The difference  $N_{t+s} - N_s$  simply counts the number  $N'_t$  of events in time  $t$  of the renewal process  $\mathcal{N}'$  with the same distribution  $F$  of  $X_1$  but a delay  $X'_0 \sim B_s$ . Now statement (2) shows that  $X_0 \sim B_s = B_0$ . Hence we obtain  $N'_t = N_t = N_{t+s} - N_s$  in distribution, which was to be proven.

□

Because of the results above a renewal process which satisfies condition (4.6) is called **stationary renewal process**. As one would expect, also the mean residual life time  $\mathbb{E}(B_t)$  of a stationary renewal process coincides with the limit of the mean residual life time of an ordinary renewal process:

**Lemma 4.15** *For a non-negative random variable  $X$  the  $n$ th moment can be expressed by*

$$\mathbb{E}(X^n) = \int_0^{\infty} \mathbb{P}(X > x) \cdot nx^{n-1} dx$$

**Proof:** This follows simply by writing

$$\mathbb{E}(X^n) = \int_0^{\infty} \mathbb{P}(X^n > z) dz = \int_0^{\infty} \mathbb{P}(X > \sqrt[n]{z}) dz$$

and substituting  $x = \sqrt[n]{z}$  with  $nx^{n-1} dx = dz$ .

□

**Theorem 4.16** *For a stationary renewal process with  $\mathbb{E}(X_1^2) < \infty$  the mean residual life time is given by*

$$\mathbb{E}(B_t) = \frac{\mathbb{E}(X_1^2)}{2m}$$

*independently of  $t \geq 0$ .*

**Proof:** Using part (2) of theorem 4.14, we obtain

$$\begin{aligned} \mathbb{E}(B_t) &= \int_0^\infty \mathbb{P}(B_t > y) dy = \frac{1}{m} \int_{y=0}^\infty \int_{x=y}^\infty (1 - F(x)) dx dy \\ &= \frac{1}{m} \int_{x=0}^\infty \int_{y=0}^x (1 - F(x)) dy dx = \frac{1}{m} \int_{x=0}^\infty \mathbb{P}(X_1 > x) \cdot x dx \end{aligned}$$

and the statement follows from lemma 4.15.

□

**Example 4.17** Waiting time at a bus stop

Consider a bus stop where buses are scheduled to arrive in intervals of length  $T$ . However, due to traffic variations the real inter-arrival times are uniformly distributed within intervals  $[T - a, T + a]$  with some  $a > 0$ . Now suppose that somebody arrives at the bus stop "at random", i.e. without knowing the bus schedule. Then we can model the mean waiting time for the next bus by the mean residual life time  $\mathbb{E}(B_t)$  in a stationary renewal process with distribution  $X_1 \sim U(T - a, T + a)$ . We obtain

$$\mathbb{E}(X_1^2) = \frac{1}{2a} \int_{T-a}^{T+a} x^2 dx = \frac{1}{2a} \cdot \frac{1}{3} (6T^2a + 2a^3) = T^2 + \frac{a^2}{3}$$

and by theorem 4.16

$$\mathbb{E}(B_t) = \frac{T^2 + \frac{a^2}{3}}{2 \cdot T} = \frac{T}{2} + \frac{a^2}{6 \cdot T}$$

Thus the mean waiting time for random inter-arrival times (meaning  $a > 0$ ) is longer than it would be for deterministic ones (namely  $T/2$ ). This phenomenon is called the **waiting time paradox**.



## Notes

A classical presentation of renewal theory is chapter 11 in Feller [9]. The presentation in this chapter is largely adapted to Ross [17, 18] as well as Karlin and Taylor [14].

**Exercise 4.18** Prove theorem 4.3.

**Exercise 4.19** In the proof for Wald's lemma 4.11 we have used the relation  $\mathbb{E}(S) = \sum_{n=0}^{\infty} \mathbb{P}(S > n)$ . For a positive continuous distribution function  $F$ , the equivalent is  $\mathbb{E}(F) = \int_0^{\infty} (1 - F(y)) dy$ . Give a proof for these equations.

**Exercise 4.20** Show for a Poisson process  $\mathcal{N}$  with intensity  $\lambda > 0$  that

$$\mathbb{P}(N_t = k) = \frac{(\lambda t)^k}{k!} e^{-\lambda t}$$

for all  $t \geq 0$  and  $k \in \mathbb{N}_0$ , and  $\mathbb{E}(N_t) = \lambda \cdot t$ .

**Exercise 4.21** A plumber receives orders at time intervals which are distributed exponentially with parameter  $\lambda$ . As soon as he has received an order he goes to work, which takes an exponentially distributed time with parameter  $\mu$ . During work he cannot receive any orders. Assume that at time zero the plumber is working. Give a model of the plumber receiving orders in terms of a renewal process and determine the density of the renewal intervals' distribution.

**Exercise 4.22** An intelligence agency eavesdrops on telephone calls automatically. If there occurs a suspicious sequence of words, a closer investigation is initiated. The probability for such a sequence is one in a thousand for every call. The length of a call is distributed exponentially with a mean of 20 seconds. How long is the expected amount of time before a closer investigation begins? Use Wald's lemma.

**Exercise 4.23** Let  $\mathcal{N} = (N_t : t \geq 0)$  denote an ordinary renewal process with  $X_1 \sim F$ . Show that the current life time  $X_{N_t}$  satisfies

$$\mathbb{P}(X_{N_t} > x) \geq 1 - F(x)$$

for all  $x \geq 0$ .

**Exercise 4.24** Show that the age  $A_t$  of a stationary renewal process is distributed as

$$\mathbb{P}(A_t \leq x) = \frac{1}{m} \int_0^x (1 - F(y)) dy$$

independently of  $t \geq 0$ .

# Chapter 5

## Appendix

### 5.1 Conditional Expectations and Probabilities

Let  $(\Omega, \mathcal{A}, P)$  denote a probability space and  $(S, \mathcal{B})$  a measurable space. A **random variable** is a measurable mapping  $X : \Omega \rightarrow S$ , which means that  $X^{-1}(B) \in \mathcal{A}$  for all  $B \in \mathcal{B}$ . In other words,  $X$  is a random variable if and only if  $X^{-1}(\mathcal{B}) \subset \mathcal{A}$ . In stochastic models, a random variable usually gives information on a certain phenomenon, e.g. the number of users in a queue at some specific time.

Consider any real-valued random variable  $X : (\Omega, \mathcal{A}) \rightarrow (\mathbb{R}, \mathcal{B})$ ,  $\mathcal{B}$  denoting the Borel  $\sigma$ -algebra on  $\mathbb{R}$ , which is integrable or non-negative. While the random variable  $X$  itself yields the full information, a rather small piece of information on  $X$  is given by its **expectation**

$$\mathbb{E}(X) := \int_{\Omega} X \, dP$$

The conditional expectation is a concept that yields a degree of information which lies between the full information  $X$  and its expectation  $\mathbb{E}(X)$ .

To motivate the definition, we first observe that the distribution  $P^X = P \circ X^{-1}$  of  $X$  is a measure on the sub- $\sigma$ -algebra  $X^{-1}(\mathcal{B})$  of  $\mathcal{A}$ , i.e. in order to compute

$$P(X \in B) = P^X(B) = \int_{X^{-1}(B)} dP$$

we need to evaluate the measure  $P$  on sets

$$A := X^{-1}(B) \in X^{-1}(\mathcal{B}) \subset \mathcal{A}$$

On the other hand, the expectation  $\mathbb{E}(X)$  is an evaluation of  $P$  on the set  $\Omega = X^{-1}(S)$  only. Thus we can say that the expectation employs  $P$  only on the trivial  $\sigma$ -algebra  $\{\emptyset, \Omega\}$ , while  $X$  itself employs  $P$  on the  $\sigma$ -algebra  $X^{-1}(\mathcal{B})$  generated by  $X$ .

Now we take any sub- $\sigma$ -algebra  $\mathcal{C} \subset \mathcal{A}$ . According to the Radon–Nikodym theorem there is a random variable  $X_0 : \Omega \rightarrow S$  with  $X^{-1}(\mathcal{B}) = \mathcal{C}$  and

$$\int_{\mathcal{C}} X_0 dP = \int_{\mathcal{C}} X dP \quad (5.1)$$

for all  $C \in \mathcal{C}$ . This we call the **conditional expectation** of  $X$  under  $\mathcal{C}$  and write

$$\mathbb{E}(X|\mathcal{C}) := X_0$$

A conditional expectation is  $P$ -almost certainly uniquely determined by (5.1). Typical special cases and examples are

**Example 5.1** For  $\mathcal{C} = \{\emptyset, \Omega\}$ , the conditional expectation equals the expectation, i.e.  $\mathbb{E}(X|\mathcal{C}) = \mathbb{E}(X)$ . For any  $\sigma$ -algebra  $\mathcal{C}$  with  $X^{-1}(\mathcal{B}) \subset \mathcal{C}$  we obtain  $\mathbb{E}(X|\mathcal{C}) = X$ .

**Example 5.2** Let  $I$  denote any index set and  $(Y_i : i \in I)$  a family of random variables. For the  $\sigma$ -algebra  $\mathcal{C} = \sigma(\bigcup_{i \in I} Y_i^{-1}(\mathcal{B}))$  generated by  $(Y_i : i \in I)$ , we write

$$\mathbb{E}(X|Y_i : i \in I) := \mathbb{E}(X|\mathcal{C})$$

By definition we obtain for a  $\sigma$ -algebra  $\mathcal{C} \subset \mathcal{A}$ , random variables  $X$  and  $Y$ , and real numbers  $\alpha$  and  $\beta$

$$\mathbb{E}(\alpha X + \beta Y|\mathcal{C}) = \alpha \mathbb{E}(X|\mathcal{C}) + \beta \mathbb{E}(Y|\mathcal{C})$$

For  $\sigma$ -algebras  $\mathcal{C}_1 \subset \mathcal{C}_2 \subset \mathcal{A}$  we obtain

$$\mathbb{E}(\mathbb{E}(X|\mathcal{C}_2)|\mathcal{C}_1) = \mathbb{E}(\mathbb{E}(X|\mathcal{C}_1)|\mathcal{C}_2) = \mathbb{E}(X|\mathcal{C}_1) \quad (5.2)$$

Let  $\mathcal{C}_1$  and  $\mathcal{C}_2$  denote sub- $\sigma$ -algebras of  $\mathcal{A}$ ,  $\mathcal{C} := \sigma(\mathcal{C}_1 \cup \mathcal{C}_2)$ , and  $X$  an integrable random variable. If  $\sigma(X^{-1}(\mathcal{B}) \cup \mathcal{C}_1)$  and  $\mathcal{C}_2$  are independent, then

$$\mathbb{E}(X|\mathcal{C}) = \mathbb{E}(X|\mathcal{C}_1)$$

If  $X$  and  $Y$  are integrable random variables and  $X^{-1}(\mathcal{B}) \subset \mathcal{C}$ , then

$$\mathbb{E}(XY|\mathcal{C}) = X \cdot \mathbb{E}(Y|\mathcal{C}) \quad (5.3)$$

Conditional probabilities are special cases of conditional expectations. Define the **indicator function** of a measurable set  $A \in \mathcal{A}$  by

$$1_A(x) := \begin{cases} 1, & x \in A \\ 0, & x \notin A \end{cases}$$

Such a function is a random variable, since

$$1_A^{-1}(\mathcal{B}) = \{\emptyset, A, A^c, \Omega\} \subset \mathcal{A}$$

with  $A^c := \Omega \setminus A$  denoting the complement of the set  $A$ . Let  $\mathcal{C}$  denote a sub- $\sigma$ -algebra of  $\mathcal{A}$ . The conditional expectation of  $1_A$  is called **conditional probability** of  $A$ . We write

$$P(A|\mathcal{C}) := \mathbb{E}(1_A|\mathcal{C})$$

Immediate properties of conditional probabilities are

$$0 \leq P(A|\mathcal{C}) \leq 1, \quad P(\emptyset|\mathcal{C}) = 0, \quad P(\Omega|\mathcal{C}) = 1$$

$$A_1 \subset A_2 \implies P(A_1|\mathcal{C}) \leq P(A_2|\mathcal{C})$$

all of which hold  $P$ -almost certainly. For a sequence  $(A_n : n \in \mathbb{N})$  of disjoint measurable sets, i.e.  $A_n \in \mathcal{A}$  for all  $n \in \mathbb{N}$  and  $A_i \cap A_j = \emptyset$  for  $i \neq j$ , we obtain

$$P\left(\bigcup_{n=1}^{\infty} A_n \middle| \mathcal{C}\right) = \sum_{n=1}^{\infty} P(A_n|\mathcal{C})$$

$P$ -almost certainly. Let  $X : (\Omega, \mathcal{A}) \rightarrow (\mathbb{R}, \mathcal{B})$  denote a non-negative or integrable random variable and  $Y : (\Omega, \mathcal{A}) \rightarrow (\Omega', \mathcal{A}')$  a random variable. Then there is a measurable function  $g : (\Omega', \mathcal{A}') \rightarrow (\mathbb{R}, \mathcal{B})$  with

$$\mathbb{E}(X|Y) = g \circ Y$$

This is  $P^Y$ -almost certainly determined by

$$\int_{A'} g \, dP^Y = \int_{Y^{-1}(A')} X \, dP$$

for all  $A' \in \mathcal{A}'$ . Then we can define the conditional probability of  $X$  given  $Y = y$  as  $g(y)$ . We write

$$\mathbb{E}(X|Y = y) := g(y)$$

for all  $y \in \Omega'$ .

## 5.2 Extension Theorems

Throughout this book, our basic stochastic tools are either sequences of random variables (such as Markov chains or Markov renewal chains) or even uncountable families of random variables (such as Markov processes, renewal processes, or semi-regenerative processes). It is essential for our models that these random variables are dependent, and in fact we define them in terms of conditional probabilities, i.e. via their dependence structure.

It is then an immediate question whether a probability measure  $\mathbb{P}$  exists that satisfies all the postulates in the definition of a stochastic sequence or process. This question is vital as it concerns the very existence of the tools we are using.

### 5.2.1 Stochastic chains

Let  $(S, \mathcal{B})$  denote a measurable space,  $\mu$  a probability measure on  $(S, \mathcal{B})$ , and  $P_n$ ,  $n \in \mathbb{N}$ , stochastic **kernels** on  $(S, \mathcal{B})$ . The latter means that for every  $n \in \mathbb{N}$ ,  $P_n : S \times \mathcal{B} \rightarrow [0, 1]$  is a function that satisfies

(K1) For every  $x \in S$ ,  $P_n(x, \cdot)$  is a probability measure on  $(S, \mathcal{B})$ .

(K2) For every  $A \in \mathcal{B}$ , the function  $P_n(\cdot, A)$  is  $\mathcal{B}$ -measurable.

Define  $S^\infty$  as the set of all sequences  $x = (x_n : n \in \mathbb{N}_0)$  with  $x_n \in S$  for all  $n \in \mathbb{N}_0$ . A subset of  $S^\infty$  having the form

$$C_{n_1, \dots, n_k}(A) = \{x \in S^\infty : (x_{n_1}, \dots, x_{n_k}) \in A\}$$

with  $k \in \mathbb{N}$ ,  $n_1 < \dots < n_k \in \mathbb{N}_0$ , and  $A \in \mathcal{B}^k$ , is called **cylinder** (with coordinates  $n_1, \dots, n_k$  and base  $A$ ). The set  $\mathcal{C}$  of all cylinders in  $S^\infty$  forms an algebra of sets. Define  $\mathcal{B}^\infty := \sigma(\mathcal{C})$  as the minimal  $\sigma$ -algebra containing  $\mathcal{C}$ .

Now we can state the extension theorem for sequences of random variables, which is proven in Gikhman and Skorokhod [10], section II.4.

**Theorem 5.3** *There is a probability measure  $\mathbb{P}$  on  $(S^\infty, \mathcal{B}^\infty)$  satisfying*

$$\begin{aligned} \mathbb{P}(C_{0, \dots, k}(A_0 \times \dots \times A_k)) &= \int_{A_0} d\mu(x_0) \int_{A_1} P_1(x_0, dx_1) \dots \\ &\dots \int_{A_{k-1}} P_{k-1}(x_{k-2}, dx_{k-1}) P_k(x_{k-1}, A_k) \end{aligned} \quad (5.4)$$

for all  $k \in \mathbb{N}_0$ ,  $A_0, \dots, A_k \in \mathcal{B}$ . The measure  $\mathbb{P}$  is uniquely determined by the system (5.4) of equations.

The first part of the theorem above justifies our definitions of Markov chains and Markov renewal chains. The second part states in particular that a Markov chain is uniquely determined by its initial distribution and its transition matrix.

Based on this result, we may define a **stochastic chain** with state space  $S$  as a sequence  $(X_n : n \in \mathbb{N}_0)$  of  $S$ -valued random variables which are distributed according to a probability measure  $\mathbb{P}$  on  $(S^\infty, \mathcal{B}^\infty)$ .

### 5.2.2 Stochastic processes

Let  $S$  denote a Polish (i.e. a complete separable metric) space, and  $\mathcal{B}$  the Borel  $\sigma$ -algebra on  $S$ . Define  $\Omega$  as the set of all functions  $f : \mathbb{R}_0^+ \rightarrow S$ . In order to construct an appropriate  $\sigma$ -algebra on  $\Omega$ , we again start from the cylinder sets

$$C_{t_1, \dots, t_k}(A) = \{f \in \Omega : (f(t_1), \dots, f(t_k)) \in A\}$$

for  $k \in \mathbb{N}$ ,  $t_1 < \dots < t_k \in \mathbb{R}_0^+$ , and  $A \in \mathcal{B}^k$ . Denote the set of all cylinders in  $\Omega$  by  $\mathcal{C}$ . Again,  $\mathcal{C}$  forms an algebra of sets and we can define  $\mathcal{A} := \sigma(\mathcal{C})$  as the minimal  $\sigma$ -algebra containing  $\mathcal{C}$ .

Let  $\mathcal{M} = \{\mu_{t_1, \dots, t_k} : k \in \mathbb{N}, t_1, \dots, t_k \in \mathbb{R}_0^+\}$  denote a family of probability distributions with

(C1) For all  $k \in \mathbb{N}$ ,  $t_1, \dots, t_k \in \mathbb{R}_0^+$ , and  $A \in \mathcal{B}^k$

$$\mu_{t_1, \dots, t_k, t_{k+1}}(A \times S) = \mu_{t_1, \dots, t_k}(A)$$

(C2) For all  $k \in \mathbb{N}$  and permutations  $\pi : \{1, \dots, k\} \rightarrow \{1, \dots, k\}$

$$\mu_{\pi(t_1), \dots, \pi(t_k)}(\pi(A)) = \mu_{t_1, \dots, t_k}(A)$$

Then the family  $\mathcal{M}$  is called **compatible**.

**Remark 5.4** Condition (C1) ensures that the distributions are consistent with each other, while condition (C2) is merely notational.

The following extension theorem by Kolmogorov is proven in Gikhman and Skorokhod [12], section 3.2.

**Theorem 5.5** *Let  $\{\mu_{t_1, \dots, t_k} : k \in \mathbb{N}, t_1, \dots, t_k \in \mathbb{R}_0^+\}$  denote a compatible family of probability measures. Then there is a probability measure  $\mathbb{P}$  on  $(\Omega, \mathcal{A})$  with*

$$\mathbb{P}(\{f \in \Omega : (f(t_1), \dots, f(t_k)) \in A\}) = \mu_{t_1, \dots, t_k}(A) \quad (5.5)$$

*for all  $k \in \mathbb{N}$ ,  $t_1, \dots, t_k \in \mathbb{R}_0^+$ , and  $A \in \mathcal{B}^k$ . The measure  $\mathbb{P}$  is uniquely determined by the system (5.5) of equations.*

Based on this, we define a **stochastic process** with Polish state space  $S$  as a family  $X = (X_t : t \in \mathbb{R}_0^+)$  of  $S$ -valued random variables which are distributed according to a probability measure  $\mathbb{P}$  on  $(\Omega, \mathcal{A})$ . An element  $\omega \in \Omega$  is an arbitrary function  $\omega : \mathbb{R}_0^+ \rightarrow S$ . It is also called a **path** of  $X$ . If we want to state that the support of  $\mathbb{P}$  consists of a special class of functions (say right-continuous ones), then we say that  $X$  is a stochastic process with right-continuous paths. The above family  $\mathcal{M}$  of probability measures is called the set of **finite-dimensional marginal distributions** for  $X$ .

Due to theorem 5.5 a Markov process is uniquely defined by its initial distribution and the family of transition probabilities, since they determine all finite-dimensional marginal distributions. Further our constructions of Markov processes, renewal processes, and semi-Markov processes yield compatible sets of finite-dimensional marginal distributions, hence by theorem 5.5 a probability measure  $\mathbb{P}$  for the respective process.

## 5.3 Transforms

In several parts of the present book, it is essential to argue via transforms of distributions. The necessary background for these shall be presented shortly in this section. For discrete distributions on  $\mathbb{N}_0$  we will introduce  $z$ -transforms, while for distributions on  $\mathbb{R}_0^+$  the Laplace-Stieltjes transform will be useful.

### 5.3.1 $z$ -transforms

Let  $X$  denote a  $\mathbb{N}_0$ -valued random variable with distribution  $A = (a_n : n \in \mathbb{N}_0)$ , i.e.  $\mathbb{P}(X = n) = a_n$  for all  $n \in \mathbb{N}_0$ . Then the power series

$$A^*(z) := \sum_{n=0}^{\infty} a_n z^n \quad (5.6)$$

converges absolutely for  $z \in \mathbb{C}$  with  $|z| \leq 1$  and is analytic in this region. We note that  $A^*(z) = \mathbb{E}(z^X)$ . If  $A(z)$  is a given power series for a distribution  $(a_n : n \in \mathbb{N}_0)$ , then the probabilities  $a_n$  can be derived as

$$a_n = \frac{1}{n!} \left. \frac{d^n}{dz^n} A(z) \right|_{z=0}$$



for all  $n \in \mathbb{N}_0$ . Thus the mapping between discrete distributions on  $\mathbb{N}_0$  and the power series in (5.6) is bijective, and we may call  $A^*(z)$  the (uniquely determined)  **$z$ -transform** of  $X$  (also: of the distribution  $A$ ).

**Example 5.6** For a Dirac distribution on  $k \in \mathbb{N}_0$  with

$$a_n = \begin{cases} 1, & n = k \\ 0, & n \neq k \end{cases}$$

we obtain  $A^*(z) = z^k$ .

**Example 5.7** Let  $A$  denote the geometric distribution with some parameter  $p \in ]0, 1[$ , i.e.

$$a_n = (1 - p)p^n$$

for all  $n \in \mathbb{N}_0$ . The  $z$ -transform of  $A$  is given by

$$A^*(z) = (1 - p) \sum_{n=0}^{\infty} p^n z^n = \frac{1 - p}{1 - pz}$$

for all  $|z| \leq 1$ .

A very useful feature is the behaviour of the  $z$ -transform with respect to the convolution of two distributions. Let  $A = (a_n : n \in \mathbb{N}_0)$  and  $B = (b_n : n \in \mathbb{N}_0)$  denote two distributions on  $\mathbb{N}_0$ . The convolution  $C = A * B$  of  $A$  and  $B$  is defined as the distribution  $C = (c_n : n \in \mathbb{N}_0)$  with

$$c_n = \sum_{k=0}^n a_k b_{n-k}$$

for all  $n \in \mathbb{N}_0$ . For the  $z$ -transform of  $C$  we obtain

$$\begin{aligned} C^*(z) &= \sum_{n=0}^{\infty} c_n z^n = \sum_{n=0}^{\infty} \sum_{k=0}^n a_k b_{n-k} z^n = \sum_{n=0}^{\infty} a_k z^k \sum_{n=k}^{\infty} b_{n-k} z^{n-k} \\ &= A^*(z) \cdot B^*(z) \end{aligned}$$

for all  $|z| \leq 1$ .

This means that the  $z$ -transform of a convolution  $A * B$  equals the product  $A^*(z) \cdot B^*(z)$  of the  $z$ -transform of  $A$  and  $B$ . In terms of random variables we have

the following representation: Let  $X$  and  $Y$  denote two independent  $\mathbb{N}_0$ -valued random variables. Then the  $z$ -transform of the sum  $X + Y$  equals the product of the  $z$ -transforms of  $X$  and  $Y$ , i.e.

$$\mathbb{E}(z^{X+Y}) = \mathbb{E}(z^X) \cdot \mathbb{E}(z^Y)$$

for all  $|z| \leq 1$ .

### 5.3.2 Laplace–Stieltjes transforms

Let  $X$  denote an  $\mathbb{R}_0^+$ -valued random variable with distribution function  $F$ . The **Laplace–Stieltjes transform (LST)** of  $X$  (or: of  $F$ ) is defined by

$$F^*(s) := \int_0^\infty e^{-st} dF(t) = \mathbb{E}(e^{-sX})$$

for all  $s \in \mathbb{C}$  with  $\operatorname{Re}(s) \geq 0$ . The LST uniquely determines its underlying distribution.

**Example 5.8** Let  $X$  be exponentially distributed with parameter  $\lambda$ , i.e.  $X$  has the distribution function  $F(t) = 1 - e^{-\lambda t}$  with Lebesgue density  $f(t) = \lambda e^{-\lambda t}$ . Then

$$F^*(s) = \int_0^\infty e^{-st} \lambda e^{-\lambda t} dt = \frac{\lambda}{s + \lambda}$$

for  $\operatorname{Re}(s) \geq 0$ .

**Example 5.9** For the Dirac distribution  $\delta_x$  on  $x \in \mathbb{R}_0^+$  we obtain

$$\delta_x^*(s) = \int_0^\infty e^{-st} dF(t) \quad \text{with} \quad F(t) = \begin{cases} 0, & t < x \\ 1, & t \geq x \end{cases}$$

and hence

$$\delta_x^*(s) = e^{-sx}$$

for  $\operatorname{Re}(s) \geq 0$ .

Like the  $z$ -transform, the LST is very useful for dealing with convolutions. Let  $X$  and  $Y$  denote two independent  $\mathbb{R}_0^+$ -valued random variables. Then the LST of the sum  $X + Y$  equals the product of the LSTs of  $X$  and  $Y$ , i.e.

$$\mathbb{E}(e^{-s(X+Y)}) = \mathbb{E}(e^{-sX}) \cdot \mathbb{E}(e^{-sY})$$

for all  $s \in \mathbb{C}$  with  $\operatorname{Re}(s) \geq 0$ .

## Notes

For more on  $z$ -transforms see e.g. Juri [13], or the collection of results in Kleinrock [15], appendix I. For Laplace–Stieltjes transforms see chapter XIII in Feller [9] or again Kleinrock [15], appendix I.

## 5.4 Gershgorin's circle theorem

An important theorem to find bounds for the eigenvalues of a matrix has been developed by Gershgorin in 1938. For ease of reference it shall be presented in this section. Let  $A = (a_{ij})_{i,j \leq m}$  denote a square matrix of dimension  $m$  with entries  $a_{ij} \in \mathbb{C}$ . The following theorem is called **Gershgorin's circle theorem**.

**Theorem 5.10** *All eigenvalues of the matrix  $A$  lie in the union  $C := \bigcup_{i=1}^m C_i$  of the circles*

$$C_i = \left\{ z \in \mathbb{C} : |z - a_{ii}| \leq \sum_{k \neq i} |a_{ik}| \right\}$$

**Proof:** Let  $x^{(\nu)}$  denote an eigenvector to the eigenvalue  $\lambda_\nu$  of  $A$ , i.e.  $Ax^{(\nu)} = \lambda_\nu x^{(\nu)}$ . This implies

$$\sum_{k=1}^m a_{ik} x_k^{(\nu)} = \lambda_\nu x_i^{(\nu)} \quad (5.7)$$

for all  $i \leq m$ . Since an eigenvector is determined only up to a scalar multiplicative, we can assume without loss of generality that there is a component

$$x_{i_0}^{(\nu)} = \max_{1 \leq j \leq m} |x_j^{(\nu)}| = 1$$

of the vector  $x^{(\nu)}$ . Now (5.7) yields for  $i = i_0$  the relation

$$\sum_{k \neq i_0} a_{i_0,k} x_k^{(\nu)} = (\lambda_\nu - a_{i_0,i_0}) x_{i_0}^{(\nu)} = \lambda_\nu - a_{i_0,i_0}$$

which implies by the triangle inequality

$$|\lambda_\nu - a_{i_0,i_0}| \leq \sum_{k \neq i_0} |a_{i_0,k}| \cdot |x_k^{(\nu)}| \leq \sum_{k \neq i_0} |a_{i_0,k}|$$

Since every eigenvalue satisfies at least one such inequality, the proof is complete.

□

**Corollary 5.11** *If  $A$  is diagonally dominated, i.e. if*

$$|a_{ii}| > \sum_{k \neq i} |a_{ik}|$$

*holds for all  $1 \leq i \leq m$ , then the matrix  $A$  is invertible.*

**Proof:** The strict inequality of the assumption implies that  $a_{ii} \neq 0$  for all  $i \leq m$ . Applying theorem 5.10 yields a restriction

$$|\lambda| \geq |a_{ii}| - |a_{ii} - \lambda| \geq |a_{ii}| - \sum_{k \neq i} |a_{ik}| > 0$$

for every eigenvalue  $\lambda$  of  $A$ . Therefore the matrix  $A$  has no eigenvalue zero and thus is invertible.

□

# Bibliography

- [1] L. Breiman. *Probability*. Philadelphia, PA: SIAM, 1968.
- [2] L. Breuer. The Periodic BMAP/PH/c Queue. *Queueing Systems*, 38(1):67–76, 2001.
- [3] L. Breuer. Two Examples for Computationally Tractable Periodic Queues. *International Journal of Simulation*, 3(3–4):15–24, 2003.
- [4] E. Çinlar. *Introduction to stochastic processes*. Englewood Cliffs, N. J.: Prentice-Hall, 1975.
- [5] K. Chung. *Markov chains with stationary transition probabilities*. Springer-Verlag, Berlin etc., 1960.
- [6] T. Engset. Emploi du calcul des probabilités pour la détermination du nombre de sélecteurs dans les Bureaux Téléphoniques Centraux. *Rev. gén. elect.*, 9:138–140, 1921.
- [7] A. Erlang. Solution of Some Probability Problems of Significance for Automatic Telephone Exchanges. *Electroteknikeren*, 13:5–13, 1917.
- [8] W. Feller. *An introduction to probability theory and its applications. Vol. I*. New York etc.: John Wiley and Sons, 1950.
- [9] W. Feller. *An introduction to probability theory and its applications. Vol. II. 2nd ed.* New York etc.: John Wiley and Sons, 1971.
- [10] I. Gihman and A. Skorohod. *The theory of stochastic processes I*. Berlin etc.: Springer, 1974.
- [11] I. Gihman and A. Skorohod. *The theory of stochastic processes II*. Berlin etc.: Springer-Verlag, 1975.

- [12] I. Gikhman and A. Skorokhod. *Introduction to the theory of random processes*. Saunders, 1969.
- [13] E. Juri. *Theory and Application of the  $z$ -Transform Method*. Wiley, New York, 1964.
- [14] S. Karlin and H. M. Taylor. *A first course in stochastic processes*. 2nd ed. New York etc.: Academic Press, 1975.
- [15] L. Kleinrock. *Queueing systems. Vol. I: Theory*. New York etc.: John Wiley & Sons, 1975.
- [16] S. Meyn and R. Tweedie. *Markov chains and stochastic stability*. Berlin: Springer-Verlag, 1993.
- [17] S. Ross. *Applied Probability Models with Optimization Applications*. Holden-Day, San Francisco, 1970.
- [18] S. Ross. *Stochastic Processes*. John Wiley & Sons, New York etc., 1983.