

Spinal injuries



Waleed M. Awwad, MD. FRCSC
Assistant professor and Consultant
Orthopedic Surgery department

Objectives



- ❧ The ability to demonstrate knowledge of the following:
 - ❧ Basic anatomy of the spine.
 - ❧ Initial assessment and treatment of spinal injuries at the field.
 - ❧ Principle of spinal stability.
 - ❧ Understanding of neurologic syndromes caused by spinal trauma.
 - ❧ Management of Cauda equina syndrome.

Spine Pathology Red Flag Conditions



☞ Beware of:

- 1) Cauda Equina/severe neurologic injury (perianal numbness, decreased rectal tone, loss of movement in the extremities).
- 2) Tumour weakening the vertebrae (causing cord compression or vertebral fracture).
- 3) Infection weakening bone (causing disc/vertebral destruction or cord compression).
- 4) **Traumatic Spine Fracture** (causing vertebral angulation, pain, or neuro compromise).



Remember that spine fracture can occur without trauma.




Anatomy





Anatomy




Anatomy




Anatomy



Anatomy



Anatomy




Anatomy




Thoracic Vertebrae

Axial (Overhead) View





Lateral (Side) View



Anatomy






Anatomy




Anatomy




Anatomy



Anatomy



Epidemiology

- ❧ 56000 cases per year. 
- ❧ 11000 new spinal cord injuries.
- ❧ 15-20% multiple non-contiguous levels.
- ❧ 10% involving the cervical spine.
- ❧ 90% involving thoracolumbar spine.
- ❧ 25% have neurologic deficit.
- ❧ Age: mostly between 15-24 years.
- ❧ Gender: mostly males (4:1).






Fig. 56-6 Three patterns of multiple-level injury described by Calenoff et al. (see text). (From Calenoff L, Chessare JW, Rogers LF, et al *Am J Roentgenol* 130:665, 1978.)



Mechanism of Injury



- ❧ High energy trauma such as an MVA or fall from a height or a horse.
 - ❧ MVA: 40-55%
 - ❧ Falls: 20-30%
 - ❧ Sports: 6-12%
 - ❧ Others: 12-21%

- ❧ Low energy trauma in a high risk patient (ie a patient with known spinal canal compromise such as ankylosing spondylitis, Osteoporosis or metastatic vertebral lesions)

- ❧ Penetrating trauma from gunshot or knives.




Spine stability




- ❧ Cervical spine instability:
 - ❧ Compression fracture with 25% loss of height.
 - ❧ Angular displacement > 11 degrees.
 - ❧ Translation $> 3.5\text{mm}$.
 - ❧ Disc space separation $> 1.7\text{mm}$.

- ❧ Thoracic and lumbar spine: Denis three column.

The Three columns



Instability exists with disruption of any two of three columns.



Assessment



- ❧ In cases of trauma, ABCDE' s must be assessed first and treated appropriately.
- ❧ Patients should be examined with spinal collar until spinal pathology is excluded.
- ❧ Careful log rolling keeping the head, neck and pelvis in line should be done to examine the spine properly.

Assessment



- ❧ Immobilization.
- ❧ History:
 - ❧ Mechanism of injury:
 - ❧ compression, flexion, extension, distraction
 - ❧ Other injuries.
 - ❧ Seat belt.
 - ❧ Other causalities.
- ❧ Physical examination:
 - ❧ Inspection, palpation.
 - ❧ Neurologic examination.

Immobilization



Immobilization



Immobilization



NEUROLOGIC



❧ **Muscle Test**

❧ **Sensory exam**

*light touch, Sharp dull discrimination, Vibration sense, Proprioception
and two-point discrimination*

❧ **Reflexes**

Signs of Spinal Trauma



- ❧ Apnea, lower cranial nerve injury VIII-XII (high C-spine).
- ❧ Deformity of the spine or neck.
- ❧ Tenderness on palpation along spinal processes.
- ❧ Paralysis or muscle weakness (which spinal level).
- ❧ Loss of sensation (which dermatomes).
- ❧ Loss of rectal tone.
- ❧ Positive Babinski sign.

Asia Score: Brief Trauma Neurologic Survey

Patient Name _____

Examiner Name _____ Date/Time of Exam _____



STANDARD NEUROLOGICAL CLASSIFICATION
OF SPINAL CORD INJURY



MOTOR

KEY MUSCLES
(scoring on reverse side)

	R	L	
C5	<input type="checkbox"/>	<input type="checkbox"/>	Elbow flexors
C6	<input type="checkbox"/>	<input type="checkbox"/>	Wrist extensors
C7	<input type="checkbox"/>	<input type="checkbox"/>	Elbow extensors
C8	<input type="checkbox"/>	<input type="checkbox"/>	Finger flexors (distal phalanx of middle finger)
T1	<input type="checkbox"/>	<input type="checkbox"/>	Finger abductors (little finger)
UPPER LIMB TOTAL	<input type="checkbox"/>	+ <input type="checkbox"/>	= <input type="checkbox"/>
(MAXIMUM)	(25)	(25)	(50)

Comments:

L2	<input type="checkbox"/>	<input type="checkbox"/>	Hip flexors
L3	<input type="checkbox"/>	<input type="checkbox"/>	Knee extensors
L4	<input type="checkbox"/>	<input type="checkbox"/>	Ankle dorsiflexors
L5	<input type="checkbox"/>	<input type="checkbox"/>	Long toe extensors
S1	<input type="checkbox"/>	<input type="checkbox"/>	Ankle plantar flexors

Voluntary anal contraction
(Yes/No)



LOWER LIMB TOTAL	<input type="checkbox"/>	+ <input type="checkbox"/>	= <input type="checkbox"/>
(MAXIMUM)	(25)	(25)	(50)

	LIGHT TOUCH		PIN PRICK	
	R	L	R	L
C2	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
C3	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
C4	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
C5	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
C6	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
C7	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
C8	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
T1	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
T2	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
T3	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
T4	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
T5	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
T6	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
T7	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
T8	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
T9	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
T10	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
T11	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
T12	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
L1	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
L2	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
L3	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
L4	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
L5	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
S1	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
S2	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
S3	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
S4-5	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
TOTALS	<input type="checkbox"/>	+ <input type="checkbox"/>	= <input type="checkbox"/>	<input type="checkbox"/>
(MAXIMUM)	(58)	(58)	(58)	(58)

SENSORY

KEY SENSORY POINTS

0 = absent
1 = impaired
2 = normal
NT = not testable



• Key Sensory Points

Any anal sensation (Yes/No)

PIN PRICK SCORE (max: 112)

LIGHT TOUCH SCORE (max: 112)

NEUROLOGICAL LEVEL

The most caudal segment with normal function

	R	L
SENSORY	<input type="checkbox"/>	<input type="checkbox"/>
MOTOR	<input type="checkbox"/>	<input type="checkbox"/>

COMPLETE OR INCOMPLETE?

Incomplete = Any sensory or motor function in S4-S5

ASIA IMPAIRMENT SCALE

ZONE OF PARTIAL PRESERVATION

Caudal extent of partially innervated segments

	R	L
SENSORY	<input type="checkbox"/>	<input type="checkbox"/>
MOTOR	<input type="checkbox"/>	<input type="checkbox"/>

ASIA IMPAIRMENT SCALE

- A = Complete:** No motor or sensory function is preserved in the sacral segments S4-S5.
- B = Incomplete:** Sensory but not motor function is preserved below the neurological level and includes the sacral segments S4-S5.
- C = Incomplete:** Motor function is preserved below the neurological level, and more than half of key muscles below the neurological level have a muscle grade less than 3.
- D = Incomplete:** Motor function is preserved below the neurological level, and at least half of key muscles below the neurological level have a muscle grade of 3 or more.
- E = Normal:** motor and sensory function are normal

CLINICAL SYNDROMES

- Central Cord
- Brown-Sequard
- Anterior Cord
- Conus Medullaris
- Cauda Equina

Level of Cord Injury determines level of function



Prognosis for Recovery of spinal Cord Injury:

Poor prognosis for recovery if:

- pt arrives in shock
- pt cannot breath
- pt has a complete injury

Assessment



Severity of neurologic deficit

Complete

Flaccid paralysis below level of injury.

May involve diaphragm if injury above C5.

Sympathetic tone loss if fracture above T6.

Incomplete

? Any sensation.

? Sacral sparing.

Assessment

Severity of neurologic deficit


Incomplete

Central cord syndrome:

- # Characterized by disproportionately (UL>LL).
- # Mechanism: hyper-extension.
- # Occur with or without fractures.
- # Recovery: 50% regaining function.
- # Prognosis is fair.



Assessment



Severity of neurologic deficit

Incomplete

Anterior cord syndrome:

Characterized by loss of corticospinal and spinothalamic tract with preserved posterior column.

Mechanism: ischemia or infarction to spinal cord..

Common injury.

Recovery: 10%.

Prognosis is good if progressive recovery within 24hrs, absent SS after 24hrs protends a poor outcome.

Assessment

Severity of neurologic deficit

Incomplete

Brown-Sequard syndrome:

Characterized by hemicord injury with ipsilateral paralysis, loss of proprioception and light touch, and contralateral temperature and sharp pain loss.

Prognosis is good, with over 90% regaining of bowel and bladder function and ambulatory capacity.

Assessment



Severity of neurologic deficit

Incomplete

Conus Medullaris syndrome:

Seen in T12-L1 injuries.

Loss of voluntary bowel and bladder control with preserved lumbar root function.

Uncommon as pure lesion (mixed conus-cauda).

Assessment



Severity of neurologic deficit

Incomplete

Cauda Equina syndrome:

Saddle anesthesia, urinary retention and stool incontinence.

Usually due to large central disc herniation rather than fracture.

Nerve root deficit: LMN



∞ Spinal Shock

- ∞ Transient loss of spinal reflexes.
- ∞ Lasts 24-72 hours.

∞ Neurogenic shock

- ∞ Reduced tissue perfusion due to loss of sympathetic outflow and un-opposed vagal tone.
- ∞ Peripheral vasodilatation (hypotension and bradycardia).
- ∞ Rx: fluid resuscitation and vasopressors.

Imaging



- ❧ X-rays:
 - ❧ Cervical: 3 views.
 - ❧ AP, lateral and open mouth.
 - ❧ Thoraco-lumbar: 2 views.
 - ❧ AP & lateral.
 - ❧ Flexion-Extension views.
- ❧ CT: best for bony anatomy.
- ❧ MRI: best to evaluate soft tissue.

Management of Spinal Injuries



∞ Depends on:

∞ Level of injury.

∞ Degree and morphology of injury: **STABILITY**

∞ Presence of neurologic deficit.

∞ Other factors.



∞ Some general rules:

∞ *Stable* injuries are usually treated conservatively.

∞ *Unstable* injuries usually require surgery.

∞ Neurologic compression requires decompression.

Specific Injuries



Cervical spine fractures



- ❧ Descriptive: depends on mechanism of injury.
 - ❧ Flexion/extension.
 - ❧ Compression/distraction.
 - ❧ Shear.

- ❧ Presence of subluxation/dislocation

- ❧ SCI:
 - ❧ high fracture results in quadriplegia.
 - ❧ Low fracture results in paraplegia.

Cervical spine fractures



∞ SCI:

- ∞ high fracture results in quadriplegia.
- ∞ Low fracture results in paraplegia.



Cervical spine fractures





Thoraco-Lumbar fractures




- ❧ Spinal cord terminates at L1/2 disc in adult
 - ❧ L2/3 in a child
- ❧ 50% of injuries occur at Thoraco-lumbar junction.
- ❧ Common fractures:
 - ❧ Wedge fracture (flexion/compression).
 - ❧ Burst (compression).
 - ❧ Chance (flexion/distraction).

Wedge fracture



Wedge fracture



Burst fracture




Burst Fractures - characteristics

1. **Retropulsion** of posterosuperior vertebral body fragment
 - Wedge compression may **bulge** posterior cortex but not posteriorly displace
2. Sagittal fracture of vertebral body (90%)
3. Sagittal posterior element fracture (85%)
4. Widening of interpedicular distance (80%)





Burst fracture




Burst fracture



Chance fracture



Chance fracture



Chance fracture





Figure 1. Lap Belt Ecchymosis





Image courtesy of Dr. Antonio Muñoz.



© Division of Pediatric Surgery - Brown Medical School

Fracture dislocation



Pathologic fractures



- ❧ Low-energy fractures.
- ❧ Usually due to infection or tumour.
- ❧ Osteoporotic is common.
- ❧ X-rays: “winking owl” sign for infection or tumour.

Pathologic fractures




Cauda Equina Syndrome




- ❧ A surgical emergency.
- ❧ Requires full neurologic examination *including rectal examination for anal tone.*
- ❧ Investigations: X-rays initially, but *MRI is mandatory as X-rays are usually unremarkable.*
- ❧ Treatment: Emergency decompression-usually discectomy and wide laminectomy within 24 hours.

Cauda Equina Causes


- ❧ Central disc prolapse.
- ❧ Burst fractures of lumbar spine.
- ❧ Penetrating injuries such as stab wounds or bullets.
- ❧ Epidural hematoma from spinal anesthesia, or post surgery(rare).
- ❧ Tumours compressing the lower spinal nerve roots.
- ❧ Spinal Stenosis.




Disc hernia



Tumor




Burst fracture



Bullet to cauda

Cauda Equina Syndrome



Questions

